

Two new species of Chrysomelobia (Acari: Heterostigmata: Podapolipidae) parasitic on Gonioctena rubripennis Baly (Coleoptera: Chrysomelidae; Chrysomelinae) in Japan

Authors: Husband, Robert W., Kurosa, Kazuyoshi, and Seeman, Owen D.

Source: Systematic and Applied Acarology, 21(11) : 1450-1462

Published By: Systematic and Applied Acarology Society

URL: <https://doi.org/10.11158/saa.21.11.2>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Two new species of *Chrysomelobia* (Acari: Heterostigmata: Podapolipidae) parasitic on *Gonioctena rubripennis* Baly (Coleoptera: Chrysomelidae; Chrysomelinae) in Japan

ROBERT W. HUSBAND^{1*}, KAZUYOSHI KUROSA² & OWEN D. SEEMAN³

¹Biology Department, Adrian College, Adrian, MI 49221 USA. E-mail: husbandadrian@aol.com

²Nishi-Ikebukuro 5-21-15, Tokyo 171-0021, Japan. E-mail: CQW35713@nifty.com

³Queensland Museum, P.O. Box 3300, South Brisbane, QLD, 4101, Australia. E-mail: owen.seeman@qm.qld.gov.au

*Corresponding author

Abstract

Chrysomelobia matsuzawai sp. nov. and *Chrysomelobia nipponica* sp. nov. (Acari: Podapolipidae) are described from *Gonioctena rubripennis* Baly (Coleoptera: Chrysomelidae) collected in Japan. This is the first record of the genus *Chrysomelobia* Regenfuss, 1968 in Asia. Adult females of the type species for *Chrysomelobia*, *Chrysomelobia mahunkai* Regenfuss, 1968, were recollected from female specimens infesting a *Gonioctena* sp. in Germany. An updated key to all species of *Chrysomelobia* is provided.

Key words: insect parasites, *Chrysomelobia*, key, Acari, Japan, Podapolipidae

Introduction

Mites in the family Podapolipidae are common parasites of a number of families of Coleoptera, and less commonly on Blattodea and Orthoptera (e.g., Regenfuss 1968; Husband 1990; Husband & OConnor 2003). The family Podapolipidae is represented by one species each on Heteroptera (Kurosa & Husband 1994) and Hymenoptera (Husband & Sinha 1970) and has not been found on Diptera, Lepidoptera, Odonata or any aquatic insects. In most instances, each family of parasitized Coleoptera will have genera of Podapolipidae that are associated with that family only. A conspicuous exception is the genus *Podapolipus* Rovelli & Grassi, 1888 which is found on beetles in four families as well as on insects in the orders Blattodea and Orthoptera (e.g., Husband 1986). Previous reports of Podapolipidae on Chrysomelidae have been recorded as species in the genus *Chrysomelobia* Regenfuss, 1968 (*Parobia* Seeman & Nahrung, 2003) (Regenfuss 1968; Eickwort 1975; Drummond *et al.* 1984; Fain 1987; Haitlinger 1989; Houck 1992; Moraes *et al.* 1999; Husband & Moraes 1999; Seeman & Nahrung 2003, 2005, 2013; Husband & OConnor 2004; Seeman 2008). The single record of a podapolipid mite that is not a *Chrysomelobia* but is a parasite of a chrysomelid beetle is *Cassidopohpus physonotae* Husband & OConnor, 2014, a parasite of *Physonota alutacea* Beheman (Husband & OConnor 2014). Twenty-one species of *Chrysomelobia* have been reported from Australia (14), Africa (2), Europe (1), and the Americas (4) on leaf beetles belonging to the subfamily Chrysomelinae (Seeman & Nahrung 2013). The record of a single female specimen of *C. donati* Haitlinger, 1989 from a cercopid hemipteran is considered accidental. *Chrysomelobia nipponica* sp. nov. and *Chrysomelobia matsuzawai* sp. nov. are the first *Chrysomelobia* species described from Asia.

Materials and methods

Examination of chrysomelid beetles representing primarily the subfamily Chrysomelinae, by Kazuyoshi Kurosa over a number of years, yielded mites belonging to the genus *Chrysomelobia* (Podapolipidae). Mites were removed from the abdominal tergites and under the elytra. Mites mounted on slides in Hoyer's mounting medium were placed on a heated drying tray for five days and ringed with red insulating varnish.

Measurements were taken with a Zeiss compound phase contrast microscope with a stage micrometer. Measurements are given in micrometers (µm). Alveolar vestiges of setae are designated as v. Microsetae, designated as m, are no longer than the diameter of their setal alveoli. Other terminology is based on Lindquist (1986).

The holotypes are deposited in the National Museum of Nature and Science, Tsukuba, 306-0005, Japan (NSMT). Paratypes of males, larvae and females are housed with the holotypes excepting some female, male and larval paratypes that are placed in the following museums: the A.J. Cook Arthropod Research Collection, Michigan State University, East Lansing, Michigan (CARC); The Acarology Laboratory, Museum of Biodiversity, The Ohio State University, Columbus, Ohio (OSAL); United States National Museum of Natural History, Washington, D.C. (NMNH) (mite collection housed in the USDA Systematic Entomology Laboratory, Beltsville, Maryland); Queensland Museum, South Brisbane, Australia (QMBA); Tarbiat Modares University, Tehran, Iran (TMUI); University of Michigan Museum of Zoology, Ann Arbor, Michigan (UMMZ); Tyumen State University, Tyumen, Russia (TSUR) and Zoological Museum, University of Hamburg, Hamburg, Germany (ZMH).

Description of new species

Chrysomelobia matsuzawai Husband, Kurosa & Seeman sp. nov.

(Figs. 1–5)

Diagnosis. All life stages. Tibia I with seta *k*, tarsus I with seven setae and one solenidion, setae *tc'* and *tc''* with blunt tips. Adult female: trachea shorter than setae v_1 , setae v_1 slender, setae sc_2 bulbous, setae c_1 bulbous, setae c_2 long, slender, setae *e* shorter than v_1 . Coxal setae *la*, *2a* and *3b* bulbous, apodemes I, II meeting sternal apodeme. Leg I with one claw. Femur II with minute setae *d* and conspicuous *l'*. Tibia IV with a pair of long setae, tarsus IV with a single long seta. Adult male: shield C, D, EF with 4 pairs of setae, c_1 , *d*, *e* minute, c_2 developed, 5 long; setae c_1 posterior to plane of setae c_2 . Plate C, D, EF with row of setae *d* slightly anterior to row of setae *e*. Genital capsule posterodorsal, shield C, D, EF with broadly concave posterior margin, setae ps_1 not evident; tibiae I, II, III with spine-like setae, femur II with minute setae *d* and longer *l'*, leg IV enlarged basally, convex lateral margin, about 2/3 length of leg III, tibia III setae v'' nearly 1/2 width of idiosoma, tibia IV setae v' , v'' shorter than seta *d*. Legs I, II, III with two claws. Three tarsus IV setae plus a curved claw. Larva: dorsal gnathosomal setae nearly 1/2 length of dorsal gnathosomal setae in adult females.


Description

Female (Figs. 1, 2, n=18)

Gnathosoma. Length 60–70, width 54–63 (Table 1). Cheliceral stylets 50–57 (one 46). Pharynx width 10–12. Setae *ch* 17–27, *su* 5–10. Palps longer than wide, three segmented, *su-su* 18–19.

Idiosoma. Length 258–319, width 208–240, setae v_1 24–30, positioned on narrowed anterolateral margin of prodorsal shield and immediately posterior to stigmata, v_2 v. Setae sc_2 bulbous, length 10–

12, width 7–9. Idiosomal plate lengths: PD 98, C 70, D 50–58, EF 45–52; widths PD 208, C 210–220, D 190, EF 103–123, setae c_1 bulbous, length 8–9, width 4–6, c_2 90–127, d 7–8 (one 12), e 7–10, h_1 15–25, h_1-h_1 53–56. Cupule *ia* anterolateral to setae *d*, cupule *im* anterolateral to setae *e*. Stigmata at anterolateral margin of prodorsal shield. Trachea length 25–28, width 5, branching not evident. Distance between setae v_1-v_1 70–82, sc_2-sc_2 43–54, c_1-c_1 86–93, c_1-c_2 31–37, v_1-sc_2 38–55, v_2-sc_2 8–10. Venter with apodemes II meeting sternal apodeme. Coxal setae *la* bulbous 12 long, 7 wide; *2a* bulbous, 10 long, 8 wide, *2b* v, *3a* 5–8, *3b* bulbous, 10 long, 6 wide; *4b* 6. Distance between setae *la-la* 41, *2a-2a* 72, *3a-3a* 82, *3b-3b* 129.



FIGURES 1–2. *Chrysolobobia matsuzawai* Husband, Kurosa & Seeman **sp. nov.**, adult female, 1. dorsal, 2. ventral.

Legs. Femur I setae *l'* thick 18–20, *d* m, v'' 10–16, tibia I with setae *l'* m, *d* 65–75, ϕ 11–14, *k* thin 12–13. Tarsus I setae *tc'*, *tc''* eupathidial (blunt), *tc'* 22–28, *tc''* 20–30, solenidion ω 9–11. Femur II setae *l'* 9–10, *d* 2. Tibia II *l'* 11–13, *d* 20–22, v' 28, v'' 43. Tarsus II setae *pl'* 9–14, *tc'* 43–46, *pl''* 53, *u'* 10–12, *pv''* 8–9. Tibia III setae *l'* 7–10, *d* 42, v' 18, v'' 30. Tarsus III setae *tc'* 48, *pl'* 11–14, *pl''* 50, *u'* 8–12, *pv''* 5–7. Femur and genu IV fused. Tibia IV setae v' , v'' and tarsi IV *tc'* exceed 200. Tibia and tarsus IV separate. Setation for femur, genu, tibia, tarsus I, II, III, IV: 30-6(+1)-7(+1), 2-0-4-5, 0-0-4-5, 0-0-2-1.


Male (Figs. 3, 4, n=1)

Gnathosoma. Length 50, width 51. Cheliceral stylets 33, pharynx width 10, setae *ch* 9, *su* 6, *su-su* 19, palp length 15.

Idiosoma. Length 250, width 207, setae v_1 2, v_2 m, setae sc_2 3, c_1 m, c_2 5, *d* m, distance between setae v_1-v_1 32, v_2-v_2 30, sc_2-sc_2 66, c_1-c_1 60, c_1-c_2 41, $d-d$ 32, $e-e$ 78. Genital capsule posterodorsal, length 35, width 50, two internal lobes interpreted as setae ps_2 length 10, aedeagus small. Venter with apodemes II almost reaching sternal apodeme. Coxal setae minute.

TABLE 1. Maximum measurements in micrometers (μm) for *Chrysomelobia mahunkai* (*mah*), *C. nipponica* *sp. nov.* (*nip*), *C. matsuzawai* *sp. nov.* (*mat*), *C. gimlii* (*gim*), *C. pagurus* (*pag*), *C. orthagoriscus* (*ort*), *C. captivus* (*cap*), *C. alleni* (*all*), *C. lipsettae* (*lip*). Males and larvae of *C. mahunkai* have not been reported.

	<i>mah</i>	<i>nip</i>	<i>mat</i>	<i>gim</i>	<i>pag</i>	<i>ort</i>	<i>cap</i>	<i>all</i>	<i>lip</i>
FEMALE									
Idiosoma length	300	300	505	433	370	505	430	320	433
Idiosoma width	289	289	289	300	318	388	279	230	300
Gnathosoma width	70	76	63	82	85	79	69	50	83
Cheliceral stylets	74	66	55	44	60	60	55	47	67
Gnathosomal setae									
<i>ch</i>	46	41	27	50	40	40	44	26	63
<i>su</i>	16	17	9	21	19	23	21	17	28
Idiosomal setae									
<i>v</i> ₁	34	35	30	29	24	18	15	8	61
<i>sc</i> ₂ bulbous (b)	53	22	12b	10b	12b	7b	12	12b	143
<i>c</i> ₁ bulbous (b)	7	10	9b	9b	10b	6b	12	80	10
<i>c</i> ₂ bulbous (b)	142	140	127	155	145	135	136	11b	19
<i>e</i>	10	32	10	24	24	19	27	77	19
<i>h</i>	19	25	25	28	18	31	22	20	18
Coxal setae									
1 <i>a</i> bulbous (b)	20	12	12b	7b	10b	7b	m	12b	20
2 <i>a</i> bulbous (b)	6b	6b	10b	8b	10b	7b	7.5b	11b	24
3 <i>b</i> bulbous (b)	6b	5b	10b	6b	9b	7b	7b	12	12
MALE									
Idiosoma length	–	200	250	290	223	255	228	215	260
Idiosoma width	–	187	207	235	215	238	187	155	235
Gnathosoma width	–	60	51	76	65	72	57	43	70
Cheliceral stylets	–	41	33	50	48	51	48	34	54
Gnathosomal setae									
<i>ch</i>	–	19	9	15	17	16	16	9	22
<i>su</i>	–	4	6	17	20	17	14	9	17
Dorsal plate setae									
<i>v</i> ₁	–	m	2	m	2	m	m	m	2
<i>sc</i> ₂	–	4	3	v	2	m	m	m	4
<i>c</i> ₁	–	2	m	m	2	m	m	m	2
<i>c</i> ₂	–	10	5	13	8	9	10	m	6
Genital capsule length	–	23	43	43	37	37	32	32	42
Genital capsule width	–	23	48	46	47	52	43	60	60
Tibia III, setae <i>v</i> ''	–	16	140	160	38	220	120	120	130
LARVA									
Idiosomal length	–	300	310	275	330	265	231	223	158
Idiosomal width	–	230	212	198	268	210	193	187	117
Gnathosomal width	–	32	43	77	132	112	88	81	53
Cheliceral stylets	–	50	53	110	267	202	180	89	66
Gnathosomal setae									
<i>ch</i>	–	12	18	50	95	62	58	39	35
<i>su</i>	–	7	7	17	58	46	7	10	19
Dorsal plate setae									
<i>v</i> ₁	–	m	m	63	39	15	22	3	32
<i>sc</i> ₂	–	4	v	140	10	137	175	122	83
<i>e</i>	–	m	10	100	33	22	15	10	23
Seta <i>h</i> ₁	–	-	55	240	396	283	230	407	145
Seta <i>h</i> ₂	–	-	-	5	11	6	10	3	m
Seta tibia I <i>d</i>	–	17	7	43	44	38	36	43	67
Seta tibia III <i>v</i> ''	–	16	17	150	38	130	120	70	190



FIGURES 3–4. *Chrysomelobia matsuzawai* Husband, Kurosa & Seeman **sp. nov.**, male, 3. dorsal, 4. ventral.

Legs. Femur I setae l' 6, v'' 8, d m, femur II setae l' 10, d m, no femora III, IV setae. No genua I, II, III, IV setae. Tibia I solenidion ϕ 11, slender adjacent seta k 10. Tibia I setae v' spine-like, tibiae II, III setae l' spine-like 10, tibia I, II, III setae d 22, 18, 15. Tarsi I setae pl' 15, tc' 23, tc'' 30, solenidion ω 10, setae pl'' 22, s 12, pv' 3, pv'' 8. Tibia III setae v'' 140. Ambulacra I, II, III with two stout claws. Setation for femur, genu, tibia, tarsus of legs I, II, III, IV: 3-0-6 (+1)-7(+1), 2-0-4-5, 0-0-4-5, 0-0-3-3 + claw. Thickness of fused femur and genu IV 33.

Larval female (Fig. 5, $n=5$ exoskeletons containing adult females)

Gnathosoma. Length 43–52, width 37–48. Cheliceral stylets 45–53, pharynx width 8–10. Setae ch 9, su 9, $su-su$ 15.

Idiosoma. Length 270–370, width 195–265, setae v_1 m, v_2 v, sc_2 3, c_1 m, c_2 3, h_1 55, distance setae v_1-v_1 30, v_2-v_2 40, sc_2-sc_2 63, v_2-sc_2 20. Distance between setae c_1-c_2 32, h_1-h_1 8.

Legs. Femur I setae l' 5, d m, v'' 6, no genua I, II, III, IV setae. Tibia I solenidion ϕ 6, k 2, v' 5, v'' 5, d 5. Tarsus I setae tc' 12, tc'' 14, solenidion ω 5, setae pl'' 6. Femur II setae l' 6, d 3. Tibia II setae v'' 26, tibia III setae v'' 70. Ambulacra I, II, III with two stout claws.

Etymology. The species is named for Dr. Haruo Matsuzawa, specialist in Chrysomelidae, who provided many potential host beetles that yielded *Chrysomelobia* mites for this study. The species name is a noun in the genitive case.

Type material. All specimens from *Gonioctena rubripennis* Baly (Coleoptera: Chrysomelidae). *Holotype*: adult female (Kurosa Collection No. 3321–3(3/7), Shiromana, Okutama, Tokyo, Japan, 3 May 1980, coll. K. Kurosa, deposited with the type host in the National Museum of Nature and Science, Tsukuba, Japan (NSMT). *Paratypes*: 5 females, 1 male, same data as holotype (KCN 3321-1 to 3321-8); 3 females, Mineoka, Kamogawa City, Chiba Pref., Japan, 4 June 1978, coll. J. Okuma; 2 females (1 slide), Mt. Odamiyama, Oda-cho, Ehime Pref., Japan, coll. E. Yamamoto; 5 females inside of exoskeletons of larval females, Kamafuga Dam, Miyaga Pref., Japan, 27 April 1995, coll. unknown; 2 females, Bizen-shi, Okayama Pref., Japan, 7–9 1989, coll. unknown. One female paratype each is deposited at CARC, OSAL, NMNH, QMBA, TMUI, TNAU, UMMZ

and ZMH. The balance of paratypes is deposited with the holotype (NSMT). The balance of type hosts is deposited in UMMZ.




FIGURE 5. *Chrysomelobia matsuzawai* Husband, Kurosa & Seeman **sp. nov.**, larval female, dorsal, proterosoma.

Differential diagnosis. The new species appears closely related to *C. gimlii* (Seeman & Nahrung, 2005), but differs by having females with the alveolar vestige of seta v_2 situated close to seta sc_2 and setae $tc'-tc''$ on tarsus I eupathidial (alveolar vestige of seta v_2 midway between setae v_1 and sc_2 and setae $tc'-tc''$ with tapering tips in *C. gimlii*); and by having males without alveolar vestiges of setae sc_1 and lacking extremely long setae on tibia III (alveolar vestiges of setae sc_1 present and tarsus III with a very long attenuate seta in *C. gimlii*).

Remarks. The six species of *Chrysomelobia* from the Western Hemisphere (4 spp.) and Africa (2 spp.) all have setae on genera I–II, and of those six species, only *C. donati* lacks setae on genus III. These setal losses place *C. matsuzawai* **sp. nov.** within the radiation of 14 Australian species, plus the European species *C. mahunkai* Regenfuss 1968, that all lack setae on genera I–III. The distinctive bulbous setae present in several species of *Chrysomelobia* are expressed variously and help define species groups, as indicated in Seeman (2008). The bulbous setae in *C. matsuzawai* **sp. nov.** are sc_2 , c_1 , la , $2a$ and $3b$, which is the same as mites in the *gimlii* species group (*C. gimlii*, *C. orthagoriscus* Seeman, 2008, *C. pagurus* Seeman, 2008). Thus, *C. matsuzawai* **sp. nov.** is similar to these species, but differs from the other species of the *gimlii* species group in the following features. In female *C. matsuzawai* **sp. nov.**, the vestige of seta v_2 is situated close to seta sc_2 and setae $tc'-tc''$ on tarsus I are eupathidial, i.e., blunt-tipped. In species of the *gimlii* species group, seta v_2 is in a more typical position midway between setae v_1 and sc_2 and setae $tc'-tc''$ are not eupathidial, having tapering tips.

Chrysomelobia matsuzawai **sp. nov.** also differs from all other Australian species, excepting *C. lipsettae* Seeman, 2008, by having broad tracheae (width 5). The thin trachea that do not anastomose may be a synapomorphy for species of *Chrysomelobia* that infest eucalypt-feeding Paropsini; the host of *C. lipsettae* feeds on *Acacia* (Fabaceae), and *C. lipsettae* was hypothesized by Seeman (2008) to be a species intermediate between the Australian (+*C. mahunkai*) and the American and African species of *Chrysomelobia*.

***Chrysmelobia nipponica* Husband, Kurosa & Seeman sp. nov.**

(Figs. 6–10)

Diagnosis. All life stages. Tibia I with seta *k*, tarsus I with seven setae and one solenidion, setae *tc'* and *tc''* with slender tips. Adult female: tracheae broad and long, anastomosing distally, setae *c*₁ slender, setae *c*₂ long, setae *e* as long as setae *v*₁. Coxal setae *la*, *4a* slender, *2a* and *3b* bulbous, femora I, II with seta *l'* and minute setae *d*. Tibia IV with a pair of long setae, tarsus IV with a single long seta. Adult male: setae *v*₁, *v*₂, *d*, *e* minute, setae *sc*₂, *c*₁, *c*₂ short, two times diameter of setal acetabulum. Genital capsule posterodorsal, wider than long, setae *h*₁, *h*₂ minute, setae *ps*₁ minute, setae *ps*₂ internal, lobular. Femur I setae *l'* 10, *d* m, *v''* 10, femur II setae *l'* 4, *d* m. Tibiae I, II, III without spine-like setae. Tibia III setae *l''* shorter than tibia III setae *d*. Legs I, II, III with two claws. Larva: gnathosomal setae *su* near 1/2 length setae *ch*. Setae *v*₁, *v*₂, *c*₁ minute, setae *c*₂, *sc*₂ short, two times diameter of setal acetabulum.

Description

Female (Figs. 6, 7, n=16)

Gnathosoma. Length 60–67, width 63–76 (Table 1). Cheliceral stylets 60–66. Pharynx width 12–13. Setae *ch* 37–41, *su* 12–17. Distance between setae *su*–*su* 19–24. Palps longer than wide, two segmented.

Idiosoma. Length 275–300, width 222–289, with prodorsal shield narrow anteriorly, setae *v*₁ 25–35, *v*₂ *v*, *sc*₂ 19–22, *c*₁ 7–10, *c*₂ 120–140, *d* 8–10, *e* 28–32, *h*₁ 20–25. Idiosomal plate lengths: prodorsal plate 90–98, C 70–72, D 58–60, EF 40–60; widths PD 178–180, C 223–228, D 118–130, EF 112–130, setae *c*₁ in line with *c*₂. Stigmata at anterolateral prodorsal shield. Length of broad trachea leading from stigmata near 80 and anastomosing distally. Stigmata–stigmata 63–70, setae *v*₁–*v*₁ 70–76, *v*₂–*v*₂ 55–59, *sc*₂–*sc*₂ 112–114, *c*₁–*c*₁ 55–58, *d*–*d* 50–52, *e*–*e* 91–100, *h*₁–*h*₁ 56–62. Venter with apodemes II not meeting sternal apodeme. Coxal setae *la* slender 8–12, *2a* bulbous 5–6, *3a* slender 10–16, *3b* bulbous 4–5, *4b* 10–17. Distance between *la*–*la* 62–66, *2a*–*2a* 106–110, *3a*–*3a* 50–53, *3b*–*3b* 104–112.

Legs. Femur I setae *l'* thick 19–22, *d* m, *v''* 20–29, tibia I *d* 82–90, *φ* 12–13, *k* 11–13. Tarsus I setae *tc'* 35–42, *tc''* 30–39, *ω* 10–12. Ambulacrum I with one claw. Femur II setae *l'* 10–13, *d* m, tibia II *l'* 11–20, *d* 39–48, *v'* 22–30, *v''* 34–36. Tarsus II setae *pl'* 13–14, *tc'* 45–53, *tc''* 30–37, *u'* 9–12, *pv''* 6–12. Tibia III setae *l'* 8–12, *d* 34–37, *v'* 20–30, *v''* 34–37, tarsus III setae *pl'* 14–19, *tc'* 39–48, *tc''* 30–40, *u'* 9–10, *pv''* 7–10. Tibia IV setae *v'*, *v''* and tarsus IV setae *tc'* exceed 250. Setation for femur, genu, tibia, tarsus I, II, III, IV: 3-0-6(+1)-7(+1), 2-0-4-5, 0-0-4-5, 0-0-2-1.

Male (Figs. 8, 9, n=1)

Gnathosoma. Length 50, width 60. Cheliceral stylets 41, pharynx width 10, setae *ch* 19, *su* 7, distance *su*–*su* 21, palp length 12.


Idiosoma. Length 300, width 230, setae *v*₁ m, *v*₂ m, *sc*₂ 4, *c*₁ 3, *c*₂ 10, *d*, *e*, *f* m. Distance between setae *v*₁–*v*₁ 32, *v*₁–*sc*₂ 49, *sc*₂–*sc*₂ 75, *c*₁–*c*₁ 33, *c*₂–*c*₂ 114, *c*₁–*c*₂, *d*–*d* 18, *e*–*e* 63. Genital capsule posterodorsal, length 35, width 50.

Legs. Femur I setae *l'*, *d* m, *v''* 10, femur II setae *l'* 4, *d* m. No genua I, II, III, IV setae. Tibia I setae *l'* 3, *d* 30, *φ* 10, *k* 10, *v*₁ thick 6, *v''* 17, *l''* 3. Tarsus I setae *tc'* 27, *tc''* 31, *ω* 8, *pl'* 13, *pv'* 3, *s* 8, *pv''* 5, *pl''* 13. Femur II setae *l'* 4, *d* m, tibia II setae *l'* 6, *d* 19, *v*₁ 18, *v''* 19, tarsus II setae *pl'* 8, *tc'* 34, *pl''* 25, *pv'* 2, *u'* 6, *pv''* 6. Tibia III seta *l'* 16. Thickness of mid femur IV 20. Ambulacra I, II, III with two claws. Setation for femur, genu, tibia, tarsus I, II, III: 3-0-6(+1)-7(+1), 2-0-4-5, 0-0-4-5, 0-0-2-2+ claw.

Larval female (Figure 10, n=1, exoskeleton very pale)

Gnathosoma. Length 49, width 52. Cheliceral stylets 50, pharynx width 11. Setae *ch* 12, *su* 7, *su-su* 9.

Idiosoma. Length 300, width 230. Setae v_1 m, v_2 m, sc_2 4, c_1 m, c_2 4, h_1 m. Distance v_1-v_1 29, v_2-v_2 34, v_2-sc_2 25, sc_2-sc_2 58, c_1-c_2 20.




FIGURES 6–7. *Chrysomelobia nipponica* Husband, Kurosa & Seeman **sp. nov.**, adult female, 6. dorsal, 7. ventral.

Legs. Femur I seta l' 4, d m, v'' 10. Tibia I l' m, d 28, ϕ 10, k 8, v' 4, v'' 12, l'' m. Tarsus I tc' 22, tc'' 22, ω 9, pl' 9, pv' 3, s 5, pv'' 3, pl'' 8. Femur II seta l' 5, d m, Tibia II l' 3, d 7, v' 8, v'' 14. Tarsus II tc' 16, pl'' 23, u' 8, pv'' 4. Tibia III l' 2, v' 12, v'' 17. Tarsus III pl' 10, pl'' 20, tc' 12, u' 5, pv'' 3. Setaion for femur, genu, tibia, tarsus I, II, III: 3-0-6(+1)-7(+1), 2-0-4-5, 0-0-4-5.


Etymology. The specific name *nipponica* is an adjective derived from the country of origin, Japan (Nippon).

Type material. All specimens from *Gonioctena rubripennis* Baly (Coleoptera: Chrysomelidae). *Holotype*: adult female (RWH25 VIII 2015-2), Katsuura, Chiba Pref., Japan, 22–24 IV 1989, coll. K. Kurosa, deposited with the type host in the National Museum of Nature and Science, Tsukuba, Japan. *Paratypes*: 1 male, 12 females, 1 larva, same data as holotype; 2 females, Hongo-Cho, Aizu-Wakamatsu, Fukushima Pref., Japan, 26–27 V 1999, coll. unknown; 1 male, Minakami-machi, Gunma Pref., Japan, 1 IV 1999, coll. unknown; 1 female, Fujikawachi, Umemachi, Saiki-shi, Oita Pref., Japan, 3 VI 2012, coll. S. Sasaki; 1 female, Nano-shi, Yamagata Pref. Japan, 23 VI 1999, coll. unknown. 1 female paratype is deposited at each of the following CARC, OSAL, NMNH, QMBA, TMUI, TNAU, TSUR, UMMZ, ZMH. Balance of paratypes is deposited with the holotype (NSMT). Balance of type hosts deposited in UMMZ.



FIGURES 8–9. *Chrysomelobia nipponica* Husband, Kurosa & Seeman **sp. nov.**, male, 8. dorsal, 9. ventral.

Differential diagnosis. The new species appears closely related to *C. mahunkai*, but differs by having females with one ambulacral claw (two in all other *Chrysomelobia*) and seta *d* on femur II (absent in *C. nipponica* **sp. nov.**).



FIGURES 10–11. 10. *Chrysomelobia nipponica* Husband, Kurosa & Seeman **sp. nov.**, larval female. A, dorsal, proterosoma; B, dorsal, idiosomal plates; C, ventral, proterosoma; 11. *Chrysomelobia mahunkai* Regenfuss 1968, holotype, adult female, ventrodorsal.

Remarks. The host genus for both new species, *Gonioctena* (Coleoptera: Chrysomelidae), is also a host for the type species of *Chrysomelobia*, *C. mahunkai*. New adult female specimens of *C. mahunkai* were collected from an unspecified locality in Germany from *Gonioctena* sp. and the holotype female was also examined. An illustration of the holotype female loaned by Dr. Hieronymus Dastych of the University of Hamburg, Germany is provided (Fig. 11). Previously, the species was recorded from a single female collected from Tansey beetle *Chrysolina graminis* (L., 1758) (= *Chrysomela graminis*). Of the two new species, *C. matsuzawai* **sp. nov.** is not closely related to *C. mahunkai*, but in contrast, *C. nipponica* **sp. nov.** shares several similar character states with *C. mahunkai*. These similarities are the bulbous coxal setae *2b* and *3b*, dorsal setae and seta *la* unmodified, the female tibia IV with two setae and tarsus IV with one terminal seta, and the broad tracheae that anastomose distally. With the exception of the last character state, these states are also shared with some Australian species, particularly the *husbandi* species group. Female *C. mahunkai* and *C. nipponica* **sp. nov.** are distinct from the *husbandi* species group by having well-developed setae *la* (they are minute in the *husbandi* species group). Male *C. nipponica* **sp. nov.** differ from the *husbandi* species group by their large leg IV that bears a terminal claw (leg IV is diminutive and lacks a claw in the *husbandi* species group). Female *C. nipponica* **sp. nov.** are distinguished from *C. mahunkai* by the presence of two ambulacral claws on leg I in *C. mahunkai* (one in all other *Chrysomelobia*) and the presence of seta *d* on femur II in *C. nipponica* **sp. nov.** (absent in *C. mahunkai*). The male and larval stages for *C. mahunkai* remain unknown, so cannot be compared with *C. nipponica* **sp. nov.**, but we anticipate males and larvae of *C. matsuzawai* **sp. nov.** and *C. nipponica* to be similar. The divided plate C in the larva is absent in all Australian species but is present in *C. eickworti* Husband & OConnor, 2004, *C. labidomerae* Eickwort, 1975 and *C. peruviansis* Husband and Moraes, 1999 and may be present in *C. nipponica* **sp. nov.**

Key to species of *Chrysomelobia*

1. Female & male: at least 1 seta on genua I, II and IV and femur IV 2
 - Female & male: setae absent on genua I–IV and femur IV 7
- 2(1). Female: genu I with 3 setae; femur II with 1 seta; femur III without setae *C. donati* Haitlinger
 - Female & male: genu I with 4 setae; femur II with 3 setae; femur III with 2 setae 3
- 3(2). Female: coxal seta *4b* absent. Male: tarsus I without setae *ft'* and *ft''* *C. elytrosphaerae* Fain
 - Female: coxal seta *4b* present. Male: tarsus I with at least 1 *ft* seta 4
- 4(3). Female: genu IV with 2 setae (*v''* present). Male: with 4 prodorsal setae or vestiges of setae (*sc*₁ present); post-genital shield posterior to genital capsule expansive; fused telofemur-genu IV with 2 setae *C. eickworti* Husband & OConnor
 - Female: genu IV with 1 seta (*v''* absent). Male: with 3 prodorsal setae or vestiges of setae (*sc*₁ absent); post-genital shield posterior to genital shield elongate; fused telofemur-genu IV with 1 seta 5
- 5(4). Female: Cheliceral stylets 37–46. Male: idiosomal plates reticulate; tibia III, seta *v''* short (< 50). *C. oneili* Moraes, Husband & Lofego
 - Female: Cheliceral stylets > 50. Male: idiosomal plates smooth; tibia III, seta *v''* long (> 70) 6
- 6(5). Female: seta *h*₁ 30–40. Male: ventral gnathosomal setae 18–23; seta *sc*₂ close to margin of prodorsal shield; genu IV with 1 seta (*v'* present). *C. peruviansis* Husband & Moraes
 - Female: seta *h*₁ 17–19. Male: ventral gnathosomal setae 10–13; seta *sc*₂ well within margin of prodorsal shield; genu IV without setae (*v'* absent) *C. labidomerae* Eickwort
- 7(1). Female: tarsus IV with 2 or 3 long terminal setae. Male: seta *c*₂ minute 8
 - Female: tarsus IV with 1 long terminal seta. Male: seta *c*₂ developed, > 3 long 14
- 8(7). Female: seta *sc*₂, *c*₂, *la*, *2a* and *3b* slender 9
 - Female: seta *sc*₂, *c*₂, *la*, *2a* and *3b* bulbous 10
- 9(8). Female & male: seta *3a* absent. *C. vafer* Seeman
 - Female & male: seta *3a* present. *C. verecundus* Seeman

- 10(8). Female & male: tibia II lacking seta *l'*. Female: tibia and tarsus IV partially or completely fused. Male: dorsal shield C-D-E divided or with folds marking a weak division; tarsus IV, setae *u'* and *pv''* absent...
C. armstrongi Seeman
- Female & male: tibia II with seta *l'*. Female: tibia and tarsus IV separate. Male: dorsal shield C-D-E entire; tarsus IV, setae *u'* and *pv''* present..... 11
- 11(10). Female & male: tibia IV with 1 seta (*v''* present)..... *C. alipilus* (Seeman & Nahrung)
- Female & male: tibia IV without setae (*v''* absent)..... 12
- 12(11). Female: tarsus IV with 3 terminal setae; setae *d* and *e* < 40..... *C. nahrungae* Seeman
- Female: tarsus IV with 2 terminal setae; setae *d* and *e* > 45..... 13
- 13(12). Female: intercoxal setae further apart (*la-la* 37, *2a* 62–65). Male: tarsus IV with claw, *u'*, and 3 setae (minute seta *pv'* absent). Larva: setae *sc*₂ 27–31 and *c*₂ 25–29 long..... *C. alleni* Seeman & Nahrung
- Female: intercoxal setae closer together (*la-la* 26–31, *2a* 47–51). Male: tarsus IV with claw, *u'*, and 4 setae (minute seta *pv'* present). Larva: setae *sc*₂ 13–18 and *c*₂ 13–18 long... .. *C. aquariolus* Seeman
- 14(7). Female: coxal setae *2a* and *3b* slender... .. *C. lipsettae* Seeman
- Female: coxal setae *2a* and *3b* bulbous or minute... .. 15
- 15(14). Female: seta *la* slender; tracheae broad, anastomosing distally..... 16
- Female: seta *la* minute or bulbous; tracheae narrow, not anastomosing distally..... 17
- 16(15). Female: ambulacra with 2 claws; femur II without seta *d*... .. *C. mahunkai* Regenfuss
- Female: ambulacra with 1 claw; femur II with seta *d*... .. *C. nipponica* **sp. nov.**
- 17(15). Female: seta *sc*₂, *c*₁ and *la* bulbous. Male: tarsus IV with terminal claw, with 3–4 setae... .. 18
- Female: seta *sc*₂ and *c*₁ slender; seta *la* minute. Male: tarsus IV lacking terminal claw, with 1–2 setae... .. 21
- 18(17). Female: vestigial seta *v*₂ closely associated with seta *sc*₂; tarsus I setae *tc'-tc''* eupathidial (blunt-tipped). Male: plate C-D-EF with 5 pairs of setae; post-sternal apodeme well-developed... ..
C. matsuzawai **sp. nov.**
- Female: vestigial seta *v*₂ not closely associated with seta *sc*₂, about half way between setae *v'* and *sc*₂; tarsus I setae *tc'-tc''* not eupathidial (tips tapered). Male: plate C-D-EF with 4 pairs of setae; post-sternal apodeme developed or absent..... 19
- 19(18). Female: seta *sc*₂ and *c*₁ 5–7 long, 4–5 wide, with mediolateral projection 4–5 long (if broken then obvious stub present). Male: tibia II, seta *v''* 54–58, tarsus III, seta *tc'* 43–47. *C. orthagoriscus* Seeman
- Female: seta *sc*₂ 8–12 long, 5–6 wide; seta *c*₁ 7–10 long, 5.5–7 wide, mediolateral projection absent or a minute stub. Male: tibia II, seta *v''* either < 40 or > 80 long, tarsus III, seta *tc'* either < 35 or > 50 long..... 20
- 20(19). Female: distance between setae *v*₁–*sc*₂ 52–60, *v*₂–*sc*₂ 26–31. Male: seta *c*₂ 10–13; tibia II, seta *v''* 34–39; tarsus III, seta *tc'* 50–56... .. *C. gimlii* (Seeman & Nahrung)
- Female: distance between setae *v*₁–*sc*₂ 40–44, *v*₂–*sc*₂ 12–21. Male: seta *c*₂ 5–9; tibia II, seta *v''* 80–140; tarsus III, seta *tc'* 31–34... .. *C. pagurus* Seeman
- 21(17). Female: setae *2a* and *3b* bulbous... .. 22
- Female: no coxal setae bulbous... .. 23
- 22(21). Female: setae *2a* and *3b* 5–6 long, 3–4 wide; distance between *h*₁–*h*₁ 31–41. Male: tarsus II, seta *tc''* < 50... .. *C. husbandi* (Seeman & Nahrung)
- Female: setae *2a* and *3b* 6–7.5 long, 4.5–5 wide; distance between *h*₁–*h*₁ 22–29. Male: tarsus II, seta *tc''* > 80... .. *C. captivus* (Seeman & Nahrung)
- 23(21). Female & male: femur II without setae... .. *C. cubile* Seeman
- Female & male: femur II with minute seta... .. 24
- 24(23). Female: seta *v*₂ vestigial but distinct; setae *h*₁ length 33–43. *C. intrusus* Seeman & Nahrung
- Female: seta *v*₂ absent; setae *h*₁ length 15–19... .. *C. lawsoni* (Seeman & Nahrung)

Acknowledgements

We are thankful for assistance with obtaining potential host Chrysomelidae for Podapolipidae and advice by Barry O'Connor and Mark O'Brien of the University of Michigan Museum of Zoology, Ann Arbor, Michigan and for technical assistance from Adrian College librarians Richard Geyer, David Cruse and Noelle Keller. Our cordial thanks are also due to the following Japanese entomologists: Takeshi Matsuzawa, Jun Okuma, Shigeo Sasaki and Eiji Yamamoto.

References

- Drummond, F.A., Casagrande, R.A., Chauvin, R., Hsiao, T.H., Lashomb, J.H., Logan, P.A. & Atkinson, T.H. (1984) Distribution and new host records of a race of *Chrysomelobia labidomerae* Eickwort (Acari: Tarsonemina; Podapolipidae) attacking the Colorado potato beetle in Mexico. *International Journal of Acarology*, 10(3), 179–180.
<http://dx.doi.org/10.1080/01647958408683372>
- Eickwort, G.C. (1975) A new species of *Chrysomelobia* (Acari: Tarsonemina; Podapolipidae) from North America and the taxonomic position of the genus. *Canadian Entomologist*, 107, 613–626.
<http://dx.doi.org/10.4039/Ent107613-6>
- Fain, A. (1987) *Chrysomelobia elytrosphaerae* nov. spec. (Acari: Podapolipidae) parasite d'un Chrysomelidae neotropical. *Bulletin de l'Institut royal des Sciences naturelles de Belgique*, 57, 191–195.
- Haitlinger, R. (1989) New species *Chrysomelobia donati* and *Coccipolipus arturi* (Acari, Prostigmata, Podapolipidae) connected with insects from Cameroon and Sumatra. *Wiadomosci Parazytologiczne*, 35, 161–164.
- Houck, M.A. (1992) Morphological variation in an ectoparasite: partitioning ecological and evolutionary influences. In: Sorensen, J.T. & Footit, R. (Eds.), *Ordination in the Study of Morphology, Evolution and Systematics of Insects: Applications and Quantitative Genetic Rationals*. Amsterdam, Elsevier, pp. 277–308.
- Husband, R.W. (1986) New *Podapolipus* spp. (Acari: Podapolipidae) from Australia and Hawaii. *International Journal of Acarology*, 12, 187–209.
<http://dx.doi.org/10.1080/01647958608683465>
- Husband, R.W. (1990) New species of *Podapolipoides* (Acari: Podapolipidae), ectoparasites of grasshoppers (Orthoptera: Acrididae) in Australia and New Zealand, with keys to world species. *Annals of the Entomological Society of America*, 83, 371–393.
<http://dx.doi.org/10.1093/aesa/83.3.371>
- Husband R.W. & Moraes, G.J. de (1999) A new species of *Chrysomelobia* (Acari: Podapolipidae) from *Platyphora testudo* (Demay) (Coleoptera: Chrysomelidae) from Peru, with a key to known species of the genus. *International Journal of Acarology*, 25, 309–315.
<http://dx.doi.org/10.1080/01647959908684169>
- Husband, R.W. & OConnor, B.M. (2003) A new genus and species of mite (Acari: Tarsonemina: Podapolipidae), ectoparasite of the Peruvian cockroaches, *Blaberus parabolicus* (Walker) and *Eublaberus distantis* (Kirby) (Blattodea: Blaberidae). *International Journal of Acarology*, 29, 331–338.
<http://dx.doi.org/10.1080/01647950308684349>
- Husband, R.W. & OConnor, B.M. (2004) A new species of *Chrysomelobia* Regenfuss (Acari: Podapolipidae) from *Ceralces* sp. (Coleoptera: Chrysomelidae) from Tanzania, with a key to species of *Chrysomelobia*. *International Journal of Acarology*, 30, 17–23.
<http://dx.doi.org/10.1080/01647950408684363>
- Husband, R.W. & OConnor, B.M. (2014) A new species of Podapolipidae (Acari: Heterostigmata) parasitic on *Physonota alutacea* (Boheman) (Coleoptera: Chrysomelidae: Cassidinae) in Mexico and Central America. *Systematic & Applied Acarology*, 19, 435–446.
<http://dx.doi.org/10.11158/saa.19.4.7>
- Husband, R.W. & Sinha, R.N. (1970) A revision of the genus *Locustacarus* with a key to genera of the family Podapolipidae (Acarina). *Annals of the Entomological Society of America*, 63, 1152–1162.
<http://dx.doi.org/10.1093/aesa/63.4.1152>
- Kurosa, K. & Husband, R.W. (1994) *Cydnipolipus miyamotoi*, a new genus and species of podapolipid mite parasitic on *Aethus indicus* (Heteroptera: Cydnidae). *Journal of the Acarological Society of Japan*, 1, 21–32.
<http://dx.doi.org/10.2300/acari.3.21>
- Lindquist, E.E. (1986) The world genera of Tarsonemidae (Acari: Heterostigmata): a morphological phylogenetic and systematic revision, with a reclassification of family-group taxa in Heterostigmata. *Memoirs of the Entomological Society of Canada*, 136, 1–517.
<http://dx.doi.org/10.4039/entm118136fv>
- Moraes, G.J.de, Husband, R.W. & Lofego, A.C. (1999) A new species of *Chrysomelobia* (Acari: Podapolipidae) from Central America. *Systematic & Applied Acarology*, 4, 131–136.
<http://dx.doi.org/10.11158/saa.4.1.19>
- Rovelli, G. & Grassi, B. (1888) Di un singolare Acaride "*Podapolipus reconditus*, nobis". *Bullittino della*

- Societa Entomologica Italiana*, 20, 50–63.
- Regenfuss, H. (1968) Untersuchungen zur Morphologie, Systematic and Okologie der Podapolipidae (Acarina: Tarsonemini). *Zeitschrift für wissenschaftliche Zoologie*, 177, 183–282.
- Seeman, O.D. (2008) Systematics and phylogeny of *Chrysomelobia* species (Acari: Podapolipidae), sexually transmitted parasites of chrysomelid beetles. *Invertebrate Systematics*, 22, 55–84.
<http://dx.doi.org/10.1071/is06035>
- Seeman, O.D. & Nahrung, H.F. (2003) *Parobia husbandi* gen. and sp. nov. (Acari: Podapolipidae), subelytral parasites of paropsine beetles (Coleoptera: Chrysomelidae). *Australian Journal of Entomology*, 42, 334–342.
<http://dx.doi.org/10.1046/j.14406055.2003.00380.x>
- Seeman, O.D. & Nahrung, H.F. (2005) New *Parobia* (Acari: Podapolipidae) parasitic on pestiferous paropsine beetles (Coleoptera: Chrysomelidae) in eucalypt plantations. *Systematic & Applied Acarology*, 10, 111–135.
<http://dx.doi.org/10.11158/saa.10.1.12>
- Seeman, O.D. & Nahrung, H.F. (2013) Two new species of *Chrysomelobia* Regenfuss 1968 (Acariformes: Podapolipidae) from *Paropsis charybdis* (Coleoptera: Chrysomelidae). *Systematic Parasitology*, 86, 257–270.
<http://dx.doi.org/10.1007/s11230-013-9447-2>

Submitted: 16 May 2016; accepted by Anne Baker: 9 Sep. 2016; published: 17 Oct. 2016