

Discovery of the Male of *Canaea bibarra* (Lepidoptera: Thyrididae: Striglininae)

Authors: Huang, Lan-Lan, Owada, Mamoru, and Wang, Min

Source: Florida Entomologist, 98(1) : 200-201

Published By: Florida Entomological Society

URL: <https://doi.org/10.1653/024.098.0134>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Discovery of the male of *Canaea bibarra* (Lepidoptera: Thyrididae: Striglininae)

Lan-Lan Huang¹, Mamoru Owada² and Min Wang^{1,*}

Abstract

The male of *Canaea bibarra* (Chu & Wang, 1991) (Lepidoptera: Thyrididae: Striglininae) was discovered in Hainan, China for the first time. The male adult is very similar to female except for the paler ground color and monopectinate antenna. The male adults and the genitalia are described, with notes on their distribution and how they differ from related species.

Key Words: Lepidoptera; Thyrididae; Striglininae; *Canaea*; Hainan; China; Vietnam

Resumen

Machos de *Canaea bibarra* (Chu y Wang, 1991) (Lepidoptera: Thyrididae: Striglininae) fue descubierto en Hainan, China por primera vez. El macho adulto es muy similar a la hembra con la excepción del color de fondo más pálido y las antenas monopectinadas. Se describen los machos adultos y su genitalia, con notas sobre su distribución y cómo se diferencian de otras especies relacionadas.

Palabras Clave: Lepidoptera; Thyrididae; Striglininae; *Canaea*; Hainan; China; Vietnam

The genus *Canaea* Walker, 1864 (Lepidoptera: Thyrididae: Striglininae) was revised by Whalley (1971) on a worldwide basis, and 10 species with 10 subspecies were enumerated from West Malaysia to New Caledonia and the eastern coast of Australia. Chu & Wang (1991) described a new genus and species of striglinine moth, *Phthina bibarra* Chu & Wang, 1991, from Hainan on the basis of 3 female specimens. Robinson et al. (1994) illustrated *Canaea semitessellata* Walker, 1864 and *Canaea* sp., and added China (Hainan) to the distributional range of *Canaea*, but they did not refer to *Phthina bibarra*. In the checklist of Australian Lepidoptera, the genus *Phthina* Chu & Wang, 1991 was synonymized with *Canaea* by Shaffer & Nielsen (1996) without any comments.

In the course of a research project on the lepidopteran fauna of Hainan, we were able to collect males of *Canaea bibarra* (Chu & Wang 1991) in Jianfengling National Nature Reserve and Yinggeling Nature Reserve, Hainan Province, China. In addition, 2 males of this same species were collected in central Vietnam in joint research by the National Science Museum, Tokyo (NSMT) and the Institute of Ecology and Biological Resources, Hanoi (IEBR). In this paper, we describe features of the male for the first time.

Canaea bibarra (Chu & Wang, 1991) (Figs. 1–3)

Phthina bibarra Chu et Wang 1991: 340, 346, Fig. 20, pl. 2, Fig. 20; Zhu (Chu) & Wang 1996: 234–236, Fig. 186, pl. 15, Fig. 10.

SPECIMENS EXAMINED

CHINA, Hainan, Jianfengling National Nature Reserve, 900 m, 1 male, 23. V. 2004, Min Wang leg. Yinggeling Nature Reserve, 500 m,

1 male, 26. VII. 2005, Liu-Sheng Chen leg. (in SCAU). VIETNAM, Thua Thien Hue, Back Ma, 1,200 m, 2 males, 7–11. VI. 2002, Mamoru Owada leg. (in NSMT and IEBR).

MALE ADULT

Wingspan 27–30 mm. Head greyish white; antenna monopectinate, yellowish brown; labial palpus long, greyish brown, upturned. Thorax and tegula with yellowish brown scales. Abdomen yellowish brown, underside paler, fore tibia with furcella, hind tibia with two pairs of spurs. Forewing ground colour yellowish ochre, distal half brown with some irregular semitransparent spots, basal half whitish brown, middle part with an greyish brown stripe, near the stripe with a small crescent-shaped greyish brown stripe, oval-shaped semitransparent spot in the middle of these two stripes; apical angle slightly round, fringe scales grey.

Hind wing with some irregular pale yellowish white spots, and two greyish brown stripes, one is longer in the middle, another is shorter near the outer margin of hindwing; basal area yellowish white, with some black brown lines formed as reticulation. Underside of both forewing and hindwing paler than upperside.

Male genitalia: Uncus long, tenuous, apex pointed; gnathos broad, just meeting in middle, not merging; tegumen broad, rectangle-shaped; valva broad basally, tapering, apex slightly pointed, basal part slightly invaginate; costa sclerotized, with three processes, inner one long triangle, strongly toothed, central one short, and outer one longest, smooth, with Y-shaped apex; sacculus broad, with an arc-shaped incision, basala process of sacculus long, sclerotized; saccus exposure, V-shaped; aedeagus slightly slender, shrinking at middle, vesica membranous exposure.

¹Department of Entomology, College of Natural Resources & Environment, South China Agricultural University, Guangzhou 510642, China

²Department of Zoology, National Museum of Nature and Science, Amakubo 4-1-1, Tsukuba, 305-0005 Japan

*Corresponding author; E-mail: minwang@scau.edu.cn



Figs. 1-3. 1. Male adult of *Canaea bibarra* Chu et Wang. 2. Male genitalia. 3. Aedeagus.

DISTRIBUTION

China (Hainan), Vietnam (Thua Thien Hue) new record.

NOTES

The wing pattern of males from Hainan and Vietnam are identical with those of females (Chu & Wang 1991, Fig. 20; Zhu (Chu) & Wang 1996, pl. 15, Fig. 10), and the ground color is paler. This species can be distinguished from other *Canaea* species by the oval-shaped semi-transparent spot in the middle of forewing. Jianfeng, the type locality of *C. bibarra*, is located in the southwestern coast of Hainan, and is distant ca. 280 km from Back Ma, Thua Thien Hue, across the sea of Tonkin Bay. The genus *Canaea* has not been recorded from Vietnam (Konvička et al. 1998).

The structure of male genitalia of *C. bibarra* is similar to that of *C. ignotalis* (Röber, 1891) from Sulawesi, Salajar and Buru, but distinguished from the latter by the longer uncus, the shape of sclerotized processes of costa, and the membranous vesica of aedeagus. The sclerotized patch in corpus bursae of female genitalia of *C. bibarra* (Chu & Wang 1991, Fig. 20e) lacks rows of sclerotized plates, which are present (Whalley 1991, Figs. 50, 52) in *C. ignotalis*.

Acknowledgments

We are very grateful to Dr. Liu-sheng Chen, Shihezi University, for collecting the material in Hainan. Mamoru Owada expresses his hearty thanks to Dr. Ta Huy Thinh, IEBR, Hanoi, for his kind cooperation to the field researches in Vietnam.

References Cited

- Chu HF, Wang LY. 1991. The Thyrididae (Lepidoptera) of China. II. Striglininae. *Sinozoologia* 8: 325-348, pls. 1-2.
- Konvička MK, Spitzer Jaroš J. 1998. Notes of Striglininae (Lepidoptera, Thyrididae), tropical leaf moths from Vietnam, with a description of a new species. *Tropical Zoology* 11: 1-9.
- Robinson GS, Tuck KR, Shaffer M. 1994. A Field Guide to the Smaller Moths of South-East Asia. Kuala Lumpur: Malayan Nature Society, 309 pp., 32 pls.
- Shaffer M, Nielsen ES. 1996. Thyrididae. In Nielsen ES, Edwards ED, Rangsi TV [eds.], Checklist of the Lepidoptera of Australia. Monographs on Australian Lepidoptera 4: 161-163. CSIRO Division of Entomology, Canberra.
- Whalley PES. 1971. A revision of the genus *Canaea* Walker (Lepidoptera, Thyrididae). *Bulletin of the British Museum (Natural History)*, (Entomol. Series) 26(3): 161-179, pls. 1-12.
- Zhu (Chu) HF, Wang LY. 1996. Lepidoptera, Bombycidae, Saturniidae, Thyrididae. *Fauna Sinica, Insecta* 5. Science Press, Beijing, 302 pp., 18 pls.