

Feeding by the Social Wasp *Polybia scutellaris* (Hymenoptera: Vespidae) on *Syzygium jambos* (Myrtaceae) Fruits in Minas Gerais, Brazil

Authors: Brügger, Bruno Pandelo, Castro, Bárbara Monteiro de Castro e, Prezoto, Fábio, Serrão, José Eduardo, and Zanuncio, José Cola

Source: Florida Entomologist, 100(1) : 172-173

Published By: Florida Entomological Society

URL: <https://doi.org/10.1653/024.100.0125>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Feeding by the social wasp *Polybia scutellaris* (Hymenoptera: Vespidae) on *Syzygium jambos* (Myrtaceae) fruits in Minas Gerais, Brazil

Bruno Pandelo Brügger^{1,*}, Bárbara Monteiro de Castro e Castro², Fábio Prezoto³, José Eduardo Serrão⁴, and José Cola Zanuncio¹

Polybia scutellaris (White) (Hymenoptera: Vespidae), a eusocial wasp native to South America, is a generalist predator (Tellería 1996; Fernández Corujo et al. 2010) that feeds on many insect species, including pest species (Barros et al. 2015). Workers forage for food resources (proteins, carbohydrates), material for nest building (plant fiber, clay, water) (Jeanne 1996), water for thermoregulation of the colony (Kovac et al. 2009), and nectar for energy (Beggs 2001; Wegner & Jordan 2005). Social wasps can thrive in diverse habitats, including artificial or highly modified environments, such as agricultural areas (De Souza et al. 2012), and their role as a predator can be exploited as a component in pest management programs (Prezoto et al. 2006). However, these insects may also damage fruit, such as jaboticaba (*Plinia cauliflora* [Mart.] Kausel; Myrtaceae) (De Souza et al. 2010), guava (*Psidium guajava* L.; Myrtaceae) (Brugger et al. 2011), and mango (*Mangifera indica* L.; Anacardiaceae) (Barbosa et al. 2014), during foraging, which renders the fruit unsuitable for sale (Brugger et al. 2011). The wasps also are attracted to food and garbage, and forage in sites such as trade fairs, camp grounds, and picnic areas, and, therefore, frequently come into contact with people and represent a risk (Dvořák & Landolt 2006).

The foraging behavior of *P. scutellaris* on iamb fruit, *Syzygium jambos* (L.) Alston (Myrtaceae), was recorded in Feb 2016 for 5 d, hourly from 7 AM to 6 PM, on a *S. jambos* tree in Santo Antônio do Aventureiro, Minas Gerais, Brazil (21.750060°S, 42.800032°W).

Thirty-two foraging *P. scutellaris* workers were monitored, of which 26 visited ripe ground fall iamb fruits and 6 visited fruit still on the branches. Twenty wasps fed through preexisting damage and 12 pierced the skin of iamb fruits. Each visit to fruits lasted for 18 ± 7 (7–32) s, and visits occurred most often late in the morning (11 AM) and early afternoon (2 PM) (Fig. 1).

Iamb fruit on the soil used by *P. scutellaris* had preexisting damage and was fermenting (Dvořák & Landolt 2006; Landolt et al. 2014). Ground fall fruits were visited by this wasp due to recruitment and information on available food resources (D'Adamo & Lozada 2014; Sibrina et al. 2014; Santoro et al. 2015).

Temperature, light, and humidity affect social wasp foraging (Nadeau & Stamp 2003; Monceau et al. 2013), and wasp activity observed in this study was more intense in the warmer periods of the




Fig. 1. Daytime foraging activity of *Polybia scutellaris* on *Syzygium jambos* fruits. No data were collected at 1 PM.

day. Diurnal variation in the intensity of activity may be a response to environmental conditions such that the energy costs and duration of foraging are reduced by optimizing body temperature (Kovac et al. 2015; Agrawal & Karsai 2016). Also, damage to iamb fruit by foraging wasps reduces their commercial value, and actions to reduce damage may be required, such as removing wasp colonies during fruiting.

We acknowledge Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES), and Fundação de Amparo à Pesquisa do Estado de Minas Gerais (FAPEMIG) for financial support.

Summary

We report on the foraging behavior of *Polybia scutellaris* (White) (Hymenoptera: Vespidae) and the damage it causes to the fruit of *Syzygium jambos* (L.) Alston (Myrtaceae). Of 32 wasps monitored, 26 visited fruit on the ground and 6 visited fruit on the branches. Foraging was most frequent in late morning and early afternoon.

Key Words: biological control; pre-harvest damage; foraging

¹Universidade Federal de Viçosa, Departamento de Entomologia/BIOAGRO, Viçosa, Minas Gerais 36.570-900, Brasil; E-mail: brunopb2002@yahoo.com.br (B. P. B.), zanuncio@ufv.br (J. C. Z.)

²Universidade Federal de Viçosa, Departamento de Fitotecnia, Viçosa, Minas Gerais 36.570-900, Brasil; E-mail: barbaramcastro@hotmail.com (B. M. C. C.)

³Universidade Federal de Juiz de Fora, Departamento de Zoologia/LABEC, Juiz de Fora, Minas Gerais 36.036-900, Brasil; E-mail: fabio.prezoto@ufjf.edu.br (F. P.)

⁴Universidade Federal de Viçosa, Departamento de Biologia Geral, Universidade Federal de Viçosa, Viçosa, Minas Gerais 36.570-900, Brasil;

E-mail: jeserrao@ufv.br (J. E. S.)

*Corresponding author; E-mail: brunopb2002@yahoo.com.br (B. P. B.)

Sumário

Polybia scutellaris (Write) (Hymenoptera: Vespidae) preda insetos pragas, porém pode danificar cascas de frutos de *Syzygium jambos* (L.) Alston (Myrtaceae), tornando-os comercialmente inviáveis. Trinta e dois indivíduos dessa vespa foram observados, 26 visitando frutos amadurecidos no solo e seis naqueles fixos nos galhos. O forrageio de *P. scutellaris* foi mais frequente no final da manhã e início da tarde.

Palavras Chave: controle biológico; dano pré-colheita; forrageio

References Cited

- Agrawal D, Karsai I. 2016. The mechanisms of water exchange: the regulatory roles of multiple interactions in social wasps. *PLoS One* 11: e0145560.
- Barbosa BC, Paschoalini MF, Prezoto F. 2014. Temporal activity patterns and foraging behavior by social wasps (Hymenoptera, Polistinae) on fruits of *Mangifera indica* L. (Anacardiaceae). *Sociobiology* 61: 239–242.
- Barros EC, Bacci L, Picanco MC, Martins JC, Rosado JF, Silva GA. 2015. Physiological selectivity and activity reduction of insecticides by rainfall to predatory wasps of *Tuta absoluta*. *Journal of Environmental Science and Health Part B* 50: 45–54.
- Beggs J. 2001. The ecological consequences of social wasps (*Vespula* spp.) invading an ecosystem that has an abundant carbohydrate resource. *Biological Conservation* 99: 17–28.
- Brugger BP, Araújo LSS, De Souza AR, Prezoto F. 2011. Social wasps (*Synaldis cyanea*) damaging *Psidium* sp. (Myrtaceae) fruits in Minas Gerais State, Brazil. *Sociobiology* 57: 533–535.
- D'Adamo P, Lozada M. 2014. How context modification can favor the release of past experience in *Vespula germanica* wasps, enabling the detection of a novel food site. *Journal of Insect Behavior* 27: 395–402.
- De Souza AR, Venâncio DFA, Prezoto F. 2010. Social wasps (Hymenoptera: Vespidae: Polistinae) damaging fruits of *Myrciaria* sp. (Myrtaceae). *Sociobiology* 55: 297–299.
- De Souza AR, Venâncio DFA, Prezoto F, Zanuncio JC. 2012. Social wasps (Hymenoptera: Vespidae) nesting in eucalyptus plantations in Minas Gerais, Brazil. *Florida Entomologist* 95: 1000–1002.
- Dvořák L, Landolt PJ. 2006. Social wasps trapped in the Czech Republic with syrup and fermented fruit and comparison with similar studies (Hymenoptera Vespidae). *Bulletin of Insectology* 59: 115–120.
- Fernández Corujo VL, Alicia Basilio, M, Galati BG. 2010. Pollen content accumulated in nests of *Polybia scutellaris* (Hymenoptera, Vespidae). *Grana* 49: 308–313.
- Jeanne RL. 1996. Regulation of nest construction behaviour in *Polybia occidentalis*. *Animal Behaviour* 52: 473–488.
- Kovac H, Stabentheiner A, Schmaranzer S. 2009. Thermoregulation of water foraging wasps (*Vespula vulgaris* and *Polistes dominulus*). *Journal of Insect Physiology* 55: 959–966.
- Kovac H, Stabentheiner A, Brodschneider R. 2015. What do foraging wasps optimize in a variable environment, energy investment or body temperature? *Journal of Comparative Physiology A* 201: 1043–1052.
- Landolt PJ, Cha DH, Werle CT, Adamczyk JJ, Meagher RL, Gilbride RL, Clepper TS, Reed HC, Teal PEA, Sampson BJ. 2014. *Polistes* spp. (Hymenoptera: Vespidae) orientation to wine and vinegar. *Florida Entomologist* 97: 1620–1630.
- Monceau K, Maher N, Bonnard O, Thiéry D. 2013. Predation pressure dynamics study of the recently introduced honeybee killer *Vespa velutina*: learning from the enemy. *Apidologie* 44: 209–221.
- Nadeau H, Stamp N. 2003. Effect of prey quantity and temperature on nest demography of social wasps. *Ecological Entomology* 28: 328–339.
- Prezoto F, Santos-Prezoto HH, Machado VL, Zanuncio JC. 2006. Prey captured and used in *Polistes versicolor* (Olivier) (Hymenoptera: Vespidae) nourishment. *Neotropical Entomology* 35: 707–709.
- Sabrina M, D'Adamo P, Lozada M. 2014. The influence of past experience on wasp choice related to foraging behavior. *Insect Science* 21: 759–764.
- Santoro D, Hartley S, Suckling DM, Lester PJ. 2015. Nest-based information transfer and foraging activation in the common wasp (*Vespula vulgaris*). *Insectes Sociaux* 62: 207–217.
- Tellería MC. 1996. Plant resources foraged by *Polybia scutellaris* (Hym. Vespidae) in the Argentine pampas. *Grana* 35: 302–307.
- Wegner GS, Jordan KK. 2005. Comparison of three liquid lures for trapping social wasps (Hymenoptera: Vespidae). *Journal of Economic Entomology* 98: 664–666.