



Evaluating the Use of Phenylacetonitrile Plus Acetic Acid to Monitor *Pandemis pyrusana* and *Cydia pomonella* (Lepidoptera: Tortricidae) in Apple

Authors: Basoalto, Esteban, Hilton, Richard, Judd, Gary J. R., Knight, Alan L., El-Sayed, Ashraf M., et al.

Source: Florida Entomologist, 100(4) : 761-766

Published By: Florida Entomological Society

URL: <https://doi.org/10.1653/024.100.0424>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Evaluating the use of phenylacetone nitrile plus acetic acid to monitor *Pandemis pyrusana* and *Cydia pomonella* (Lepidoptera: Tortricidae) in apple

Esteban Basoalto^{1,*}, Richard Hilton², Gary J. R. Judd³, Alan L. Knight⁴, Ashraf M. El-Sayed⁵, and David Maxwell Suckling^{5,6}

Abstract

Recent studies have demonstrated that key herbivore-induced foliage volatiles from apple, *Malus domestica* Borkhausen (Rosaceae), can be attractive to conspecific adult male and female tortricid leafrollers when used together with acetic acid lures. Our study reported here was conducted in Washington State during 2013–2014 and assessed the attractiveness of both sexes of the leafroller, *Pandemis pyrusana* Kearfott (Lepidoptera: Tortricidae), to blends of acetic acid with phenylacetone nitrile. Interestingly, traps baited with phenylacetone nitrile plus acetic acid caught significantly more males, females, and total moths than the number of males caught in traps baited with a commercial sex pheromone lure. The evaporation rate of the acetic acid co-lure was shown to be an important factor affecting catches of *P. pyrusana* with phenylacetone nitrile. Adding phenylacetone nitrile to traps baited with pear ester, ethyl (*E, Z*)-2,4-decadienoate and acetic acid significantly reduced both total and female moth catches of codling moth, *Cydia pomonella* (L.) (Lepidoptera: Tortricidae). However, neither *C. pomonella* nor *P. pyrusana* catch were impacted when phenylacetone nitrile and acetic acid were added to traps baited with pear ester plus (*E, E*)-8,10-dodecadien-1-ol, the sex pheromone of codling moth. These results support our broader work to develop lures to improve monitoring and management of a number of tortricid pests attacking horticultural crops worldwide.

Key Words: kairomones; host plant volatiles; codling moth; leafroller; trapping; sex pheromone

Resumen

Estudios recientes han demostrado que volátiles foliares claves inducidos por herbívoros en manzano, *Malus domestica* Borkhausen (Rosaceae), pueden ser atractivos para hembras y machos adultos conspecíficos de enrolladores tortricidos cuando se usan junto con cebos de ácido acético. Nuestro estudio, descrito aquí, se llevó a cabo en el estado de Washington durante 2013–2014 evaluando la atracción de ambos sexos del enrollador, *Pandemis pyrusana* Kearfott (Lepidoptera: Tortricidae), a mezclas de ácido acético con fenilacetone nitrilo. Curiosamente, las trampas cebadas con fenilacetone nitrilo y ácido acético capturaron significativamente más machos, hembras y polillas totales que el número de machos atrapados en trampas provistas de un cebo comercial de feromonas sexuales. Demostrándose que la tasa de evaporación del co-attractante de ácido acético era un factor importante que afectaba las capturas de *P. pyrusana* con fenilacetone nitrilo. La adición de fenilacetone nitrilo a las trampas cebadas con éster de pera, etil (*E, Z*)-2,4-decadienoato y ácido acético redujo significativamente tanto la captura total como la de hembras de polilla de la manzana, *Cydia pomonella* (L.) (Lepidoptera: Tortricidae). Sin embargo, ni la captura de *C. pomonella* ni la de *P. pyrusana* se vieron afectadas cuando se añadieron fenilacetone nitrilo y ácido acético a las trampas cebadas con éster de pera y (*E, E*)-8,10-dodecadien-1-ol, feromona sexual de polilla de la manzana. Estos resultados respaldan nuestro trabajo más amplio para desarrollar cebos que mejoren el monitoreo y manejo de varias especies de plagas de la familia Tortricidae, que atacan cultivos hortícolas en todo el mundo.

Palabras Clave: kairomones; volátiles de plantas hospederas; polilla de la manzana; enrollador de hojas; trapeo; feromona sexual

Pandemis spp. leafrollers (Lepidoptera: Tortricidae) are an important group of pests attacking pome fruits (apple, *Malus domestica* Borkhausen [Rosaceae], and pear, *Pyrus communis* (L.) [Rosaceae], in North America (Chapman 1973; Dombroskie & Sperling 2012), Europe, and Asia (Dickler 1991). Both species are bivoltine, and overwinter-

ing larvae in the fall through spring and summer generations feed on the skin of developing fruits adjacent to leaves, thereby creating culls (Brunner 1996). In the western United States, *Pandemis pyrusana* (Kearfott) (Lepidoptera: Tortricidae), *Pandemis limitata* (Robinson) (Lepidoptera: Tortricidae), and *Choristoneura rosaceana* Harris (Lepi-

¹Instituto de Producción y Sanidad Vegetal, Facultad de Ciencias Agrarias, Universidad Austral de Chile, Campus Isla Teja, Valdivia 5090000, Chile; E-mail: esteban.basoalto@uach.cl (E. B.)

²Oregon State University, 530 Hanley Rd., Central Point, Oregon 97502, USA; E-mail: richard.hilton@oregonstate.edu (R. H.)

³Agriculture and Agri-Food Canada, Summerland Research and Development Centre, Box 5000, 4200 Highway 97, Summerland, British Columbia, Canada V0H1Z0; E-mail: gary.judd@agr.gc.ca (G. J. R. J.)

⁴Yakima Agricultural Research Laboratory, USDA-ARS, 5230 Konnowac Pass Rd, Wapato, Washington 98951, USA; E-mail: alan.knight@ars.usda.gov (A. K.)

⁵New Zealand Institute for Plant & Food Research Limited, Lincoln, Canterbury Agriculture & Science Centre, Gerald St, Lincoln 7608, New Zealand; E-mail: ashraf.el-sayed@plantandfood.co.nz (A. M. E-S.)

⁶School of Biological Sciences, University of Auckland, Tamaki Campus, Building 733, Auckland, New Zealand; E-mail: max.suckling@plantandfood.co.nz (D. M. S.)

*Corresponding author; E-mail: esteban.basoalto@uach.cl (E. B.)

doptera: Tortricidae), are the key leafroller pest species (Madsen & Madsen 1980; Brunner & Beers 1990). Insecticides often are used to manage these pests, although sex pheromone mating disruption has been developed and implemented on a relatively small acreage (Knight & Turner 1999; Judd & Gardiner 2004). Sex pheromone-baited traps are used to monitor adult populations to assess density and phenology of these pests, but pest managers feel these traps are insensitive to local population densities within the orchards (Brunner 1996). Alternative lures, including acetic acid, have been evaluated to monitor both sexes of *P. pyrusana* (Knight 2001), but acetic acid alone was found to be a relatively weak lure and was not adopted by Washington State growers (Alway 2003). In addition, acetic acid added to traps with pome fruit and green leaf volatiles, or terpenes (substances typically released by healthy or insect-infested fruit or foliage) as attractants for leafrollers did not catch significantly more moths than traps baited with acetic acid alone (Knight et al. 2014). Yet, traps baited with a mixture of (*E*, *E*)-8,10-dodecadien-1-ol (the sex pheromone of codling moth, *Cydia pomonella* [L.] [Lepidoptera: Tortricidae]), pear ester, ethyl (*E*, *Z*)-2,4-decadienoate, and acetic acid were evaluated in commercial orchards as a promising new tool (Knight et al. 2014). This multispecies approach to simultaneously monitor codling moth and leafrollers was found to be more effective in characterizing local population densities of leafroller immature stages compared with sex pheromone-baited traps (Knight et al. 2014). Because moth counts were generally low across the orchards monitored in that study, we presumed that an improved monitoring approach could be developed with the identification of more effective attractants for leafrollers.

Several recent studies have identified that the combination of acetic acid with either phenylacetoneitrile or 2-phenylethanol produced effective bisexual lures for a number of tortricid pests, including *Pandemis heparana* (Denis & Schiffermüller) (Lepidoptera: Tortricidae) in Sweden, *P. limitata* in Canada, and *P. pyrusana* in Washington State (Giacomuzzi et al. 2013; Giacomuzzi et al. 2016; Judd et al. 2017a; Giacomuzzi et al. 2017). However, the purported pheromone potency of these binary lures has perhaps been overstated (El-Sayed et al. 2016). For example, traps baited with the sex pheromone of *C. rosaceana* caught significantly more moths (3.5×) than traps with 2-phenylethanol and acetic acid (Knight et al. 2017). Similarly, traps baited with the sex pheromone of *P. limitata* caught significantly more moths (1.5×) than traps baited with the same binary lure (Judd et al. 2017a). Previously, moth catches of the eye-spotted bud moth, *Spilonota ocellana* (Denis & Schiffermüller) (Lepidoptera: Tortricidae), with acetic acid and phenylacetoneitrile had been reported to be increased significantly (> 4× greater) when added to the sex pheromone lure (El-Sayed et al. 2016). However, these data were obtained from an orchard treated the previous year with sex pheromone dispensers that could continue to emit pheromone the following year (Porcel et al. 2014). A more recent study demonstrated that traps baited with phenylacetoneitrile and acetic acid caught only 14% as many total moths as traps baited with sex pheromones in untreated orchards (Judd et al. 2017b).

Nevertheless, adoption of these new binary lures may improve our ability to monitor leafroller populations in orchards treated with sex pheromones for mating disruption. For example, the moth catch in orchards treated with sex pheromone dispensers for mating disruption of *C. rosaceana*, was significantly greater (23×) in traps baited with acetic acid and 2-phenylethanol than sex pheromone lures (Knight et al. 2017). But, the relative effectiveness of these new binary lures may not be consistent across all species. For example, moth catches in traps baited with acetic acid and phenylacetoneitrile were similar to traps baited with sex pheromone in apple orchards treated with sex pheromone dispensers for disruption of *S. ocellana* (Judd et al. 2017b).

A second useful attribute of these lures is their attraction to female moths. The sex ratio of moths caught in baited traps has been close to 1:1 in most studies. As previously shown with *Cydia pomonella* (Lepidoptera: Tortricidae), the catch of female moths can allow pest managers to develop more accurate phenological models and action thresholds (Knight & Light 2005a; Knight & Light 2005b).

Codling moth is generally the key tortricid pest of pome fruit orchards, and growers use a greater density of traps for this species than other tortricid leafrollers (Knight 2008). Some efforts to reduce monitoring costs have previously considered combining sex pheromone lures of *C. pomonella* and leafrollers in a single trap (Knight & Christianson 1999). Several studies have already examined the possible interaction of these new attractants with sex pheromones and other plant volatiles. For example, traps baited with leafroller sex pheromone caught significantly fewer males when lures loaded with either 2-phenylethanol or phenylacetoneitrile alone (*C. rosaceana*) or acetic acid plus 2-phenylethanol (*P. limitata*) were added (Knight et al. 2017; Judd et al. 2017a). The addition of (*E*)-4,8-dimethyl-1,3,7-nonatriene to traps to enhance the catch of *C. pomonella* has previously been reported and may be most useful in orchards treated with mating disruption technologies incorporating pear ester (Knight et al. 2015). However, co-monitoring *C. pomonella* and *C. rosaceana* in traps including (*E*)-4,8-dimethyl-1,3,7-nonatriene significantly reduced the catch of latter species (Knight et al. 2017). These studies suggest that the optimal development of a multi-species trapping approach combining sex pheromones with plant and food volatiles will likely require extensive testing of blend combinations. Furthermore, the correlation of moth catch and lure with pest pressure within orchards remains to be established (Knight et al. 2014). Herein, we report studies conducted from 2012–2014 to measure the response of adult *P. pyrusana* to acetic acid and phenylacetoneitrile compared with sex pheromone baited traps, and to assess the utility of combining lures for *P. pyrusana* and *C. pomonella* in the same trap.

Materials and Methods

CHEMICALS AND LURES

Sachet lures loaded with 100 mg phenylacetoneitrile (> 99% purity, Sigma Aldrich, St. Louis, Missouri, USA) consisted of a heat-sealed, semi-permeable polyethylene bag (45 × 50 mm, 150 μm wall thickness), with a piece of cellulose acetate filter (15 × 45 mm, Moss Packaging Co. Ltd., Wellington, New Zealand) inserted as the carrier substrate. Three acetic acid lures were used in our studies. Two lures were made by drilling either 1.0 or 3.2 mm holes in the cap of 8 ml Nalgene vials (Nalge Nunc International, Rochester, New York, USA) and loading each vial with 2 small cotton balls and 5 ml of glacial acetic acid (99% purity, Sigma Aldrich). A 3rd acetic acid lure consisted of a plastic membrane cup (3.4 cm diam, Pherocon AA, Trécé Inc, Adair, Oklahoma, USA) loaded with 0.72 ml of this acid. Also, halobutyl grey septa lures consisting of Pherocon CM-DA loaded with 3.5 mg pear ester, Pherocon CM-DA combo loaded with 3.5 mg of both the codling moth sex pheromone (*E*, *E*)-8,10-dodecadien-1-ol and pear ester (*E*, *Z*)-2,4-decadienoate, and Pherocon PLR (#3147) loaded with 1.0 mg of the *P. pyrusana* sex pheromone (94:6 blend of (*Z*)-11-tetradecenyl acetate and (*Z*)-9-tetradecenyl acetate) were obtained from Trécé Inc.

FIELD EXPERIMENT PROTOCOLS

Studies were conducted in 2 apple orchards: the USDA research farm situated east of Moxee, Washington, USA (46.5061°N,

120.1675°W), and in a commercial orchard located southwest of Naches, Washington, USA (46.7228°N, 120.7009°W) in 2012, 2013, and 2014. Sex pheromone dispensers for mating disruption of *P. pyrusana* and *C. pomonella* were not used in either orchard. Orange delta traps with polybutane-coated sticky inserts (Pherocon VI, Trécé Inc.) were used in all studies. Five replicates of each lure treatment were randomized and spaced 20 to 30 m apart. Traps were placed at a 2 m height in the 4 m canopy. All lures were placed on the liner. Acetic acid vials with 3.2 mm holes were used in all studies unless specified otherwise. Traps baited with a blank sachet lure were used as a negative control in each study. Moths were sexed in all studies.

MULTI-SPECIES LURE COMBINATIONS

Studies were conducted to compare moth catches of *P. pyrusana* and *C. pomonella* in blank traps and in traps baited with phenylacetoneitrile, pear ester, and acetic acid lures either alone or in binary or ternary lure combinations. The 2012 study included 6 lure treatments: blank, phenylacetoneitrile alone, pear ester plus phenylacetoneitrile, pear ester plus acetic acid, phenylacetoneitrile plus acetic acid, and phenylacetoneitrile plus pear ester plus acetic acid. Two additional lure treatments were included in the 2013 study: acetic acid alone and pear ester alone. Traps in 2012 were initially placed in the Moxee orchard on 1 Aug to monitor *C. pomonella*, and traps were checked and re-randomized on 8 Aug. This study was terminated on 13 Aug and traps were moved to the Naches orchard to monitor *P. pyrusana*. Traps were checked and re-randomized in this orchard on 22 and 30 Aug, and the study was terminated on 12 Sep. The 2013 study was conducted from 20–27 Jun at both the Moxee and Naches orchards and from 5–12 and 17–27 Aug in the Moxee orchard.

A 3rd study was conducted from 7–21 Jul 2014 in the Moxee orchard to evaluate moth catches of *C. pomonella* in traps baited with or without phenylacetoneitrile added to the sex pheromone-pear ester combinational lure plus the acetic acid cup lure. A 4th study was conducted in the Naches orchard from 30 Aug to 10 Sep 2014, to compare the magnitude of the total bisexual moth catches of *P. pyrusana* in traps with phenylacetoneitrile plus the 3.2 mm hole acetic acid vial lure with male catches in traps baited with *P. pyrusana* sex pheromone.

ACETIC ACID LURE PERFORMANCE

The mean weekly weight loss (evaporation rate) of acetic acid from the 3 lures was recorded from 30 Aug to 26 Sep 2013. Individual lures (N = 10) were placed in red Pherocon VI delta traps (Trécé Inc.) without sticky liners that were hung in the canopy of 5 linden trees, *Tilia cordata* Miller (Malvaceae), at the Yakima Agricultural Research Laboratory, Wapato, Washington, USA. Similarly, the weight loss of sachet lures loaded with phenylacetoneitrile (N = 5) was recorded from 28 Aug to 19 Sep 2013.

A field study was conducted in the Moxee orchard in 2013 to compare moth catches in delta traps baited with sachet lures loaded with phenylacetoneitrile and 1 of 3 different acetic acid lures: the membrane cup lure and vials with 1.0-mm and 3.2-mm apertures. Experiments were run for 1 wk beginning on 3 dates: 28 Aug, 4 Sep, and 12 Sep.

STATISTICAL ANALYSES

Count data were subjected to a square root transformation prior to analysis of variance (ANOVA) to normalize the variances. A randomized, complete block design was used in most studies with date and orchard as the blocking variable. Following a significant F-test in the ANOVA, means were separated with Tukey's HSD test with

an alpha value of 0.05. A completely randomized design was used to test the effect of adding phenylacetoneitrile to traps baited with the sex pheromone, pear ester, and acetic acid for *C. pomonella*. These data could not be normalized and a Kruskal-Wallis nonparametric test was used instead. Treatments with zero catches or only incidental catch (mean total catch < 0.2 moths) were not included in these analyses.

Results

MULTI-SPECIES LURE COMBINATIONS

Consistent data were collected in 2012 and 2013 when comparing catches of *C. pomonella* with phenylacetoneitrile alone and with binary and ternary blends including pear ester and acetic acid (Table 1). Pear ester in combination with acetic acid caught significantly more total and female *C. pomonella* than other lures. Adding phenylacetoneitrile to pear ester and acetic acid significantly reduced total and female moth collections. Moth abundance was low in traps baited with phenylacetoneitrile only and in binary blends with either acetic acid or pear ester but did not differ significantly from each another. In 2013, *C. pomonella* abundance from traps baited with only phenylacetoneitrile was similar to that from traps with acetic acid or pear ester alone. The addition of phenylacetoneitrile to traps baited with the sex pheromone-pear ester combination lure (Pherocon CM-DA) plus acetic acid did not significantly reduce male ($K = 0.10$; $P = 0.75$), female ($K = 2.47$; $P = 0.12$), or total ($K = 0.40$; $P = 0.53$) moth catches in 2014 (Fig. 1).

Counts of *P. pyrusana* were more varied in traps than those of coding moth during both years of the study (Table 1). During 2013, traps baited with phenylacetoneitrile plus acetic acid, or in combination with pear ester, caught significantly more total and female moths compared with the rest of the treatments and did not differ among themselves. However, moth catches in 2012 were lower than in 2013 and no significant difference among treatments was found for female moth collections. Also, during that period total moth catch, in traps baited with pear ester plus acetic acid, did not differ significantly from the 4 treatments that included phenylacetoneitrile.




Fig. 1. Mean (SE) number of male, female, and total *Cydia pomonella* caught in traps with a sex pheromone plus pear ester combo lure and an acetic acid (AA) co-lure versus in traps with these same lures with the addition of a phenylacetoneitrile (PAN) sachet lure, 7–21 Jul 2014, Moxee Washington, USA. 'N.S.' denotes a nonsignificant difference in moth catches between the 2 types of lures, $P > 0.05$.

Table 1. Comparison of catches of *Cydia pomonella* and *Pandemis pyrusana* in delta traps unbaited or baited with individual components or combinations of phenylacetoneitrile (PAN), pear ester (PE), and acetic acid (AA), N = 10 for codling moth, Moxee, Washington, USA and N = 15 for *Pandemis* leafroller, Naches, Washington, USA 2012, and N = 20 for both species, Moxee and Naches, Washington, USA 2013.

Lures ^a	Mean (SE) moth catch per trap ^b							
	2012				2013			
	<i>Cydia pomonella</i>		<i>Pandemis pyrusana</i>		<i>Cydia pomonella</i>		<i>Pandemis pyrusana</i>	
Total	Females	Total	Females	Total	Females	Total	Females	
Blank	0.2 (0.1)	0.1 (0.1)	0.0 (0.0)	0.0 (0.0)	0.1 (0.1)	0.0 (0.0)	0.0 (0.0)	0.0 (0.0)
AA	—	—	—	—	0.6 (0.2)bc	0.5 (0.2)bc	0.6 (0.2)b	0.4 (0.2)b
PE	—	—	—	—	0.7 (0.2)bc	0.3 (0.2)bc	0.2 (0.1)b	0.1 (0.1)b
PAN	0.2 (0.2)	0.0 (0.0)	0.9 (0.5)b	0.7 (0.5)	0.4 (0.2)bc	0.1 (0.1)bc	0.2 (0.1)b	0.2 (0.1)b
PE + PAN	0.7 (0.3)c	0.6 (0.2)c	0.5 (0.2)b	0.3 (0.2)	0.6 (0.2)bc	0.1 (0.1)c	0.1 (0.1)b	0.1 (0.1)b
PE + AA	15.7 (3.1)a	8.6 (1.4)a	2.2 (0.8)ab	1.2 (0.5)	5.2 (1.5)a	2.8 (0.9)a	0.8 (0.2)b	0.4 (0.2)b
PAN + AA	0.4 (0.2)c	0.2 (0.1)c	4.9 (1.4)a	1.7 (0.6)	0.2 (0.1)c	0.2 (0.1)bc	7.1 (1.6)a	3.8 (1.1)a
PAN + PE + AA	5.1 (0.9)b	2.6 (0.6)b	2.6 (1.5)ab	0.7 (0.5)	1.9 (0.6)b	1.0 (0.3)b	7.8 (1.5)a	4.7 (1.0)a
ANOVA ^c	$F_{3,35} = 42.55$ $P < 0.0001$	$F_{3,35} = 34.33$ $P < 0.0001$	$F_{4,68} = 4.92$ $P < 0.01$	$F_{4,68} = 2.03$ $P = 0.10$	$F_{6,130} = 11.72$ $P < 0.0001$	$F_{6,130} = 9.76$ $P < 0.0001$	$F_{6,130} = 39.84$ $P < 0.0001$	$F_{6,130} = 27.12$ $P < 0.0001$

Means in a column followed by a different letter were significantly different in a randomized complete block ANOVA, Tukey's HSD test, $P < 0.05$.

^aPhenylacetoneitrile was loaded on a felt pad in plastic sachets at 100 μ l. Pherocon CM DA lures contained 3.0 mg pear ester. Acetic acid (5 ml) was loaded on cotton balls in plastic vials with a 3.2 mm hole.

^bExperiments were repeated on 2 dates in the Moxee orchard: 1–8 and 8–13 Aug 2012; and on 4 orchard-dates: 20–27 Jun in both the Moxee and Naches orchards, and from 5–12 and 17–27 Aug in the Moxee orchard during 2013.

^cData from blank traps for both species and with PAN lures for *C. pomonella* in 2012 were not included in the analyses.

COMPARISON OF ACETIC ACID AND PHENYLACETONITRILE WITH SEX PHEROMONE

Pandemis pyrusana from traps placed in the Naches orchard were considerably greater in 2014 compared with 2013 (Fig. 2). Total moth catch was significantly different and nearly 6-fold higher in traps baited with phenylacetoneitrile and acetic acid than the sex pheromone ($F_{1,8} = 54.64$; $P < 0.001$). Mean catch of male ($F_{1,8} = 9.30$; $P < 0.05$) and female ($F_{1,8} = 20.01$; $P < 0.01$) moths were significantly greater with this binary lure compared with male collections.

LURE PERFORMANCE

The average evaporation rate of acetic acid from the co-lures used with phenylacetoneitrile was a significant factor affecting moth catches




Fig. 2. Mean (SE) number of male, female, and total *Pandemis pyrusana* caught in traps baited with either the sex pheromone lure (males only) or with phenylacetoneitrile plus an acetic acid co-lure (males, females, and total moths). '*' denotes a significant mean difference as compared to male catch in the sex pheromone-baited trap, $P < 0.05$.

of *P. pyrusana* (Table 2). Evaporation rates ranged 18-fold across the 3 dispensers tested. Moth catches were significantly lower in traps containing acetic acid lures with the lowest evaporation rate (plastic cup lure) and did not differ between traps with different hole sizes (Table 2). Mean evaporation rate of phenylacetoneitrile from the sachet lure was 6.3 ± 0.2 mg per d.

Discussion

Our results show that the use of a binary lure of phenylacetoneitrile plus acetic acid has potential use in developing improved pest management of *P. pyrusana*. Traps baited with phenylacetoneitrile and acetic acid caught greater numbers of moths than traps baited with sex pheromone in an orchard not treated with leafroller sex pheromones. In comparison, sex pheromone lures were consistently more attractive for males and caught significantly more total numbers of moths than the use of acetic acid with either phenylacetoneitrile or 2-phenylethanol when tested with *C. rosaceana*, *S. ocellana*, or *P. limitata* (Knight et al. 2017; Judd et al. 2017a; b). These data may suggest that the commercial lures for *P. pyrusana* are not optimized, or that other species in the *P. limitata* (Robinson) group with different sex pheromone blends are more dominant in the 2 orchards studied (Dombroskie & Sperling 2012). The sex pheromones of *P. limitata* and *P. pyrusana* have been identified as 91:9 and 94:6 blends of (Z)-11-tetradecenyl acetate and (Z)-9-tetradecenyl acetate, respectively (Roelofs et al. 1976, 1977). However, previous studies conducted in 1996 with these 2 blend ratios found that either blend caught similar numbers of *Pandemis* adults in a different Washington apple orchard (Knight unpublished). Curiously, at least 1 commercial lure has previously been reported to be relatively ineffective for *P. pyrusana* due to contamination with < 1.0% of (Z)-9-dodecenyl acetate (Brunner & Fisher 1998). Because of the uncertainty in comparative performance of sex pheromone lures, it is likely that further studies are needed to validate our findings.

Apple orchards are commonly attacked by more than 1 species of leafroller, so this group of pests must be carefully monitored

Table 2. Comparison of catches of *Pandemis* leafroller in delta traps baited with phenylacetoneitrile plus 1 of 3 different acetic acid co-lures, N = 15, Moxee Washington, USA, 2013.

Type of acetic acid lure ^a	Mean (SE) daily weight loss (mg) from lure	Mean (SE) weekly moth catch ^b	
		Total	Females
Plastic cup	4.9 (0.1)c	10.2 (2.2)b	3.3 (0.8)b
1.0-mm vial	16.1 (2.0)b	26.8 (7.2)a	9.1 (2.4)a
3.2-mm vial	88.6 (3.3)a	17.1 (3.5)a	6.8 (1.4)a
ANOVA	$F_{2,27} = 500.8$ $P < 0.0001$	$F_{2,40} = 12.56$ $P < 0.001$	$F_{2,40} = 10.42$ $P < 0.001$

Means in a column followed by a different letter were significantly different in a randomized complete block ANOVA, Tukey's HSD test, $P < 0.05$.

^aAcetic acid lures included the proprietary Pherocon AA (Trécé Inc., Adair, Oklahoma, USA) plastic cup and polypropylene vials filled with 5 ml of acetic acid and with either 1.0 or 3.2 mm holes. Phenylacetoneitrile sachet lures released an average (SE) of 6.3 (0.2) mg d⁻¹.

^bExperiments were repeated on 3 dates: 28 Aug–4 Sep, 4–11 Sep, and 12–19 Sep 2013, in the Moxee orchard.

during the season (Chapman 1973). Studies have found either that acetic acid plus 2-phenylethanol performs better than phenylacetoneitrile and acetic acid (Knight et al. 2017), similarly (Giacomuzzi et al. 2013, 2016; Giacomuzzi et al. 2017), or worse (El-Sayed et al. 2016) with different tortricid species. A ternary blend of acetic acid and both compounds also can significantly increase moth catches compared with either binary blend (Judd et al. 2017b). Clearly, additional studies are needed to optimize these aromatic plus acetic acid lures for various geographical regions attacked by different suites of tortricid leafrollers. In addition, more studies will be needed to assess their utility in monitoring tortricids in orchards treated with their respective sex pheromones. Ultimately, we expect that developing improved timing models and pest thresholds based on these lures can significantly improve management of orchards with reduced insecticide inputs.

The ability of pest managers to monitor multiple species within single traps could fuel the adoption of labor-saving, remote electronic traps (Guarnieri et al. 2011; Kim et al. 2011). Our study found that adding pear ester to traps baited with phenylacetoneitrile and acetic acid did not impact the catch of *P. pyrusana*. Nor did it affect the catch of *C. pomonella* when phenylacetoneitrile and acetic acid were added to traps baited with the combination sex pheromone plus pear ester lure. This latter combination lure is considered to be the industry standard for orchards using mating disruption on codling moth (Knight et al. 2005). Future studies should consider the creation of bait stations for even more co-occurring tortricid pests within orchards, such as the oriental fruit moth *Grapholita molesta* (Busck) (Lepidoptera: Tortricidae). Development of binary pheromone lures combining the sex pheromone of *C. pomonella* and *G. molesta* already has been developed (Knight et al. 2015), and testing of various host plant volatiles for this latter species have been reported (Lu et al. 2014; Yu et al. 2015). Further testing and refinement of this multi-species approach could lead to the development of both lower-cost monitoring programs and perhaps multi-species management tools, i.e., lure and kill strategies.

Acknowledgments

We would like to thank Duane Larson and Chey Temple, Agricultural Research Service, Wapato, Washington, USA, for their technical assistance in the laboratory and field. We would like to acknowledge our appreciation to Bill Lingren, Trécé Inc., Adair, Oklahoma, USA, for donating the commercial lures used in our studies. This project was supported with partial funding from the Washington Tree Fruit Research Commission, Wenatchee, Washington, USA.

References Cited

- Alway T. 2003. Exploring alternatives to organophosphate insecticides with apple and pear: the Areawide II Project in Washington State. *Compact Fruit Tree* 36: 91–93.
- Brunner JF. 1996. Management of leafrollers in Washington apple orchards, pp. 257–263. *In* Washington State Horticultural Association [Eds], *Proceedings Washington Horticultural Association 91st Annual Meeting*, Wenatchee, Washington, USA, 4–6 Dec 1995. Western Washington Horticultural Association, Pullman, Washington, USA.
- Brunner JF, Beers EH. 1990. *Pandemis* and obliquebanded leafrollers. Washington State University Extension Bulletin 1582, Wenatchee, Washington, USA.
- Brunner JF, Fisher J. 1998. Inhibition of leafroller captures in pheromone traps, pp. 26–27. *In* Tree Fruit Research & Extension Center [Eds], *Proceedings of the 79th Western Orchard Pest and Disease Management Conference*, Portland, Oregon, USA, 7–9 Jan 1998. Washington State University, Wenatchee, Washington, USA.
- Chapman PJ. 1973. Bionomics of the apple-feeding Tortricidae. *Annual Review of Entomology* 18: 73–96.
- Dickler E. 1991. Tortricid pests of pome and stone fruits, Eurasian species, pp. 435–452. *In* van der Geest LPS, Evenhuis HH [Eds], *Tortricoid Pests*. Elsevier, Amsterdam, The Netherlands.
- Domroskie JJ, Sperling FAH. 2012. Phylogeny of Nearctic *Pandemis* (Lepidoptera: Tortricidae), with focus on species boundaries in the *P. limitata* group. *Annals of the Entomological Society of America* 105: 768–780.
- El-Sayed AM, Knight AL, Byers JA, Judd GJR, Suckling DM. 2016. Caterpillar-induced plant volatiles attract conspecific adults in nature. *Scientific Reports* 6: 37555.
- Giacomuzzi V, Abraham J, Angeli S. 2013. Feeding damage of *Pandemis heparana* induces the release of specific volatile compounds from apple plants, p. 453. *In* Tielkes E [Ed], *Proceedings of the Conference of International Research in Food Security, Natural Resource Management and Rural Development*. University of Hohenheim, Germany, 17–19 Sep 2013. Cuvillier Verlag, Göttingen, Germany.
- Giacomuzzi V, Cappellin L, Khomenko I, Biasioli F, Schutz S, Tasin M, Knight AL, Angeli S. 2016. Emission volatile compounds from apple plants infested with *Pandemis heparana* larvae, antennal response of conspecific adults, and preliminary field trial. *Journal of Chemical Ecology* 42: 1265–1280.
- Giacomuzzi V, Mattheis J, Basoalto E, Angeli S, Knight AL. 2017. Survey of conspecific herbivore-induced volatiles from apple as possible attractants for *Pandemis pyrusana* (Lepidoptera: Tortricidae). *Pest Management Science* 73: 1837–1845. doi: 10.1002/ps.4548.
- Guarnieri A, Maini S, Molari G, Rondelli V. 2011. Automatic trap for moth detection in integrated pest management. *Bulletin of Insectology* 64: 247–251.
- Judd GJR, Gardiner MGT. 2004. Simultaneous disruption of pheromone communication and mating in *Cydia pomonella*, *Choristoneura rosaceana* and *Pandemis limitata* (Lepidoptera: Tortricidae) using Isomate-Cm/LR in apple orchards. *Journal of the Entomological Society of British Columbia* 101: 3–14.
- Judd GJR, Knight AL, El-Sayed AM. 2017a. Trapping *Pandemis limitata* (Lepidoptera: Tortricidae) moths with mixtures of acetic acid, caterpillar-induced apple leaf volatiles and sex pheromone. *The Canadian Entomologist*, in press.
- Judd GJR, Knight A, El-Sayed AM. 2017b. Development of kairomone-based lures and traps targeting *Spilonota ocellana* (Lepidoptera: Tortricidae) in apple orchards treated with sex pheromones. *The Canadian Entomologist*, in press.
- Kim Y, Jung S, Kim Y, Lee Y. 2011. Real-time monitoring of oriental fruit moth, *Grapholita molesta*, populations using a remote sensing pheromone trap in apple orchards. *Journal of Asia-Pacific Entomology* 14: 259–262.

- Knight AL. 2001. Monitoring the seasonal population density of *Pandemis pyrusana* (Lepidoptera: Tortricidae) within a diverse fruit crop production area in the Yakima Valley, WA. *Journal of the Entomological Society of British Columbia* 98: 217–225.
- Knight AL. 2008. Codling moth areawide integrated pest management, pp. 159–190. In Koul O, Cuperus GW, Elliot N [Eds], *Areawide Pest Management: Theory and Implementation*. CAB International, Wallingford, United Kingdom.
- Knight AL, Turner JE. 1999. Mating disruption of *Pandemis* spp. (Lepidoptera: Tortricidae). *Environmental Entomology* 28: 81–87.
- Knight A, Christianson B. 1999. Using traps and lures in pheromone-treated orchards. *Good Fruit Grower* 150: 45–50.
- Knight AL, Light DM. 2005a. Timing of egg hatch by early-season codling moth (Lepidoptera: Tortricidae) predicted by moth catch in pear ester- and codlemone-baited traps. *The Canadian Entomologist* 137: 728–738.
- Knight AL, Light DM. 2005b. Developing action thresholds for codling moth (Lepidoptera: Tortricidae) with pear ester- and codlemone-baited traps in apple orchards treated with sex pheromone mating disruption. *The Canadian Entomologist* 137: 739–747.
- Knight AL, Hilton R, Light DM. 2005. Monitoring codling moth (Lepidoptera: Tortricidae) in apple with blends of ethyl (*E, Z*)-2, 4-decadienoate and codlemone. *Environmental Entomology* 34: 598–603.
- Knight AL, Hilton R, Basoalto E, Stelinski LL. 2014. Use of glacial acetic acid to enhance bisexual monitoring of tortricid pests with kairomone lures in pome fruits. *Environmental Entomology* 43: 1628–1640.
- Knight AL, Basoalto E, Katalin J, El-Sayed AM. 2015. A binary host plant volatile lure combined with acetic acid to monitor codling moth (Lepidoptera: Tortricidae). *Environmental Entomology* 44: 1434–1440.
- Knight AL, Judd GJR, Basoalto E, El-Sayed AM. 2017. Development of 2-phenyl-ethanol plus acetic acid lures to monitor obliquebanded leafroller (Lepidoptera: Tortricidae) under mating disruption. *Journal of Applied Entomology*, in press. doi: 10.1111/jen.12393.
- Lu PF, Qiao HL, Xu ZC, Cheng J, Zong SX, Luo YQ. 2014. Comparative analysis of peach and pear fruit volatiles attractive to the oriental fruit moth, *Cydia molesta*. *Journal of Plant Interactions* 9: 388–395.
- Madsen HF, Madsen BJ. 1980. Response of four leafroller species (Lepidoptera: Tortricidae) to sex attractants in British Columbia orchards. *Canadian Entomologist* 112: 427–430.
- Porcel M, Sjöberg S, Swiergiel W, Dinwiddie R, Rämert B, Tasin M. 2014. Mating disruption of *Spilonota ocellana* and other apple orchard tortricids using a multispecies reservoir dispenser. *Pest Management Science* 71: 562–570.
- Roelofs WL, Cardé A, Hill A, Cardé RT. 1976. Sex pheromone of the threelined leafroller, *Pandemis limitata*. *Environmental Entomology* 5: 649–652.
- Roelofs WL, Lagier RF, Hoyt SC. 1977. Sex pheromone of the moth, *Pandemis pyrusana*. *Environmental Entomology* 6: 353–354.
- Yu H, Feng J, Zhang Q, Xu H. 2015. (*Z*)-3-hexenyl acetate and 1-undecanol increase male attraction to sex pheromone trap in *Grapholita molesta* (Busck) (Lepidoptera: Tortricidae). *International Journal of Pest Management* 61: 30–35.