



Genomic Structure of the Luciferase Gene from the Bioluminescent Beetle, *Nyctophila* cf. *Caucasica*

Authors: Day, John C., Chaichi, Mohammad J., Najafil, Iraj, and Whiteley, Andrew S.

Source: Journal of Insect Science, 6(37) : 1-8

Published By: Entomological Society of America

URL: <https://doi.org/10.1673/031.006.3701>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.



Genomic structure of the luciferase gene from the bioluminescent beetle, *Nyctophila cf. caucasica*

John C. Day¹, Mohammad J. Chaichi², Iraj Najafil³ and Andrew S. Whiteley¹

¹ CEH-Oxford, Mansfield Road, Oxford, Oxfordshire, OX1 3SR, UK

² Department of Chemistry, Mazandaran University, Babolsar, Iran.

³ Biological Control Centre, Amol, Iran

Abstract

The gene coding for beetle luciferase, the enzyme responsible for bioluminescence in over two thousand coleopteran species has, to date, only been characterized from one Palearctic species of Lampyridae. Here we report the characterization of the luciferase gene from a female beetle of an Iranian lampyrid species, *Nyctophila cf. caucasica* (Coleoptera:Lampyridae). The luciferase gene was composed of seven exons, coding for 547 amino acids, separated by six introns spanning 1976 bp of genomic DNA. The deduced amino acid sequences of the luciferase gene of *N. caucasica* showed 98.9% homology to that of the Palearctic species *Lampyris noctiluca*. Analysis of the 810 bp upstream region of the luciferase gene revealed three TATA boxes and several other consensus transcriptional factor recognition sequences presenting evidence for a putative core promoter region conserved in Lampyrinae from –190 through to –155 upstream of the luciferase start codon. Along with the core promoter region the luciferase gene was compared with orthologous sequences from other lampyrid species and found to have greatest identity to *Lampyris turkistanicus* and *Lampyris noctiluca*. The significant sequence identity to the former is discussed in relation to taxonomic issues of Iranian lampyrids.

Keywords: Coleoptera, Lampyridae, phylogeny, promoter, *Lampyris turkistanicus*, *Lampyris noctiluca*

Correspondence to: jcd@ceh.ac.uk, jchaichi@yahoo.com, najafil2004@yahoo.com, aswhi@ceh.ac.uk

Received: 14.7.2005 | Accepted: 8.2.2006 | Published: 27.10.2006

Copyright: Creative Commons Attribution 2.5

ISSN: 1536-2442 | Volume 6, Number 37

Cite this paper as:

Day JC, Chaichi MJ, Najafil I, Whiteley AS. 2006. Genomic structure of the luciferase gene from the bioluminescent beetle, *Nyctophila cf. caucasica*. 8pp. *Journal of Insect Science* 6:37, available online: insectscience.org/6.37

Introduction

Bioluminescence is a process by which living organisms convert chemical energy into light. Although evident in land animals, bioluminescent organisms predominate in marine environments with only a few groups of terrestrial animals exhibiting the necessary components to generate visible light. These elements are an enzyme, luciferase, and a substrate, luciferin, which are structurally diverse in nature. The majority of bioluminescent beetle species belong to the family Lampyridae of which the firefly species *Photinus pyralis* is the most studied. Firefly luciferase (EC 1.13.12.7) from *P. pyralis* is a 62-kDa enzyme that catalyses emission of yellow-green light ($\lambda_{\max} = 560$ nm) upon reaction of D-luciferin, ATP and molecular oxygen (White 1971; McElroy and DeLuca 1983; Baldwin 1996; Conti et al 1996; Wood 1995). The cDNA for the *P. pyralis* luciferase was first characterized in 1985 (de Wet et al 1985) and over the years has been extensively studied, including the resolution of its tertiary structure (Conti et al 1996). This wealth of information has been facilitated by the use of luciferase in a range of applications exploiting the bioluminescent function of this enzyme and the requirement of ATP within the reaction. However, work has centred upon a few luciferase sequences obtained primarily from Nearctic and Oriental species. An expansion of luciferase studies to include Palearctic species would present a more complete dataset for phylogenetic studies as well as providing novel sequences for expression purposes. To date only one luciferase sequence from a Palearctic species has been characterized, the European glow-worm *Lampyris noctiluca* (Sala-Newby et al 1996). *Lampyris noctiluca* belongs to the tribe Lampyrini, which is composed of five genera, *Diaphanes*, *Lampyris*, *Nyctophila*, *Pelania* and *Pyrocoelia*. The genus *Nyctophila* was established by Olivier (Olivier, 1884), with the type specimen *N. reichii* Du Val being described in 1859, and is comprised of about 30 known species, most of which are described from Europe and the Middle East (Geisthardt and Satô, in press).

In this study we identify and examine a novel luciferase gene from *Nyctophila* cf. *caucasica* collected from the Amol forest, northern Iran and compare the luciferase gene sequence and promoter region with that of other Lampyridae species.

Materials and Methods

Specimens, taxonomy and DNA extraction

N. caucasica male and female adult specimens were provided from a maintained colony at Mazandaran University, originally collected from Amol forest, Mazandaran Province, Northern Iran (36°28'N, 52°21'E) and shipped in alcohol to Oxford, England. Total genomic DNA was extracting from a single female specimen using the High Pure PCR Template Preparation Kit (Roche, www.roche.com) according to the manufacturer's instructions.

PCR and genome walking of the luciferase gene from *N. caucasica*

Based upon the luciferase sequence from *L. noctiluca* (GenBank accession # X89479) two PCR primers, noctlucF1 and noctlucR2 (Table 1) were used to amplify the first 1kb of the luciferase gene from *N. caucasica* (Figure 1). PCR was carried out under the following conditions: initial denaturation at 94 °C for 2 min, ten cycles at 94 °C for 15 sec, 60 °C for 30 sec, and 72 °C for 6 min, twenty cycles at 94 °C for 15 sec, 60 °C for 30 sec, and 72 °C for 6 min plus an additional 5 sec per cycle and a final extension at 72 °C for 7 min. PCR products were ligated into the pGEM®-T Easy Vector System (Promega, www.promega.com/) and ligation mixtures transformed into competent cells of *Escherichia coli* DH5 α . Complete nucleotide sequences of PCR products were determined using a dye termination kit and an automatic sequencer (Beckman Coulter, www.beckman.com). From the sequence four primers, Luc5'GW1, Luc5'GW2, Luc3'GW1 and Luc3'GW2 (Table 1) were designed to amplify the remaining downstream sequence of luciferase from *N. caucasica* using the Universal GenomeWalker™ protocol (Clontech, www.clontech.com). PCR products were cloned and sequenced as described above. Two primers, NycLuc F1 and NycLuc R1 (Table 1) were used to amplify the entire gene and confirm continuity of sequence.

Table 1. PCR primers used in the amplification of luciferase from *Nyctophila caucasica*.

Oligonucleotide	Sequence (5' - 3')
noctluc F1	CACTAACGCGCTAATATCATTG
noctluc R3	AAGACAATACGAAATCCAC
NycLuc5' GW1	TTGTCCTGGAAACCTGTGCATACCTCTT
NycLuc5' GW2	TGTTCTCCAGCAGTCCATCCTCCAAG
NycLuc3' GW1	TCTTGAAACTATMAAAATGTAATTGTAT
NycLuc3' GW2	ACTGCGATTTTAACGGTTATACCATTTTC
NycLuc F1	GTATAATGTCTCTAGGTCTCTCTC
NycLuc R1	TATACGTGAAAATGAAAATACAGTGT

```

ATGGAAGATGCAAAAATATTATGCACGGTCCACCGCCATTCTATCCTTTGGAGGATGGAAGTCTGGAGAACAAATGCACAAAGCAATGAAGAGGTATGCACAGGTTCCAGGGACAATT 120
.....G.....
M E D A K N I M H G P P P F Y P L E D G T A G E Q L H K A M K R Y A Q V P G T I
GCTTTTgtaagtaaaattattatcttattcattatcaaatccaatgcaaatcttttgcgaatttgagagtttaacctttttttgtacagACYGATGCACATGACAGGTAATAA 240
.....T.....G.....
A F ..... T D A H A E V N
TTACATATCCGAATATTTGAAATGGCTTGCCGGTTAGCCGAAACTATGAAGAGGTACGGACTTGGTTTACAACACCACATTGCTGTTTGCAGCGAAAATTTCTTCAGTTTTTTATGC 360
.....T.....
I T Y S E Y F E M A C R L A E T M K R Y G L G L Q H H I A V C S E N S L Q F F M
CTGTATGCGGTGCTCTATTTATGGAGTTGGAGTTGCACCAACAACAGATATTACAATGAACgtaagccctttgtcagctctaaaaaatgatgtatayactctctacagGTGAATTAT 480
.....G.....T.....T.....
P V C G A L F I G V G V A S T N D I Y N E ..... R E L
ACAACAGTTTGTCCATATCACAACCTACAATAGTATTCTGTTCCAAAAGAGCGCTGCAAAAATCCTAGGGTACAAAAGAAATTACCTATAATTCAGAAAATTTGTTATTCTGGATTCTC 600
.....C.....
Y N S L S I S Q P T I V S C S K R A L Q K I L G V Q K K L P I I Q K I V I L D S
GAGAGATTATATGGGAAAACATCTATGTACTCGTTCATGAATCTCATTACCTCGAGGTTTTAATGAATATGATTACATACCGGATTCATTGACCGCGAAAACAGCAACAGCACTTA 720
R E D Y M G K Q S M Y S F I E S H L P A G F N E Y D Y I P D S F D R E T A T A L
TAATGAATTCATCGGATCTACCGGATACCCAGGGAGTTGAGCTTACTCACAACAACATTTGTTAGATTCTCAGTGCAGgtacgtatggctctgaaactataaaaatgtaattg 840
.....T.....G.....C.....TG.G.....
I M N S S G S T G L P K G V E L T H Q N V C V R F S H C R
tattcaaaaattgtagAGATCTCTGTTGGTAATCAAATTTATCCCGATCTGCGGATTTAACCGTTTATACCATTTCATCATGGTTTTGGAATGTTTACAACACTAGGATATTTAAGC 960
.....A.....
D P V F G N Q I I P D T A I L T V I P F H H G F G M F T T L G Y L T
TGTGGATTTCGTATTGCTTATGTATAGATTGAAGAGGAATTTTTTACGATCACTTCAAGATTATAAAATCAAAGTGCCTGCTGGTACCTACTCTGTTTTCATTCTTGGCCAAA 1080
.....A.....
C G F R I V L M Y R F E E E L F L R S L Q D Y K I Q S A L L V P T L L F S F F A K
AGCACCTTAGTCGACAAATACGATTTATCCAACCTACATGAAATGCTTCTGTTGGAGCTCCCTCGCGAAAAGTGGAGAAGCTGAGCAAAACGgtaagtcgcgataactaagtaact 1200
.....T.....G.....
S T L V D K Y D L S N L H E I A S G G A P L A K E V G E A V A K R
cagtttctaataaggcttttagTTTTAAGCTTCCGGGCATACGACAAGGGTATGGAAGTACTGAAACTACCTCAGCTATTATAATTACACCAGAAGGGGATGATAAACAGGAGCATGT 1320
.....G.....A.....
F K L P G I R Q G Y G L T E T T S A I I I T P E G D D K P G A C
GGTAAAGTTGTTCCATTCTTTCTGCCAAAATGTTGATCTGGATACCGGTAACCTTTGGTGTATATCAGAGGGGAAATATGTGTGAAAGGCCCAATGATAATGAAGGTTACGTA 1440
G K V V P F F S A K I V D L D T G K T L G V N Q R G E L C V K G P M I M K G Y V
AACAACCCAGAAGCAACAGTGCATTGATAGACAAAGACGGATGTTACTCTGTTGATAGCTTACTACGACAAAGATGGTCACTTCTCATAGTAGATCGTTTGAATCGTTAATT 1560
.....T..T.....
N N P E A T S A L I D K D G W L H S G D I A Y Y D K D G H F F I V D R L K S L I
AAATCAAAGTTATCAGgtaaaacgctgatttttagtatacgttaggtataattacctaacaatgtctatagGTACCGCTGCCGAATTAGAATCGATATTGCTGCAACATCCGTTTATA 1680
.....C.....
K Y K G Y Q ..... V P P A E L E S I L L Q H P F I
TTTGTGACAGGTGTTGCAGGAATCCCGACCCAGATGCCGGTGAACCTCTCTGACGCGTGTGTTGCTTAGAGGAAGGCAAAACGATGACTGAACAAAGATGATGGATTATGTTGCGGgt 1800
F D A G V A G I P D P D A G E L P A A V V V L E E G K T M T E Q E V M D Y V A
aaattaaccatttatgCGGTACATAGTACATCCTTATTATAGGACAAGTAACTGCTTCTAAGCGTTTACGTTGAGGAGTAAAGTTTGTGGACGAAGTACCTAAAGGTCTAACTGGAA 1920
.....G.....
G Q V T A S K R L R G G V K F V D E V P K G L T G
AGATTGATCAAGAAAATCAGGGAGATTCTCATGATGGGAAAAAATCCAAATTTGTA 1979
.....G.....C..T.....
K I D G R K I R E I L M M G K K S K L *

```

Figure 1. The nucleotide sequence and genomic organization of the luciferase gene from *Nyctophila caucasica* (upper sequence) aligned with the cDNA luciferase sequence obtained from *Lampyrus noctiluca*, GenBank accession number X89479 (lower sequence). Sequence identity is illustrated with a dot and deletions are indicated with a dash. Exon sequences are shown in bold uppercase, introns in lowercase. Amino acid differences between *N. caucasica* and *L. noctiluca* are shown underlined.

Sequence and phylogenetic analysis

DNA sequences from *N. caucasica* were edited and aligned using Sequencher 4.0.5 (Gene Codes Corporation, www.genecodes.com). A consensus sequence was aligned with the cDNA sequence from *L. noctiluca* (GenBank accession # X89479) in order to determine exon/intron positions. The exon positions were further confirmed by comparisons with the *L. noctiluca* gene sequence (accession # AY447204) (Li et al 2003b). All sequences used in the phylogenetic analysis were obtained from the DDBJ/GenBank/EMBL

database and accession numbers are as follows: *Cratomorphus distinctus* (AY633557) *Lampyrus turkistanicus* (AY742225); *Hotaria papariensis* (AF486802); *Hotaria parvula* (L39929); *Hotaria tsushiman* (AF486801); *Hotaria unmunšana* (AF420006); *Lampyrus noctiluca* (X89479); *Luciola cruciata* (P13129); *Luciola lateralis* (U51019); *Luciola mingrelica* (S61961); *Photinus pyralis* (M15077); *Photuris pennsylvanica* (U31240); *Pyrocoelia miyako* (L39928); *Pyrocoelia rufa* (AF328553); *Phrixothrix hirtus* (AF139645); *Phrixothrix vivianii* (AF139644) and

Pyrophorus plagiophthalmus (S29355). Alignments were carried out using ClustalW followed by manual modification and phylogenetic analysis was carried out using PAUP Vers 4.0b8 (Swofford 2001). The tree was rooted with CG6178 a sequence regarded as a non-bioluminescent ortholog of beetle luciferase from the *Drosophila* genome (Ohba et al 2004).

Results and Discussion

PCR amplification strategies and DNA sequencing were successfully used to isolate and characterize the luciferase gene from a single female *N. caucasica*. *Lampyris noctiluca* PCR primers LnocF1 and Lnoc R2 were sufficiently conserved to generate a PCR product from *N. caucasica* 1008 bp in size. Sequencing provided information for genome walking primers that amplified products upstream and downstream of the luciferase gene in *N. caucasica*. Primers designed at the ends of the genome walking products were used to amplify the entire luciferase gene and both upstream and downstream regions of the gene as one continuous fragment 3086 bp in length. Sequencing revealed the PCR product to be composed of 810 bp of 5' sequence upstream of the luciferase start codon and 242 bp of 3'

sequence downstream of the stop codon. From the cloned PCR fragments two alleles were identified denoted as *Ncau1* and *Ncau2*. Four synonymous transitions were identified between the two alleles of which only one was located within an exon. The luciferase gene was composed of seven exons, coding for 547 amino acids, separated by six introns and spanning 1976 bp of genomic DNA (Fig. 1 and Figure 2). The entire 3086 bp sequence, including primer sequence, was deposited in GenBank, accession # DQ072141.

To date, the entire luciferase gene has been characterized from six different Lampyridae species and in all species the luciferase gene is composed of seven exons divided by six introns. The introns are relatively conserved in size, with the most size variation occurring in intron 1, the largest intron present in the *N. caucasica* luciferase gene (Table 2). One of the most interesting differences between the luciferase gene sequences is that found between the two populations of *P. rufa* reported by Li *et. al.* (Li et al 2003a) in which three amino acid substitutions occur with extensive variation between intron sequences from individuals of Chinese and




Figure 2. Luciferase gene characterization from *Nyctophila caucasica*. A. Genome organization of the luciferase gene from *N. caucasica* and PCR primer positions. B. Deduced luciferase amino acid sequence alignment of *N. caucasica* (*Nyct.*) and *Lampyris noctiluca* (*Lamp.*) (Sala-Newby *et al* 1996).

Table 2. The size of the six introns in the luciferase gene from *Nyctophila caucasica* reported in this paper, compared to those in other lampyrid species.

Species	Intron size (bp)					
	1	2	3	4	5	6
<i>Nyctophila caucasica</i>	89	48	52	45	54	47
<i>Hotaria species</i> ¹	52	52	52	51	49	52
<i>Lampyrus noctiluca</i> ²	87	48	49	45	54	52
<i>Luciola lateralis</i> ³	77	54	49	50	46	52
<i>Photinus pyralis</i> ⁴	57	51	48	49	43	47
<i>Pyrocoelia rufa</i> I ⁵	74	51	48	45	54	52
<i>Pyrocoelia rufa</i> II ⁵	84	53	54	46	54	48

¹Intron size and position conserved in three *Hotaria* species, *H. unmunzana*, *H. papariensis* and *H. tsushima*, Choi et al. 2003

²Li et al., (2003b).

³Cho et al., 1999.

⁴de Wet et al., 1987.

⁵Li et al. (2003a) characterised the luciferase gene from two different populations of *P. rufa*, a Korean specimen (*P. rufa* I) and a Chinese population (*P. rufa* II)

```

-810                                     gtataatgtc
-800  tcctaggttc ttctcgttga agctaggtac aacggcttag gctggtagct
-750  tcaggagcac gaatagaaat ggtgggattt ctgcgcataa aagcgaataa
-700  ccattccttt tcagcttctt ttttttctt cccactgatg aggtaattca
-650  agtttcaatg cgacagcata ctgataagca agtttcttta cttgttttgt
-600  gctgagacca taatgcattt tgaatgcagt tttgaaatat tcacaaacca
-550  tttcttctg  ttgcttgctg aacacttctt taacagcgac attaggattg
-500  tatttaatgg tttttacgta cttacatgac gtgctagtga tgattttggg
-450  atgttacatg catttgctac aagtctaag gaactaccct tcctaacttc
-400  ttcgctttgt tattcttaag aaagttaag taggtaagaa gttaagtaag
-350  cgaataagag atttgatagg taagtcttaa aatgtcgtat agaattttga
-300  aagacaaaac ggtttgctaa aagatttaa aagaaagtta aagtatctca
-250  aaaattcgtt aatggcaata aagttgctga agtagttgtt aaacgcgcta
-200  agatttttat tatatacatt tatgtcctca tgtttataaa agcaatattt
-150  acgtggtata aactgtcaaa taaaatttaa gtaatgtgat tacgagtcaa
-100  aggtctttca caaaaagta ttaagatac ataatcatal aatgtgacaa
-50  ttcgagaata ctggttctg agacactaac gcgctaatat cattgcaaga
+1  ATGGAAGATG CAAAAATAT TATGCACGGT CCACGCCAT TCTATCCTTT

```

Figure 3. The 5′-flanking nucleotide sequence of the luciferase gene from *Nyctophila caucasica*. Nucleotides are numbered from the translation initiator ATG (bold) with A being position +1. The four putative TATA boxes at positions -63, -144, -166 and -190 and the CCAAT boxes at position -159 and -236 are indicated (underlined). The consensus binding sequences of transcription factor GATA elements are presented double underlined. Also shown is the 5′ end of the *Lampyrus noctiluca* cDNA end product indicated by italicised bases (GenBank accession number X89479).

Korean origin (Table 2). This suggests the possibility of cryptic speciation and/or population isolation and highlights the importance of the luciferase gene, especially the intron sequences, as an informative marker for species determination and population genetic studies.

The 810 bases upstream of the *N. caucasica* luciferase gene were examined for putative promoter sites. Four TATA boxes, two CCAAT boxes and two GATA motifs were identified in the upstream region (Figure 3). Despite extensive sequence differences, comparisons with *P. pyralis* luciferase flanking sequence revealed that three motifs were conserved in both species, two TATA boxes at positions -190 and -166 (positions refer to *N. caucasica* sequence) along with a conserved CCAAT box at position -159 (Figure 4). This suggests the presence of a putative core promoter conserved in Lampyrinae from -190 through to -155. It was not possible to infer a similar core promoter region in members of the Lucolinae due to insufficient identity with *Luciola lateralis* upstream flanking sequence (GeneBank accession numbers U49182 and U51019).

The deduced amino acid sequences of the luciferase gene of *N. caucasica* showed 98.9% homology to that of *L. noctiluca* (Table 3). Phylogenetic analysis with other bioluminescent beetle luciferases further confirmed that the

deduced amino acid sequences of the *N. caucasica* luciferase gene belonged to the subfamily Lampyrinae (Figure 5). Furthermore, with both *Lampyrus* and *Nyctophila* along with *Pyrocoelia* species belonging to Lampyrini the luciferase molecular data supports the taxonomic classification of these species down to the tribe level. However, the most identity shared was with *Lampyrus turkistanicus*, both at a nucleotide level and an amino acid sequence level, 0.992 and 0.998 respectively, that was greater than the sequence identity with species of its own genus (0.981 and 0.987). The strength of the nucleotide sequence identity, 0.992, suggests a misidentification of *L. turkistanicus*.

Table 3. Pairwise identity matrix for the deduced amino acid sequences (above) and CDS (below) among *Nyctophila caucasica* and selected other beetle luciferases. GenBank accession numbers are given in the Materials and Methods.

		Ncau1	Ncau2	Lturk	Lnoc	Ppyr
Ncau1	-	1	0.998	0.989	0.843	
Ncau2	0.999	-	0.998	0.989	0.843	
Lturk		0.992	0.992	-	0.987	0.841
Lnoc		0.984	0.984	0.981	-	0.84
Ppyr		0.771	0.771	0.771	0.768	-

To date, structural and biochemical studies of beetle luciferin have concentrated on that of *Photinus pyralis*. To our knowledge, luciferin has only been characterized from one other lampyrid, *L. turkistanicus* (Hadj-Mohammadi and Chaichi

```

nyct tgaagacaaaacggtttgctaaaag-atttaaaaagaaagttaaagtat -255
phot ctgc...a.t...tagg.a...gcccg...gtg..a.gt.gcc..a--- -257

nyct ctcaaaaattcggtaatggcaataaagttgctgaagttagttgtaaacgc -205
phot .c.t.....t..c...ta.....aa...a.a.t.g.c....t. -207

nyct gctaagatttttat tatatacattttatgtc-ctcatgttt tataaaagcaa -156
phot a.a..C..... ..........ta..ag..g...C..... ..... -157

nyct tatttacgtggta--taaactgtcaataaaaatttaagtaatgtgattac -108
phot .....aa.c...aac..caaa.a.....acg.....a -107

nyct gagtcaaaggctctttcacaanaaggtatttaagatacataatcatataat -58
phot ...c.....c.ct.g.....ca..gg...tcct.tg. -57

nyct gtgacaattcgagaataactggttcctgagacactaacgcgctaataatcat -8
phot ..t...-...tt...gtc.c.cg.a.t...t.g.att.cgg..ctg. -8

nyct tgcaagaATGGAAAGATGCAAAAAATATTATGCACGGTCCACCGCCATTCT 43
phot ..gt.a.....C..C.....C..A.A.A.A..C..GG..... 43

```

Figure 4. The 5'-flanking nucleotide sequence of the luciferase gene from *Nyctophila caucasica* aligned with orthologous sequence from *Photinus pyralis* (GeneBank accession # M15077). Nucleotides are numbered from the translation initiator ATG (bold) with A being position +1. Two putative TATA boxes and a CCAAT box conserved in both sequences are highlighted in grey.




Figure 5. A phylogenetic tree based upon amino acid sequences of the *Nyctophila caucasica* luciferase and fifteen known beetle luciferases. The maximum parsimony tree was obtained by a heuristic search with 1000 bootstrap replicates. Branch numbers refer to bootstrap values.

1996). Recently the luciferase mRNA (Alipour et al 2004) has been studied providing both enzyme and substrate information comparable to that of *P. pyralis*. However, the lampyrid species that the luciferin was characterized from along with the luciferase mRNA may be in doubt. The high DNA sequence identity of *L. turkistanicus* luciferase to the luciferase gene sequence from *N. caucasica* strongly suggests the possibility of taxonomic confusion. Furthermore, the specimens used in this current study were those obtained from the same forest area that provided specimens for both the luciferase mRNA and luciferin characterisation of *L. turkistanicus*. These data combined with the fact that *L. turkistanicus* has never been reported in Iran (pers. comm. M. Geisthardt) provides strong evidence that *N. caucasica* has, in the past, been misidentified as *L. turkistanicus*. With time we hope to resolve this issue and will eventually provide a bioluminescent system characterized at the

luciferin and luciferase level to the same extent as that of *P. pyralis*.

Acknowledgements

We are grateful to Michael Giesthardt for both the identification of male specimens of *N. caucasica* and invaluable contributions to both the work and the manuscript preparation.

References

- Alipour BS, Hosseinkhani S, Nikkhah M, Naderi-Manesh H, Chaichi MJ, Osaloo SK. 2004. Molecular cloning, sequence analysis, and expression of a cDNA encoding the luciferase from the glow-worm, *Lampyris turkistanicus*. *Biochemical and Biophysical Research Communications* 325: 215-222.
- Cho KH, Lee JS, Choi YD, Boo KS. 1999. Structural polymorphism of the luciferase gene in the firefly, *Luciola lateralis*. *Insect Molecular Biology* 8: 193-200.

- Choi YS, Lee KS, Bae JS, Lee KM, Kim SR, Kim I, Lee SM, Sohn HD, Jin BR. 2002. Molecular cloning and expression of a cDNA encoding the luciferase from the firefly, *Hotaria unimunsana*. *Comparative Biochemistry and Physiology. Biochemistry and Molecular Biology* 132: 661-670.
- Conti E, Franks NP, Brick P. 1996. Crystal structure of firefly luciferase throws light on a superfamily of adenylate-forming enzymes. *Structure* 4: 287-298.
- De Wet JR, Wood KV, Helinski DR, Deluca M. 1985. Cloning of firefly luciferase cDNA and the expression of active luciferase in *Escherichia coli*. *Proceedings of the National Academy of Sciences of the United States of America* 82: 7870-7873.
- Geisthardt M, Satô M. (in press) Lampyridae. In Löbl I. and Smetana A. (eds): *Catalogue of Palaearctic Coleoptera* Apollo Books, Stenstrup.
- Hadj-mohammadi MR, Chaichi MJ. 1996. Separation, identification and determination of luciferin in the Iranian firefly, *Lampyrus turkestanicus* by HPLC and spectroscopic methods. *Photochemistry and Photobiology* 64: 821-822.
- Li J, Choi YS, Feng Z, Kim I, Lee SM, Kim JG, Kim KY, Sohn HD, Jin BR. 2003a. Genomic structure of the luciferase gene and phylogenetic analysis of the firefly, *Pyrocoelia rufa*. *International Journal of Industrial Entomology* 7: 181-189.
- Li J, Choi YS, Lee SC, Lee SM, Kim JG, Kim I, Kim KY, Sohn HD, Jin BR. 2003b. Genomic structure of the luciferase gene of the firefly, *Lampyrus noctiluca*. *International Journal of Industrial Entomology* 7: 155-159.
- McElroy WD, Deluca MA. 1983. Firefly and bacterial luminescence: basic science and applications. *Journal of Applied Biochemistry* 5: 197-209.
- Oba Y, Ojika M, Inouye S. 2004. Characterization of CG6178 gene product with high sequence similarity to firefly luciferase in *Drosophila melanogaster*. *Gene* 329: 137-145.
- Olivier E. 1884. Essai d'une Révision des espèces européennes and circaméditerranéennes de la famille des Lampyrides. *L'Abeille* 22: 1-55 and 1-4 (Notes compl.).
- Sala-Newby GB, Thomson CM, Campbell AK. 1996. Sequence and biochemical similarities between the luciferases of the glow-worm *Lampyrus noctiluca* and the firefly *Photinus pyralis*. *Biochemical Journal* 313: 761-767.
- Swofford DL. 2001. *PAUP*. Phylogenetic Analysis Using Parsimony (*and other Methods)* Version 4. Sinauer Associates, Sunderland, Massachusetts.
- Wood KV. 1995. The chemical mechanism and evolutionary development of beetle bioluminescence. *Photochemistry and Photobiology* 62: 662-673.