



## **Stimulation by Proopiomelanocortin-Derived Peptides of LH Release by Bullfrog Dispersed Anterior Pituitary Cells**

Authors: Aida, Tomomi, Oguchi, Atsuko, Iwamuro, Shawichi, Yamamoto, Kazutoshs, and Kikuyama, Sakae

Source: Zoological Science, 14(5) : 827-831

Published By: Zoological Society of Japan

URL: <https://doi.org/10.2108/zsj.14.827>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

# Stimulation by Proopiomelanocortin-Derived Peptides of LH Release by Bullfrog Dispersed Anterior Pituitary Cells

Tomomi Aida, Atsuko Oguchi, Shawichi Iwamuro,  
Kazutoshi Yamamoto and Sakae Kikuyama\*

*Department of Biology, School of Education, Waseda University, Tokyo 169-50, Japan*

**ABSTRACT**—Labeling and immunoprecipitation experiments confirmed that three proopiomelanocortin (POMC)-derived peptides, the N-terminal peptide of POMC (NPP), joining peptide (JP) and adrenocorticotrophic hormone (ACTH), were released by the bullfrog (*Rana catesbeiana*) anterior pituitary. The effects of these three peptides on luteinizing hormone (LH) release by bullfrog dispersed anterior pituitary cells were studied. NPP and ACTH, but not JP, enhanced LH release concentration-dependently. Approximately 6 hr elapsed before the gonadotrophs responded to NPP and ACTH by releasing LH, whereas their response to human GnRH (hGnRH) was faster, suggesting that the modes of action of these two peptides and hGnRH differ. These results raise the possibility that NPP and ACTH act as paracrine factors in the bullfrog pituitary to enhance LH release either directly or indirectly.

## INTRODUCTION

It has been established that post translational processing of proopiomelanocortin (POMC) generates several biologically active peptides, as well as peptides with biological functions that are obscure (Hanneman *et al.*, 1989). We isolated substantial amounts of the joining peptide (JP) and N-terminal peptide of POMC that does not contain  $\gamma$ -melanophore-stimulating hormone (NPP) from the anterior lobe of the bullfrog pituitary (Iwamuro *et al.*, 1992), which suggested that they are products of the anterior pituitary cells, presumably the corticotrophs. These two peptides have been generally regarded as products of the intermediate, not the anterior, lobe (Eipper and Mains, 1980). The NPP thus obtained was demonstrated to have a stimulatory effect on the frog interrenal tissue (Iwamuro *et al.*, 1992). Elevation of adrenal function by NPP has also been reported in mammals (Estivariz *et al.*, 1982; Lis *et al.*, 1981) and teleosts (Takahashi *et al.*, 1985). The biological function of JP, however, remains to be determined.

Evidence is growing that, in mammals, different types of anterior pituitary cell interact with each other by producing paracrine factors (Denef *et al.*, 1989; Schwartz and Cherney, 1992). It has also been suggested that pituitary hormones act as paracrine factors in amphibians. We found in the bullfrog that prolactin enhanced gonadotropin-releasing hormone (GnRH)-induced luteinizing hormone (LH) release (Oguchi *et al.*, 1997) and that the  $\alpha$ -subunit of glycoprotein hormones stimulated the prolactin release (Oguchi *et al.*, 1996).

As a step toward elucidating further functions of POMC-derived peptides, we examined the effects of NPP, JP and adrenocorticotrophic hormone (ACTH) on LH release by bullfrog dispersed anterior pituitary cells and found that NPP and ACTH, but not JP, enhanced LH release.

## MATERIALS AND METHODS

### Hormones and antisera

Bullfrog NPP (fNPP) and JP (fJP) consisting of 48 and 35 amino acids, respectively, were purified as described by Iwamuro *et al.* (1992). Bullfrog ACTH (fACTH), a 39 amino acid polypeptide, was prepared by solid-phase chemistry (Taker Chaise Co. Ltd., Shigeo, Japan), according to the amino acid sequence deduced from bullfrog POMC cDNA (Aida *et al.*, 1994). Human GnRH (hGnRH) and ovine corticotropin-releasing factor (oCRF) were purchased from the Peptide Institute (Osaka, Japan). Polyclonal anti-fACTH and anti-fJP antisera were raised in rabbits as described by Tanaka *et al.* (1992) and Iwamuro *et al.* (1993), respectively, and an anti-fNPP antiserum was raised in a female albino rabbit by the lymph node injection technique (Goudie *et al.*, 1966). Before using the latter antiserum, its specificity was confirmed by immunoblotting and immunohistochemical staining. Briefly, Western blot analysis revealed that the anti-fNPP antiserum recognized purified fNPP, but neither fJP nor fACTH, and immunostaining of the bullfrog pituitary revealed that it stained the cells similar to those stained with anti-fACTH antiserum.

### Labeling and immunoprecipitation

Two distal pituitary lobes of adult bullfrogs were cut into small pieces and incubated with  $^{35}\text{S}$ -methionine (Amersham, England) in methionine-free 70% Dulbecco's modified Eagle's medium (Gibco, Detroit, MI) containing 0.1% BSA (Fraction V, Sigma, St. Louis, MO) and oCRF ( $10^{-7}$  M) for 5 hr at 25°C. The medium was decanted, diluted with immunoprecipitation buffer (10 mM Tris-HCl, 0.1% SDS, 0.1% Triton X-100, 2 mM EDTA, pH 7.4) and immunoprecipitated with the anti-fNPP, anti-fJP or anti-fACTH antiserum, as described previously (Miura *et al.*, 1982). The immunoprecipitates were sepa-

\* Corresponding author: Tel. +81-3-5286-1517;  
FAX. +81-3-3207-9694.

rated electrophoretically on a 15-25% SDS-polyacrylamide gradient slab gel (Daiichi Pure Chem. Co., Tokyo, Japan), according to the method of Laemmli (1970), using Tris-tricine buffer (pH 8.3), and then visualized fluorographically (Bonner and Laskey, 1974). The individual fNPP, fJP and fACTH preparations were subjected separately to SDS-polyacrylamide gel electrophoresis in the same manner and each band was stained with Coomassie brilliant blue R (Fluka Chemicals, Germany). Molecular marker used in this experiment was the product of Daiichi Pure Chem. Co. (Tokyo, Japan).

#### Cell incubation and hormone assay

Adult male bullfrog dispersed anterior pituitary cells were prepared as described previously (Oguchi *et al.*, 1996), and suspended in 70% Eagle's minimum essential medium (MEM) without phenol red (Nissui Pharmaceutical, Tokyo, Japan) containing 0.1% BSA and 20 mM HEPES (Sigma, St. Louis, MO). The volume of the suspension was adjusted with this medium so that 1 ml contained  $8 \times 10^4$  cells, and  $2 \times 10^4$  cells in 250  $\mu$ l medium were plated in each well of a 96-multiwell plate (Corning Inc., NY). Preincubation was continued for 24 hr unless otherwise stated. After preincubation, the medium was replaced with MEM containing the required test substance, fNPP, fJP, fACTH or hGnRH, and the 96-multiwell plate was incubated at 25°C in a humidified atmosphere of 95% air-5% CO<sub>2</sub> for various times. The medium was collected from each well, centrifuged at 800 rpm for 5 min and the supernatant was subjected to radioimmunoassay for bullfrog LH (fLH) (Oguchi *et al.*, 1997).

#### Statistics

The data were expressed as means  $\pm$  SEM and analyzed using analysis of variance and Student's *t*-test. Differences at  $P < 0.05$  were considered significant.


## RESULTS

In order to confirm that the anterior pituitary of the bullfrog actually released the POMC-derived peptides that were to be tested for biological activity, immunoprecipitation analysis was performed. As shown in Fig. 1, each antiserum precipitated polypeptides of varying molecular sizes. The peptides that migrated to 4-5-kDa positions were concluded to be fNPP, fJP and fACTH, as electrophoretic bands of purified fNPP and fJP and synthetic fACTH appeared at comparable positions (data not shown).


The amounts of immunoassayable LH released by the dispersed pars distalis cells into the culture media containing hGnRH, fNPP, fJP or fACTH were followed for 24 hr. As shown in Fig. 2, hGnRH stimulated LH release markedly during the initial 6 hr of incubation and moderately during the last 18 hr of the incubation period. Slight, but non-significant, enhancement by fNPP and fACTH of LH release was observed after incubation for 6 hr, followed by marked and significant elevation during the next 6 hr, whereas, no stimulation of LH release by fJP was observed throughout the 24 hr incubation period.

As shown in Fig. 3, fNPP and fACTH both enhanced LH release by the dispersed pituitary cells in a concentration-dependent manner, whereas, fJP, at the concentrations examined ( $10^{-12}$ - $10^{-7}$  M), did not stimulate LH release (data not shown). The minimum effective concentrations of fNPP and fACTH were approximately  $10^{-11}$  and  $10^{-12}$  M, respectively.


Experiments were conducted to ascertain whether it took



**Fig. 1.** Immunoprecipitation analysis of the POMC-derived peptides released by bullfrog anterior lobes. Anterior pituitaries were incubated with <sup>35</sup>S-methionine in the presence of  $10^{-7}$  M oCRF and the media were subjected to immunoprecipitation with anti-fNPP, anti-fJP or anti-fACTH antisera, followed by SDS-PAGE and fluorography. Lanes 1, 2 and 3: immunoprecipitates with anti-fNPP; anti-fJP; and anti-fACTH, respectively.



**Fig. 2.** Time courses of fNPP-, fJP- and fACTH-induced LH release by bullfrog dispersed pituitary cells. The concentrations of each test substance used was  $10^{-7}$  M, and hGnRH ( $10^{-6}$  M) was used as a positive control. After preincubation for 24 hr, incubation was continued for 3, 6, 12 or 24 hr. ●, control; ▲, fNPP; △, fJP; ○, fACTH; ■, hGnRH. The values are means  $\pm$  SEM ( $n = 6$ ). \*Significantly different from the control values ( $P < 0.001$ ).



**Fig. 3.** Effects of various concentrations of fNPP (A) and fACTH (B) on LH release by bullfrog dispersed pituitary cells. After preincubation for 24 hr, the cells were incubated for 12 hr. The values are means  $\pm$  SEM (n=6). \*Significantly different from the control values ( $P < 0.001$ ).

**Table 1.** Effects of preincubation and/or incubation with fNPP and fACTH on LH release by dispersed pituitary cells

Treatment		LH (ng/10 <sup>4</sup> cells)
Preincubation	Incubation	
None	None	12.2 $\pm$ 1.6 <sup>a</sup>
None	fNPP	13.8 $\pm$ 2.9 <sup>a</sup>
fNPP	None	50.8 $\pm$ 0.6 <sup>b</sup>
fNPP	fNPP	67.4 $\pm$ 3.1 <sup>c</sup>
None	fACTH	14.0 $\pm$ 2.2 <sup>a</sup>
fACTH	None	54.5 $\pm$ 3.3 <sup>b</sup>
fACTH	fACTH	74.6 $\pm$ 3.1 <sup>c</sup>

The cells were preincubated for 6 hr and incubated for 6 hr with various combinations of fNPP ( $10^{-7}$  M), fACTH ( $10^{-7}$  M) or medium alone and the amounts of LH released during the 6-hr incubation period were determined by RIA. The values represent the means  $\pm$  SEM of 6 determinations and those with the same superscripts do not differ significantly from each other ( $P > 0.05$ ).

the cells longer to respond to the POMC-derived peptides than to hGnRH. The results summarized in Table 1 indicate there was a delay before the cells responded to fNPP and fACTH and started to release amounts of LH above the control levels. Cells preincubated with fNPP or fACTH and then incubated with medium only, released significantly more LH than cells preincubated with medium only and then incubated with fNPP or fACTH. Less LH was released by cells preincubated with fNPP or fACTH than by cells preincubated and incubated with these peptides, but the differences were relatively small.

## DISCUSSION

The results of this study demonstrated that fNPP and fACTH, but not fJP, stimulated LH release by bullfrog dispersed anterior pituitary cells in culture. It is generally accepted that

proteolytic cleavage of the N-terminal portion of POMC in the intermediate lobe of the pituitary generates NPP,  $\gamma$ -melanophore-stimulating hormone and JP and that the N-terminal portion of POMC is not processed in the anterior lobe (Eipper and Mains, 1980). In a previous study, we isolated substantial amounts of NPP and JP from the bullfrog anterior pituitary (Iwamuro *et al.*, 1992), which suggested that the N-terminal fragment of POMC is processed, at least, partially in the frog distal lobe to generate JP and NPP. This was confirmed by our labeling and immunoprecipitation experiments, which showed that bullfrog dispersed anterior lobe cells released NPP-, JP- and ACTH-like peptides into the culture medium.

NPP and ACTH stimulated LH release by dispersed pituitary cells and their respective minimum effective concentrations were  $10^{-11}$  and  $10^{-12}$  M. As the circulating concentrations of ACTH in the frog, *Rana ridibunda*, have been reported to be around  $8 \times 10^{-12}$  M (Vaudry *et al.*, 1975), it is possible that the released POMC-derived peptides stimulate LH release *in vivo* by acting on the neighboring gonadotrophs.

It is well known that considerable transformation of the anuran pituitary structure occurs during metamorphosis. Before metamorphosis, the dorsal side of the entire anterior pituitary is attached to the median eminence. As metamorphosis proceeds, the caudal part of the pars distalis gradually becomes detached from the median eminence and finally, is connected to the median eminence only at the rostral part by the portal vessel (Etkin *et al.*, 1965). It is noteworthy that bullfrog immunoreactive ACTH cells are localized in the rostral region (Tanaka *et al.*, 1992), which is considered to be situated upstream of the pituitary circulation. Therefore, it is quite possible that POMC-derived peptides released by ACTH cells reach the LH cells, which are distributed in the dorso-caudal region of the adenohypophysis (Tanaka *et al.*, 1991, 1992), via the pituitary circulation and enhance LH release.

We observed marked enhancement of LH release by hGnRH within 6 hr of culture, whereas it took 12 hr for fNPP and fACTH to enhance LH release indicating that the modes of action of hGnRH and these two POMC-derived peptides differ. However, whether fNPP and fACTH act indirectly on LH cells remains to be elucidated.

From our concentration-response data, the LH-releasing activity of purified fNPP was estimated to be approximately one-tenth of that of synthetic fACTH. In a previous study, we measured the amount of ACTH in the fNPP material using a radioimmunoassay (RIA) that crossreacted fully with fACTH (Jégou *et al.*, 1983; Vaudry *et al.*, 1984) and found that the residual amount of ACTH comprised less than 0.005% of the purified fNPP (Iwamuro *et al.*, 1992). Accordingly, the activity of the NPP preparation can not be accounted for by contamination with ACTH and fNPP is considered to possess intrinsic LH-releasing activity.

Bullfrog NPP is known to have a relatively high sequence homologies with NPPs of other vertebrates (Iwamuro *et al.*, 1992): it shows 100, 85 and 50% sequence homology with *Rana ridibunda* (Hilario *et al.*, 1990), *Xenopus laevis* (Martens, 1987) and human (Seidah *et al.*, 1981) NPPs, respectively. The amino acid sequences of human (Cochet *et al.*, 1982), monkey (Patel *et al.*, 1988), bovine (Nakanishi *et al.*, 1979), porcine (Boileau *et al.*, 1983), rat (Drouin *et al.*, 1985), mouse (Uhler and Herbert, 1983) and salmon (Kitahara *et al.*, 1988) JPs, deduced from their cDNA sequences, are highly variable, indicating that the structure of JP has diverged considerably during evolution. In fact, fJP exhibits 86 and 54% homology with *Rana ridibunda* (Hilario *et al.*, 1990) and *Xenopus laevis* (Martens, 1987) JPs, but only 9% homology with human JP (Chang *et al.*, 1980). In our study, fJP did not show LH-releasing activity, whereas fNPP and fACTH did. The lack of phylogenetic structural conservation of JP might explain its lack of bioactivity. In contrast, the fact that the amino acid sequences of NPP are conserved throughout the vertebrate phylum suggests that this peptide may play important roles. In fact, NPP has been shown to be involved in the regulation of adrenocortical activity in the rat (Lis *et al.*, 1981), rainbow trout (*Salmo gairdneri*) (Takahashi *et al.*, 1985) and frog (*Rana ridibunda*) (Iwamuro *et al.*, 1992). Recently, the structure of the N-terminal portion (26 amino acids) of POMC was reported to be responsible for directing POMC to the regulatory pathway (Cool *et al.*, 1995).

In conclusion, our data show that the NPP and ACTH stimulate LH release by the bullfrog anterior pituitary and raise the possibility that these peptides act on the gonadotrophs in a paracrine fashion.

#### ACKNOWLEDGMENTS

This work was supported by a grant from Waseda University. The authors thank Drs. K. Wakabayashi, Y. Hanaoka and S. Tanaka of Gunma University for the gift of the goat anti-rabbit IgG-serum (HAC-RBA2-05GTP91) and the fLH RIA kit.

#### REFERENCES

- Aida T, Iwamuro S, Kikuyama S, Miura S (1994) Cloning of a bullfrog proopiomelanocortin cDNA. *Zool Sci* 11 (Suppl): 8 (Abstract)
- Boileau G, Barbeau C, Jeannotte L, Chrétien M, Drouin J (1983) Complete structure of the porcine proopiomelanocortin mRNA derived from the nucleotide sequence of cloned cDNA. *Nucleic Acids Res* 11: 8063–8071
- Bonner WM, Laskey RA (1974) A film detection method for tritium-labelled proteins and nucleic acids in polyacrylamide gels. *Eur J Biochem* 46: 83–88
- Chang ACY, Cochet M, Cohen SN (1980) Structural organization of human genomic DNA encoding the proopiomelanocortin peptide. *Proc Natl Acad Sci USA* 77: 4890–4894
- Cochet M, Chang ACY, Cohen SN (1982) Characterization of the structural gene and putative 5'-regulatory sequences for human proopiomelanocortin. *Nature* 297: 335–338
- Cool DR, Fenger M, Snell CR, Loh YP (1995) Identification of the sorting signal motif within proopiomelanocortin for the regulated secretory pathway. *J Biol Chem* 270: 8723–8729
- Denef C, Maertens P, Allaerts W, Mignon A (1989) Cell-to-cell communication in peptide target cells of the anterior pituitary. *Methods Enzymol* 168: 47–71
- Drouin J, Chamberland M, Charron J, Jeannotte L, Nemer M (1985) Structure of the rat proopiomelanocortin (POND) gene. *FEBS Lett* 193: 54–58
- Eipper BA, Mains RE (1980) Structure and biosynthesis of proadrenocorticotropin/endorphin and related peptides. *Endocrine Rev* 1: 1–27
- Estivariz FE, Iturriza F, McLean C, Hope J, Lowry PJ (1982) Stimulation of adrenal mitogenesis by N-terminal proopiomelanocortin peptides. *Nature* 297: 419–422
- Etkin W, Kikuyama S, Rosenbluth J (1965) Thyroid feedback to the hypothalamic neurosecretory system in frog larvae. *Neuroendocrinology* 1: 45–64
- Goudie RB, Horne CHW, Wilkinson PC (1966) A simple method for producing antibody specific to a single selected diffusible antigen. *Lancet* 2: 1224–1226
- Hanneman E, Bunzow J, Salon J, Grandy D, Albert P, Machida C, Civelli O (1989) Peptides encoded by the proopiomelanocortin gene. In "Peptide Hormones as Prohormones: Processing, Biological Activity, Pharmacology" Ed by J Martinez, Ellis Horwood Limited, New York, pp 53–82
- Hilario E, Lihmann I, Vaudry H (1990) Characterization of the cDNA encoding proopiomelanocortin in the frog *Rana ridibunda*. *Biochem Biophys Res Commun* 173: 653–659
- Iwamuro S, Hayashi H, Delbende C, Vaudry H, Kikuyama S (1992) Purification and characterization of joining peptide and N-terminal peptide of pro-opiomelanocortin from the pars distalis of the bullfrog pituitary. *Peptides* 13: 729–735
- Iwamuro S, Tanaka S, Sugihara S, Nomizu M, Hayashi H, Kikuyama S (1993) Secretion of immunoreactive joining peptide by the pituitary of the bullfrog *Rana catesbeiana*. *Ann NY Acad Sci* 680: 552–554
- Jégou S, Tonon MC, Leroux P, Delarue C, Leboulenger F, Pelletier G, Côté J, Ling N, Vaudry H (1983) Immunological characterization of endorphins, adrenocorticotropin and melanotropins in frog hypothalamus. *Gen Comp Endocrinol* 51: 246–254
- Kitahara N, Nishizawa T, Aida K, Okazaki H, Andoh T, Soma GI (1988) Absence of a gamma-melanocyte-stimulating-hormone sequence in proopiomelanocortin mRNA in chum salmon *Oncorhynchus keta*. *Comp Biochem Physiol* 91B: 365–370
- Laemmli UK (1970) Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* 227: 680–685
- Lis M, Hamet P, Gutkowska J, Maurice G, Seidah NG, Larivier N, Chrétien M, Genest J (1981) Effect of N-terminal portion of proopiomelanocortin on aldosterone release by human adrenal

- adenoma *in vitro*. J Clin Endocrinol Metab 52: 1053–1058
- Martens GJM (1987) Structural organization of the proopiomelanocortin gene in *Xenopus laevis*. 5'-end homologies within the toad and mammalian genes. Eur J Biochem 165: 467–472
- Miura S, Mori M, Amaya Y, Tatibana M (1982) A mitochondrial protease that cleaves the precursor of ornithine carbamoyltransferase. Purification and properties. Eur J Biochem 122: 641–647
- Nakanishi S, Inoue A, Kita T, Nakamura M, Chang ACY, Cohen SN, Numa S (1979) Nucleotide sequence of cloned cDNA for bovine corticotropin-beta-lipotropin precursor. Nature 287: 423–427
- Oguchi A, Tanaka S, Yamamoto K, Kikuyama S (1996) Release of  $\alpha$ -subunit of glycoprotein hormones from the bullfrog pituitary: Possible effect of  $\alpha$ -subunit on prolactin cell function. Gen Comp Endocrinol 102: 141–146
- Oguchi A, Tanaka S, Aida T, Yamamoto K, Kikuyama S (1997) Enhancement by prolactin of the GnRH-induced release of LH from dispersed anterior pituitary cells of the bullfrog (*Rana catesbeiana*). Gen Comp Endocrinol 107: 128–135
- Patel PG, Sherman TG, Watson SJ (1988) Characterization of proopiomelanocortin cDNA from Old World monkey, *Macaca nemestrina*. DNA 7: 627–635
- Schwartz J, Cherney R (1992) Intercellular communication within the anterior pituitary influencing the secretion of hypophysial hormones. Endocr Rev 13: 453–475
- Seidah NG, Rochemont J, Hamelin J, Benjannet S, Chrétien M (1981) The missing of the pro-sequence of human proopiomelanocortin: Sequence and evidence for C-terminal amidation. Biochem Biophys Res Commun 102: 710–716
- Takahashi A, Kubota J, Kawauchi H, Hirano T (1985) Effects of N-terminal peptide of salmon proopiomelanocortin on interrenal function of the rainbow trout. Gen Comp Endocrinol 58: 328–335
- Tanaka S, Sakai M, Park M, Kurosumi K (1991) Differential appearance of the subunits of glycoprotein hormones (LH, FSH and TSH) in the pituitary of bullfrog (*Rana catesbeiana*) larvae during metamorphosis. Gen Comp Endocrinol 84: 318–327
- Tanaka S, Mizutani F, Yamamoto K, Kikuyama S, Kurosumi K (1992) The alpha-subunit of glycoprotein hormones exists in the prolactin secretory granules of the bullfrog (*Rana catesbeiana*) pituitary gland. Cell Tissue Res 267: 223–231
- Uhler M, Herbert E (1983) Complete amino acid sequence of mouse proopiomelanocortin derived from the nucleotide sequence of proopiomelanocortin cDNA. J Biol Chem 258: 257–261
- Vaudry H, Vague P, Dupont W, Leboulenger F, Vaillant R (1975) A radioimmunoassay for plasma corticotropin in frogs (*Rana esculenta* L.). Gen Comp Endocrinol 25: 313–322
- Vaudry H, Jenks BG, Van Overbeeke AP (1984) Biosynthesis, processing and release of proopiomelanocortin related peptides in the intermediate lobe of the pituitary gland of the frog (*Rana ridibunda*). Peptides 5: 905–910

(Received May 12, 1997 / Accepted May 26, 1997)