

Redescription of *Dysdera sultani* (Araneae: Dysderidae) with the first description of the female

Authors: Özkütük, Recep Sulhi, Kaya, Rahşen S., Yağmur, Ersen Aydın, Danışman, Tarık, Gücel, Salih, et al.

Source: Arachnologische Mitteilungen: Arachnology Letters, 57(1) : 21-25

Published By: Arachnologische Gesellschaft e.V.

URL: <https://doi.org/10.30963/aramit5704>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Redescription of *Dysdera sultani* (Araneae: Dysderidae) with the first description of the female

Recep Sulhi Özkütük, Rahşen S. Kaya, Ersen Aydın Yağmur, Tarık Danişman, Salih Gücel & Kadir Boğaç Kunt



doi: 10.30963/aramit5704

Abstract. During field trips made to investigate the dysderid fauna of Turkey, specimens of *Dysdera sultani* Deeleman-Reinhold, 1988, previously known from males only, were collected. Detailed descriptions and photographs of the copulatory organs of both sexes are given. In addition, photographs of some closely related Anatolian species are included.

Keywords: Anatolia, new record, spider, Turkey, woodlouse hunter

Zusammenfassung. Wiederbeschreibung von *Dysdera sultani* (Araneae: Dysderidae) mit Erstbeschreibung des Weibchens. Bei Exkursionen zur Untersuchung der Dysderiden-Fauna der Türkei wurden Exemplare von *Dysdera sultani* Deeleman-Reinhold, 1988 gefangen, von der bisher nur Männchen bekannt waren. Es werden Beschreibungen und Fotos der Kopulationsorgane beider Geschlechter der Art präsentiert. Ergänzt wird dies durch Fotos nahe verwandter anatolischer Arten.

Members of the spider genus *Dysdera* Latreille, 1804 are nocturnal ground foragers that are commonly found in leaf litter and primarily feed on terrestrial woodlice (Jocqué & Dippenaar-Schoeman 2006). The genus is represented by 260 extant species, mainly distributed in the West Palaearctic, with the exception of the Cosmopolitan *Dysdera crocata* C. L. Koch, 1838.

The Italian arachnologist Pietro Pavesi firstly recorded the genus *Dysdera* from the İstanbul Province of Turkey (Pavesi 1876). The first significant contribution on the genus, however, was given by Nosek (1905), and included the descriptions of *D. argeica* Nosek, 1905, *D. asiatica* Nosek, 1905 and *D. longimandibularis* Nosek, 1905. Until the work of Deeleman-Reinhold & Deeleman (1988), in which eight new species of *Dysdera* were described from Turkey, there were only scattered records. Today, the genus *Dysdera* is represented by 24 species in Turkey, most of which are local endemics (Demir & Seyyar 2017, Varol & Danişman 2018).

Here we provide the first description of the female of *D. sultani* Deeleman-Reinhold, 1988, a species previously known only from males, and only from two localities in Turkey and Greece (Deeleman-Reinhold & Deeleman 1988). Additionally, we provide morphological information on the species.

Material and methods

Specimens were collected in the western central Anatolian region of Turkey using a litter reducer (sifter), pitfall traps and hand collection. Specimens were preserved in 70% ethanol and deposited in the Zoology Museum of the Eskişehir Technical University. Digital images of the copulatory or-

gans were obtained using a Leica DFC295 digital camera attached to a Leica S8AP0 stereomicroscope. Five to fifteen photographs were taken in different focal planes and combined using “CombineZP” image stacking software. All measurements are given in mm. Terminology for measurements follows Chatzaki & Arnedo (2006). Terminology for the copulatory organs was adapted from Deeleman-Reinhold & Deeleman (1988).

Abbreviations

Prosoma and opisthosoma: OL – opisthosomal length, PL – prosoma length, PWmax – maximum prosoma width, PWmin – minimum prosoma width
Eyes: AME – anterior median eyes, PLE – posterior lateral eyes, PME – posterior median eyes, AMEd – diameter of anterior median eye, PLEd – diameter of posterior lateral eye, PMEd – diameter of posterior median eye
Chelicera: ChF – length of cheliceral fang, ChG – length of cheliceral groove, ChL – total length of chelicera (lateral external view)
Legs and palps: Ta – tarsus, Me – metatarsus, Ti – tibia, Pa – patella, Fe – femur, C – coxa, D – dorsal, Pl – prolateral, Rl – retrolateral, Pv – proventral, Rv – retroventral, V – ventral.

Results

Family Dysderidae C. L. Koch, 1837

Genus *Dysdera* Latreille, 1804

Dysdera sultani Deeleman-Reinhold, 1988 (Figs 1–4, 7–12, 13–14)

D. sultani Deeleman-Reinhold, in Deeleman-Reinhold & Deeleman, 1988: 214, Figs 236–238.

Material examined. TURKEY. 5 ♀♀, Konya Province, Akşehir District, Cankurtaran Village (38.26667°N, 31.40000°E), 1530 m a.s.l., 12 May 2009, leg. R. S. Kaya; 1 ♀, Konya Province, Akşehir District, Sultan Mountains (38.28075°N, 31.42944°E), 1566 m a.s.l., 15 May 2012, leg. E. A. Yağmur & R. S. Kaya; 1 ♂, 1 ♀, Eskişehir Province, Odunpazarı District, Yukarı Kalabak Village (39.50917°N, 30.41306°E), 1135 m a.s.l., 2 May 2013, leg. R. S. Özkütük; 2 ♂♂, 1 ♀, İzmir Province, Menderes District, Özdere (38.02122°N, 27.10119°E), 30 m a.s.l., 03 February 2013, leg. T. Danişman; 1 ♂, Uşak Province, Eşme District, c. 2 km east of Hardallı Vilage (38.33694°N,

Recep Sulhi ÖZKÜTÜK, Department of Biology, Faculty of Science, Eskişehir Technical University, TR- 26470 Eskişehir, Turkey; E-mail: sozkutuk@gmail.com
Rahşen S. KAYA, Department of Biology, Faculty of Arts&Science, Uludağ University, TR-16059 Görükle, Bursa, Turkey; E-mail: rahsens@gmail.com
Ersen Aydın YAĞMUR, Alaşehir Vocational School, Celal Bayar University, TR-45600 Alaşehir, Manisa, Turkey; E-mail: ersen.yagmur@gmail.com
Tarık DANIŞMAN, Department of Biology, Faculty of Sciences and Arts, Kırıkkale University, TR-71450 Yahşihan, Kırıkkale, Turkey; E-mail: tarikdani@yahoo.com
Salih GÜCEL, Institute of Environmental Sciences and Herbarium, University of Near East, Nicosia, Cyprus; E-mail: sgucel@hotmail.com
Kadir BOĞAÇ Kunt, Department of Biology, Faculty of Science, Eskişehir Technical University, TR- 26470 Eskişehir, Turkey & Zoological Collection of Cyprus Wildlife Research Institute, Taşkent, Kyrenia, Cyprus; E-mail: chaetopelma@gmail.com

submitted 22.11.2018, accepted 31.1.2019, online 20.2.2019

28.82028°E), 935 m a.s.l., 20 January–20 May 2014, leg. E. A. Yağmur & S. Örgel; 4 ♀♀, Uşak Province, Altıntaş Village (38.71778°N, 29.50750°E), 970 m a.s.l., 26 March 2015, leg. E. A. Yağmur, S. Örgel; 1 ♂, 2 ♀♀, Uşak Province, İlyaslı Village (38.60472°N, 29.20556°E), 780 m a.s.l., 25 April 2017, leg. S. Gücel & K.B. Kunt.

Other material examined. *Dysdera argaieca* Nosek, 1905: TURKEY. 3 ♂♂, 2 ♀♀, Kayseri Province, Mount Lifos (38.58666°N, 35.48638°E), 2200 m a.s.l., 19 May 2011, leg. T. Danişman; 2 ♂♂, 2 ♀♀, Kayseri Province, Hisarcık District, Mount Erciyes (38.60808°N, 35.51361°E), 1820 m a.s.l., 18 August–27 October 2012, leg. E. A. Yağmur & H. Koç (Figs 5, 15).

***Dysdera enguriensis* Deeleman-Reinhold, 1988:** TURKEY. 1 ♂, Ankara Province, Çankaya District, Dodurga Village (39.82025°N, 32.66791°E), 1090 m a.s.l., 29 May 2012, leg. K. B. Kunt (Fig. 6).

Diagnosis. *Dysdera sultani* is closely related to the Turkish endemic species *D. argaieca*, according to the basic characteristics of the male copulatory organs, such as the cylindrical shape of the bulb, the massive structure of the posterior apophysis, and the development of the internal and external sclerites on the bulb. The most obvious difference between the males of these two species is the angle and shape of the structures towards the distal part of the bulb. The vulvae of *D. sultani* and *D. argaieca* are similar, but the midline of the transverse bar is more flattened in *D. argaieca* as opposed to crescent shaped in *D. sultani*. Additionally, the posterior diverticulum is membranous and well-developed in *D. argaieca*. **Measurements (♂/♀).** OL 4.80/7.60; PL 4.30/4.50; PWmax 3.40/3.40; PWmin 2.20/2.50; AMEd 0.10/0.23; PLEd 0.10/0.19; PMEd 0.10/0.16; ChF 0.71/0.88; ChG 0.73/0.82; ChL 1.30/1.65.

Description. Medium-sized dysderid spiders. Male specimens are slightly darker than females.

Prosoma reddish brown. The cephalic region is lighter than the thoracic region and is angular laterally. The eyes are closely grouped; anterior-median eyes are more distant from each other (Fig. 1). Chelicerae brown. Cheliceral groove with three small teeth. Distance between basal and medial teeth is half of the distance between distal and medial teeth (Fig. 2).

Labium, gnathocoxae and sternum reddish brown and margins blackish brown. Labium reddish brown from the adjacent edge of the sternum to approximately the tip of the sternum; the tip much lighter, orange and concave. The surface of the sternum bright with nearly uniform weak black hairs and brown margins. Setae longer in males.

Pedichel strongly developed, sclerotized and brown dorsally and ventrally. Legs yellowish to light brown, with the first three segments of the anterior legs darker than the other segments. Front legs spineless. Femora of all legs with dorsal spines in both sexes. Leg measurements and spination are given in Tab. 1 and Tab. 2.

Scopulae present on tarsi and metatarsi of all legs. However, the scopulae of the 3rd and 4th legs are well developed and more distinct than the other legs.

Opisthosoma greyish; cylindrical and more swollen in females. The surface covered with short blackish hairs on both the dorsal and ventral sides. Distinct colour difference between the spinnerets and opisthosoma in both sexes. Spinnerets darker brown than opisthosoma and almost the same

Tab. 1: Leg measurements of *Dysdera sultani* (in mm). C – coxa, FE – femur, PA – patella, TI – tibia, ME – metatarsus, TA – tarsus

(♂/♀)	I	II	III	IV
C	1.68 / 1.70	1.40 / 1.50	0.96 / 1.10	1.18 / 1.50
Fe	2.73 / 3.30	2.30 / 3.00	2.25 / 2.50	2.92 / 3.50
Pa	2.19 / 2.20	2.00 / 2.00	1.39 / 1.40	1.73 / 1.75
Ti	2.78 / 2.80	2.64 / 2.70	1.33 / 1.55	2.08 / 2.45
Me	2.89 / 2.80	2.53 / 2.82	1.97 / 2.25	2.47 / 3.00
Ta	0.50 / 0.60	0.60 / 0.60	0.50 / 0.60	0.60 / 0.60

Tab. 2: Leg spination of legs III and IV of *Dysdera sultani*

	III	IV
♂		
Fe	0	1-2 D
Pa	0	0
Ti	1-1-1Rl 1-1 Pl 1Rv 1V 1Pv	1-1 Rl 1-1Pl 1Rv 1V 1Pv
Me	1-1-1Rl 1-1 Pl 1-1 Rv 1Pv	1-1-1 Rl 1-1 Pl 1-1-1 Rv 1-1-1Pv
♀		
Fe	0	2-2 D
Pa	0	0
Ti	1-1-1Rl 1-1 Pl 1Rv 1V 1Pv	1-1-1 Rl 1-1Pl 1Rv 1V 1Pv
Me	1-1-1-1 Rl 1-1 Pl 1-1 Rv 1Pv	1-1-1 Rl 1-1 Pl 1-1-1 Rv 1-1-1Pv

colour as the legs. Epigastral scutum is present and weakly developed. Tracheal spiracles are sclerotized.

The female palp is the same colour as the legs; palpal tarsus with a single-claw, and almost the entire surface is densely covered with thick hairs.



Fig. 1: *Dysdera sultani*. Female, prosoma, arrows indicate the front margins of the head region





Fig. 2: *Dysdera sultani*. Cheliceral teeth




Figs 3-6: *Dysdera sultani*. 3. Bulbus, nearly prolateral view; 4. Ditto, retrolateral view; 5. *Dysdera argaeica*, bulbus, prolateral view; 6. *Dysdera enguriensis*, bulbus, prolateral view. B, basal; C, crest; D, distal; DH, distal haematodocha; IS, internal sclerite; LS, lateral sheet; M, medial; P, posterior apophysis; T, tegulum

In males, palpal segments are brown and darker than in females; dorsal surface of all patellae, tibiae and tarsi covered with dense, relatively long hairs.

Tegulum cylindrical, and sperm ducts distinct (Figs 3-4). The posterior apophysis long, strongly sclerotized and hook-shaped (Figs 3-4, 7, 12). Distal haematodocha clear; triangular from anterior to posterior in retrolateral view (Fig. 4).

The internal sclerite dark brown, broad, rectangular and extending along the distal division (Fig. 4). In SEM photographs, triangular, spiny protrusions arranged like fish-scales are present from the midline to the distal border of the in-




Figs 7-12: *Dysdera sultani*, bulbus. 7. Prolateral view; 8. Nearly prolateral view; 9. Nearly, retrolateral view; 10. Retrolateral view; 11. nearly distal view; 12. nearly posterior view. C, crest; LS, lateral sheet; MA, median apophysis; P, posterior apophysis

ternal sclerite. The lateral sheet long and spine shaped. Crest membranous and comma shaped. Median apophysis short and spine-like (Figs 7-12).

Vulva strongly sclerotized. Spermathecae slightly concave posteriorly. The dorsal arch is the same width as the spermathecae. The midline is sclerotized and membranous on the margins. Transverse bar strongly convex and about twice as long as spermathecae. Except for the parts that are adjacent to the dorsal arch, the transverse bar is rectangular. Posterior diverticulum wide and membranous. A cartilaginous transition zone is present between the posterior diverticulum and transverse bar (Figs 13-14).

Distribution. *Dysdera sultani* was described by Deeleman-Reinhold (in Deeleman-Reinhold & Deeleman 1988) from Turkey and Greece based on male specimens only. The species was previously known from the type locality in the Akşehir



Figs 13–15: *Dysdera sultani*. **13.** Vulva, dorsal view (DA – dorsal arch, PD – posterior diverticulum, S – spermathecal, TB – transverse bar), scale bar 0.2 mm; **14.** Ditto, ventral view, scale bar 0.1 mm; **15.** *Dysdera argaica*, vulva, dorsal view, scale bar 0.1 mm

District of Konya Province in Turkey and from the Samos Island in Greece. Here we have provided newly collected material from central Anatolian and Aegean regions of Turkey (Fig. 16).

Such broad range from the central Anatolian high mountainous region to the coastal Aegean region and a distribution pattern from warmer areas to more hot-dry areas in *D. sultani* may be related to drying that occurred in Anatolia at the end of the Pleistocene, 15000 years ago. As a result of this drying, many species moved to higher altitudes and humid places, whereas drought-tolerant species could remain or disperse to newly dry areas (see Çıplak 2008, Korkmaz et al. 2014). Additionally, the presence of *D. sultani* on Samos Island is likely due to land bridges that appeared in the Pleistocene during the expansion of this species.

Notes. According to the classification of Deeleman-Reinhold & Deeleman (1988), *D. sultani* belongs to the *asiatica* species group which is characterized by following features:

1. The anterior margins of the prosoma are parallel to each other and the thoracic part is slightly narrowed behind the eyes.
2. Chelicerae are shorter than half the length of the prosoma.
3. Variable leg spination.

Dysdera sultani has all characteristics of the *asiatica* species group mentioned above. The presence of a lateral sheet on the male palp (lateral sheet is well developed in *D. sultani* and *D. argaica* while not well developed in *D. enguriensis*) indicates both relationships between the species and similar features of the *asiatica* species group (see Figs 5–6, 15).

Deeleman-Reinhold & Deeleman (1988) compared the holotype and paratype samples of *D. sultani* and emphasized that “the palps are uniform despite the geographical distance of 420 km between the samples”. The authors, however, indicated that the cheliceral sizes and leg spination are slightly different between the samples.



Figs 16: Map of the records of *Dysdera sultani*

Cheliceral size could be related to the particular isopod species *Dysdera* preys upon (see Pollard 1986), but we found no significant differences in cheliceral size among specimens from different localities. Additionally, as Deeleman-Reinhold & Deeleman (1988) mentioned, the number of spines on the legs may show small variations even within the same population.

Acknowledgements

We would like to thank Dr. Halil Koç and Semih Örgel for their help during the field trips. The English of the text was kindly checked by Dr. Sarah Crews (San Francisco, USA).

References

- Chatzaki M & Arnedo MA 2006 Taxonomic revision of the epigeal representatives of the spider subfamily Harpacteinae (Araneae: Dysderidae) on the island of Crete. – *Zootaxa* 1169: 1-32
- Çıplak B 2008 The analogy between interglacial and global warming for the glacial relicts in a refugium: A biogeographic perspective for conservation of Anatolian Orthoptera. In: Fattorini S (Ed) *Insect ecology and conservation*. Research Signpost, Kerala. pp. 135-163
- Deeleman-Reinhold CL & Deeleman PR 1988 Revision des Dysderinae (Araneae, Dysderidae), les espèces méditerranéennes occidentales exceptées. – *Tijdschrift voor Entomologie* 131: 141-269
- Demir H & Seyyar O 2017 Annotated checklist of the spiders of Turkey. – *Munis Entomology & Zoology* 12: 433-469
- Jocqué R & Dippenaar-Schoeman AS 2006 Spider families of the world. Musée Royal de l'Afrique Central, Tervuren. 336 pp.
- Korkmaz E M, Lunt D H, Çıplak B, Değerli N & Başibüyük H 2014 The contribution of Anatolia to European phylogeography: The centre of origin of the meadow grasshopper, *Chorthippus parallelus*. – *Journal of Biogeography* 41: 1793-1805 – doi: [10.1111/jbi.12332](https://doi.org/10.1111/jbi.12332)
- Nosek A 1905 Araneiden, Opilionen und Chernetiden. In: Penther A & Zederbauer E (eds) *Ergebnisse einer naturwissenschaftlichen Reise zum Erdschias-Dagh (Kleinasien)*. – *Annalen des Kaiserlich-Königlichen Naturhistorischen Hofmuseums in Wien* 20: 114-154
- Pavesi P 1876 Gli aracnidi Turchi. – *Atti della Società Italiana di Scienze Naturali* 19: 1-27
- Pollard SD 1986 Prey capture in *Dysdera crocata* (Araneae: Dysderidae), a long fanged spider. – *New Zealand Journal of Zoology* 13: 149-150 – doi: [10.1080/03014223.1986.10422656](https://doi.org/10.1080/03014223.1986.10422656)
- Varol İ & Danışman T 2018 Two new spider species of the family Dysderidae (Araneae) from Turkey. – *Entomological News* 127: 418-425 – doi: [10.3157/021.127.0505](https://doi.org/10.3157/021.127.0505)