

Less is more: eight new synonyms in Mediterranean spiders (Araneae), with a new Pelecopsis species from Tunisia (Linyphiidae)

Authors: Bosmans, Robert, and Hervé, Christoph

Source: Arachnologische Mitteilungen: Arachnology Letters, 61(1) : 58-64

Published By: Arachnologische Gesellschaft e.V.

URL: <https://doi.org/10.30963/aramit6109>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Less is more: eight new synonyms in Mediterranean spiders (Araneae), with a new *Pelecopsis* species from Tunisia (Linyphiidae)

Robert Bosmans & Christoph Hervé



doi: 10.30963/aramit6109

Abstract. The following eight new synonyms are proposed: *Araneus arganicola* Simon, 1909 = *Neoscona subfuscata* (C. L. Koch, 1837) **syn. nov.**; *Drassus nugatorius* Karsch, 1881 = *Odontodrassus mundulus* (O. Pickard-Cambridge, 1881) **syn. nov.**; *Drassus sockniensis* Karsch, 1881 = *Poecilochroa pugnax* (O. Pickard-Cambridge, 1874) **syn. nov.**; *Drassus tarrhunensis* Karsch, 1881 = *Megamyrmaekion caudatum* Reuss, 1834 **syn. nov.**; *Prosthesima kerimi* Pavesi, 1880 = *Zelotes tragicus* (O. Pickard-Cambridge, 1872) **syn. nov.**; *Prosthesima quadridentata* Strand, 1906 = *Scotophaeus quadridentatus* Caporiacco, 1928 = *Setaphis mollis* (O. Pickard-Cambridge, 1874) **syn. nov.**; *Steatoda erigoniformis* (O. Pickard-Cambridge, 1872) = *Euryopis albomaculata* Denis, 1951 **syn. nov.**. The position of *Drassus obscurus* Lucas, 1846 in *Drassodes* is confirmed and the first figures of the female epigyne and vulva are given. The citation of *Dactylopisthes digiticeps* (Simon, 1881) from Tunisia by Pavesi (1884) is a misidentification and concerns *Thaumatoncus indicator* Simon, 1884. A new species is described: *Pelecopsis pavesii* **spec. nov.** from Tunisia, erroneously identified by Pavesi (1884) as *Pelecopsis parumpunctata* (Simon, 1881).

Keywords: Aranidae, Gnaphosidae, North Africa, Theridiidae, type material

Zusammenfassung. Weniger ist mehr: acht neue Synonymien mediterraner Spinnen (Araneae), mit einer neuen Pelecopsis-Art aus Tunesien (Linyphiidae). Folgende acht neue Synonymien werden vorgeschlagen: *Araneus arganicola* Simon, 1909 = *Neoscona subfuscata* (C. L. Koch, 1837) **syn. nov.**; *Drassus nugatorius* Karsch, 1881 = *Odontodrassus mundulus* (O. Pickard-Cambridge, 1881) **syn. nov.**; *Drassus sockniensis* Karsch, 1881 = *Poecilochroa pugnax* (O. Pickard-Cambridge, 1874) **syn. nov.**; *Drassus tarrhunensis* Karsch, 1881 = *Megamyrmaekion caudatum* Reuss, 1834 **syn. nov.**; *Prosthesima kerimi* Pavesi, 1880 = *Zelotes tragicus* (O. Pickard-Cambridge, 1872) **syn. nov.**; *Prosthesima quadridentata* Strand, 1906 = *Scotophaeus quadridentatus* Caporiacco, 1928 = *Setaphis mollis* (O. Pickard-Cambridge, 1874) **syn. nov.**; *Steatoda erigoniformis* (O. Pickard-Cambridge, 1872) = *Euryopis albomaculata* Denis, 1951 **syn. nov.**. Die Zugehörigkeit von *Drassus obscurus* Lucas, 1846 zu *Drassodes* wird bestätigt und Epigyne und Vulva erstmals abgebildet. Die Nennung von *Dactylopisthes digiticeps* (Simon, 1881) aus Tunesien durch Pavesi (1884) beruht auf einer Fehlbestimmung und betrifft *Thaumatoncus indicator* Simon, 1884. Eine neue Art, *Pelecopsis pavesii* **spec. nov.**, wird aus Tunesien beschrieben; sie war von Pavesi (1884) fälschlicherweise als *Pelecopsis parumpunctata* (Simon, 1881) bestimmt worden.

At present 49089 species of spider are known (WSC 2021), and nearly every day new species are described. Species described in previous centuries often remain unstudied. Redescriptions of such taxons are necessary and ultimately increase the knowledge of the arachnofauna. If synonyms are discovered this leads to fewer valid species, but, in this case: less is more.

In the early days of spider systematics, the number of described genera was very limited and authors, all to the best of their knowledge, described species in large genera, such as *Araneus*, *Clubiona*, *Drassus* (= *Drassodes*), *Erigone*, *Linyphia*, *Melanophora* (= *Zelotes*), etc. The definitions and limitations of these genera have changed and improved considerably nowadays. During revisions it is difficult to assess these older species because many of them were never illustrated. The logical thing to do is to examine the type material, if available. Here, species from the Mediterranean region described by Caporiacco (1928), Denis (1951), Karsch (1881), Lucas (1846), Pavesi (1880, 1884) and Simon (1909) are reconsidered. Fortunately, type material from almost all of them was available for study. An examination of the material and original descriptions led to several new synonyms and overall fewer valid species in the region, but also increased our knowledge of the remaining species.

Materials and methods

Species were examined using a Nikon SMZ1270 stereo microscope. Details of male palps and female epigynes were

studied with an Olympus CH-2 microscope with a drawing tube. Photographs were taken with a Moticam 5MP camera attached to a Realux stereoscopic microscope.

Structures of the left palpus are depicted. All morphological measurements are given in millimetres. Somatic morphology measurements were taken using a scale reticule in the eyepiece of the stereo microscope. Measurements of the legs are taken from the dorsal side.

Male palps were detached and transferred to glycerol for examination under the microscope. Female genitalia were excised using sharpened needles and then transferred to clove oil for examination under the microscope. Later, palps and epigynes were returned to 70% ethanol.

Abbreviations

AME: anterior median eyes; ALE: anterior lateral eyes; d: dorsal; Fe: femur; Mt: metatarsus; PME: posterior median eyes; PLE: posterior lateral eyes; Pa: patella; pl: prolateral; rl: retrolatral; Ta: tarsus; Ti: tibia;
BMNH: British Museum of natural History, London;
CRB: Collection R. Bosmans;
HECO: Hope entomological Collection, Oxford;
MCSG: Museo civico di Storia naturale "G. Doria", Genova;
MNHN: Muséum national d'Histoire naturelle, Paris;
SMNS: Staatliches Museum für Naturkunde, Stuttgart;
ZMB: Zoologisches Museum, Berlin.

Systematics

Family Araneidae

Neoscona subfuscata (C. L. Koch, 1837) (Fig. 1a-g)

Atea subfuscata C. L. Koch, 1837: 4 (descr. ♂, ♀).

Araneus arganicola Simon, 1909: 26 (descr. ♂, ♀) **syn. nov.**




Fig. 1: *Neoscona subfusca* (C. L. Koch, 1837), type material of *Araneus arganicola* Simon, 1909. **a.** Male, dorsal view; **b.** Female, dorsal view; **c.** Female, ventral view; **d.** Male palp, retrolateral view; **e.** Idem, prolateral view; **f.** Idem, ventro-retrolateral view; **g.** Epigyne, ventral view

Neoscona subfusca; Grasshoff 1980: 406 (transfer from *Araneus*, ♂, ♀).

Neoscona subfusca; Levy 1998a: 336, figs 96–107 (♂, ♀, synonymy). For a complete list of synonyms, see WSC (2021).

Type material. Syntypes of *Araneus arganicola* containing 1 ♂ 2 ♀♀ from Morocco, Mogador (= Essaouira), de La Escalera leg. (MNHN 15552 and 24415); examined.

Comments. *Araneus arganicola* was described from Morocco by Simon (1909). In his description, Simon thought it was closely related to *Neoscona dalmatica*, now a junior synonym of *N. subfusca*. Differences to *N. subfusca* or to the related *N. adianta* were not given. Examination of what we here select as the type material of *Neoscona arganicola* (Fig. 1a–g) reveals it is a synonym of *N. subfusca*. In the male palp, there is one terminal spike in the median apophysis (two in *N. adianta*) and the female epigyne has a relatively wide scape (narrow in *N. adianta*). *Araneus arganicola* also has the same opisthosomal pattern as *N. subfusca*, thus it does not differ morphologically from *Neoscona subfusca* and becomes its junior synonym.

Morano & Bonal (2018) also examined material of *A. arganicola* deposited in the MNHN, but they only saw subadult specimens. In their opinion, *A. arganicola* was a probable synonym of *A. subfusca*, which is confirmed here.

In Algeria, *Neoscona adianta* was only found in coastal regions, whereas *N. subfusca* occurs in Morocco, Algeria and Tunisia, as well as in the southern parts, including oases in the Sahara Desert (see examined material below). The type locality of *N. arganicola* is in the south of Morocco, therefore its habitat preferences are the same as those of *N. subfusca*.

Examined comparative material of *Neoscona subfusca* from North Africa. ALGERIA: M'sila: Baniou S., Chott el Hodna

(35.41°N, 4.34°E), 400 m a.s.l., 1 ♀, on herbs in salt marsh, 13. May 1988, R. Bosmans leg. (CRB). Tamanrasset: Gare de l'Assekrem (23.26°N, 5.63°E), 2700 m a.s.l., 1 ♀, Jul. 1979, R. Bosmans leg. (CRB).

MOROCCO: Fès-Boulemane: Missour (33.00°N, 3.99°E), 880 m a.s.l., 2 ♀♀, pitfall traps in steppe, 13. Sep. 2002 (CRB). Souss-Massa: between Imitek and Tata (29.69°N, 3.83°E), 740 m a.s.l., 1 ♀, among stones in palm groove, 14. Feb. 2007, R. Bosmans leg. (CRB).

TUNISIA: Beja: Oued Zergha (35.40°N, 9.62°E), 2 ♂♂ 4 ♀♀, on olive trees, V. Bouters leg. (CRB). Bizerte: Teskraia E., NW Lake Ichgeul (37.17°N, 9.61°E), 495 m a.s.l., 1 ♂ 1 ♀, beating hedges, 7. May 2006, R. Bosmans leg. (CRB). Tozeur: Dhgoumes (33.98°N, 8.46°E), 15 m a.s.l., 1 ♀, in oasis, 14. Apr. 2004, K. De Smet (CRB).

Examined material of *Neoscona adianta* from North Africa. ALGERIA: Bejaia: river mouth of Oued Daas near Tichi (36.87°N, 4.82°E), 5 m a.s.l., 4 ♀♀, beating bushes in dunes, 22. May 1988, R. Bosmans leg. (CRB); Adekar E. (36.68°N, 4.67°E), 800 m a.s.l., 2 ♀♀, 17. Apr. 1982, on herbs in mixed forest, R. Bosmans leg. (CRB). Blida: Atlas Blidéen, Maison Forestière de Meurdja (36.49°N, 3.14°E), 900 m a.s.l., 3 ♂♂, sweeping herbs in garden, 30. May 1987, R. Bosmans leg. (CRB).

Distribution. Southern Europe, Africa, Russia (Europe to Central Asia), Middle East (WSC 2021).

Family Gnaphosidae

Drassodes obscurus (Lucas, 1846) (Fig. 2a)

Drassus obscurus Lucas, 1846: 214, pl. 13, fig. 1 (descr. ♀).

Amaurobius obscurus; Simon 1864: 139 (transfer).

Drassodes obscurus; Reimoser 1919: 162 (transfer).

Type material. Holotype ♀ of *Drassus obscurus* Lucas, 1846

from Algeria, El Kala (36.88°N , 8.43°E), collection Lucas (MNHN); examined.

Comments. *Drassus obscurus* was described from Algeria and the species has never been studied since. Roewer (1955) considered it a nomen nudum. In the WSC (2021), it is listed under *Drassodes*. The discovery of the holotype shows it is indeed a *Drassodes* species, belonging to the *D. lutescens* group, which is evident from the structure of the vulva shown in the Fig. 2a. The placement of the species in the genus *Drassodes* is confirmed here. A revision of the North African species of this genus has to be undertaken to establish its exact taxonomic status.

Megamyrmaekion caudatum Reuss, 1834 (Fig. 2b-c)

Megamyrmaekion caudatum Reuss, 1834: 212, pl. 18, fig. 12 (descr. juvenile).

Drassus tarrhunensis Karsch, 1881: 12, pl. 1, fig. 11 (descr. ♀) **syn. nov.**

Megamyrmaekion caudatum; Levy 2009: 14, figs 29-32 (♀, synonymy).

Type material. Holotype ♀ of *Drassus tarrhunensis* from Libya, Djebel Tarrhuna, Bir Milra (32.43°N , 1.63°E), Dec. 1878, Rohlfs leg. (ZMB 3293); examined.

Comments. *Drassus tarrhunensis* Karsch, 1881 is only known from the original description and was never recorded again. It is mentioned in the WSC (2021) as *Drassodes tarrhunensis*. The holotype female was examined and is in a poor condition but the epigyne is intact. It shows a large, lenticular pit with an anterior hood and two pairs of receptacula at each side of the depression (Fig. 2b-c). This type of epigyne shows strong similarities to that of *Megamyrmaekion caudatum* and we consider *Drassus tarrhunensis* a junior synonym of that species.

Megamyrmaekion caudatum was described from Egypt and was redescribed by Levy (2009). Levy (op. cit.) also discovered that *M. pumilum* Simon, 1885, which Simon (1885) described from the Algerian, Tunisian and Egyptian Sahara, is a synonym of *Megamyrmaekion caudatum*. Caporiacco (1933) misidentified a specimen of *Megamyrmaekion caudatum* as *Zelotes fuscorufus* (Simon, 1878) from Libya. During our numerous excursions in Morocco, Algeria and Tunisia, *Megamyrmaekion caudatum* was never collected.

Distribution. Desert areas of Algeria, Tunisia, Libya, Egypt, Israel and Iran.

Odontodrassus mundulus (O. Pickard-Cambridge, 1872)

Drassus mundulus O. Pickard-Cambridge, 1872: 234, pl. 15, fig. 11 (descr. ♂, ♀).

Drassus nugatorius Karsch, 1881: 12, pl. 1, fig. 10 (♀) **syn. nov.** *Poecilochroa monodi* Fage, 1929: figs 2-3 (descr. ♂, ♀); synonymy by Levy (1999).

Odontodrassus mundulus; Levy 1999: 448, figs 43-46 (transfer). **Type material.** Syntypes 1 ♂, 2 ♀♀ of *Drassus mundulus* from Israel, plains of the Jordan (HECO, B. 243, t. 46, 1 ♀, t. 69, 1 ♀) and Egypt, Cairo (30.05°N , 31.23°E) (HECO, B. 241, t. 85, 1 ♂, 1 ♀); examined by Levy (1999).

Syntypes ♂♂, ♀♀ of *Poecilochroa monodi* from Algeria and Tunisia (MNHN, B. 623, t. 5218, t. 13198, t. 19524) and from Egypt (MNHN B. 623, t. 6234); examined by Levy (1999).

Holotype ♀ of *Drassus nugatorius* from Libya, palm garden near Sockna (29.07°N , 15.78°E), 4. Feb. 1879, Rohlfs leg. (ZMB 3294); examined.

Description. See Levy (1999).

Comments. *Drassus nugatorius* Karsch, 1881 is only known from the original description and has never been recorded again. In the WSC (2021), it is listed under *Drassodes*. The holotype female was examined and is in good condition. The epigyne is very typical and is identical to that of *Odontodrassus mundulus*. Moreover, the drawings of the epigynes by Pickard-Cambridge (1872, fig. 11), Karsch (1881, fig. 10) and Fage (1929, fig. 2) all show the curved copulatory ducts forming a letter X. *Drassus nugatorius* Karsch, 1884 thus becomes a junior synonym of *O. mundulus*. In the Maghreb, *O. mundulus* was reported from Libya by Caporiacco (1928, 1936a, 1936b) and from Algeria by Denis (1966).

New records. ALGERIA: El Bayadh: Brezina E. (33.09°N , 1.25°E), 800 m a.s.l., 1 ♂, stones in steppe, 8. Feb. 1987, R. Bosmans leg. (CRB). Illizi: Iherir (35.40°N , 8.40°E), 1200 m a.s.l., 1 ♀, 12. Feb. 1996, K. De Smet leg. (CRB); 70 km S to Illizi, Oued Djeret (26.29°N , 8.63°E), 700 m a.s.l., 1 ♀, 10. Feb. 1985 (CRB). M'sila: Aïn Oghrab (33.09°N , 1.25°E), 650 m a.s.l., 1 ♂, pitfall traps in open *Pinus* forest, 20. Mar.-23. Jun. 1990, R. Bosmans leg. (CRB). Tamanrasset: Massif de l'Ahaggar, Oued Tit (22.90°N , 4.91°E), 1 ♀, 17. Feb. 1988, K. De Smet leg. (CRB).

MOROCCO: Marrakesh-Safi: Aït Barka (31.48°N , 7.44°E), 1300 m a.s.l., stones in *Pinus* forest, 6. Jun. 1999, R. Bosmans leg. (CRB). Souss-Massa: Barrage Youcef Ben-Tachfine (29.84°N , 9.49°E), 100 m a.s.l., 1 ♀, 27. Apr. 2012, R. Bosmans leg. (CRB).

TUNISIA: Kairouan: road Aïn Djeloula-Oglet Tarfa ($35.763611^{\circ}\text{N}$, 9.955833°E), 60 m a.s.l., 1 ♂, 23. Jan. 1995, R. Bosmans leg. (CRB).

JORDAN: At Tafila: Dana (30.67°N , 35.62°E), 1550 m a.s.l., 1 ♂, stones in irrigated garden, 17. Nov. 2007, R. Bosmans leg. (CRB). Al Karak: Shawbak castle (30.53°N , 33.65°E), 1250 m a.s.l., 1 ♂ 1 ♀, stones on slope to castle, 17. Nov. 2007, R. Bosmans leg. (CRB).

Distribution. *Odontodrassus mundulus* is known from Morocco, Algeria, Tunisia, Libya, Egypt and Israel. It is reported here for the first time in Jordan.

Poecilochroa pugnax (O. Pickard-Cambridge, 1874) (Fig. 2d-e)

Drassus pugnax O. Pickard-Cambridge, 1874: 399, pl. 52, fig. 25 (descr. ♂).

Drassus sockniensis Karsch, 1881: 12, pl. 1, fig. 9 (descr. ♀) **syn. nov.**

Poecilochroa pugnax; Simon 1908: 423 (descr. ♀).

Poecilochroa lesserti Denis, 1947: 62 pl. 3, figs 8-10 (descr. ♂, ♀); synonymy by Levy (1999).

Poecilochroa pugnax; Levy 1999: 433, figs 10-14 (♂, ♀).

Type material. Holotype ♂ of *Drassus pugnax* from Egypt, Cairo (30.05°N , 31.23°E) (HECO, b.254); examined and re-described by Levy (1999).

Syntypes 1 ♂ and 1 ♀ of *Poecilochroa lesserti* from Egypt, Siwa, (BMNH, ♂ 1936.7.10.136a, ♀ 1936.2.12.1049); examined by Levy (1999).

Holotype ♀ of *Drassus sockniensis* from Libya, palm garden near Sockna, 4. Feb. 1879, Rohlfs leg. (ZMB 3293); examined.

Comments. The holotype female of *Drassus sockniensis* is in very bad condition, but the epigyne and vulva is intact (Fig. 2d-e) and shows very large receptacula typical for the genus

Poecilochroa. In Libya, two species of *Poecilochroa* have been observed: *Poecilochroa pugnax* (O. Pickard-Cambridge, 1874) and *P. senilis* (O. Pickard-Cambridge, 1872). Both species are similar in having elongated, sausage-shaped receptacula. They differ by the presence of ducts reaching half the length of the receptacula in *P. senilis* and less than one third the length of the receptacula in *P. pugnax* (Levy 1999). In the type specimen of *Drassus sockniensis*, ducts are limited to the basal third of the receptacula (Fig. 2e), just like in *P. pugnax*, and we consider *P. sockniensis* (Karsch, 1881) and *P. pugnax* synonymous.

Setaphis mollis (O. Pickard-Cambridge, 1874)

Prosthesima mollis O. Pickard-Cambridge, 1874: 381, pl. 51, fig. 9 (descr. ♀).

Echemus pharetratus Karsch, 1881: 11, pl. 1, fig. 8 (descr. ♀; synonymy by Bosmans & Janssen 1999).

Prosthesima quadridentata Strand, 1906: 613 (descr. ♀); Strand, 1908: 75 (descr. ♀) **syn. nov.** and removed from nomen dubium, contra Nentwig et al. (2020: 24).

Scotophaeus quadridentatus Caporiacco, 1928: 85, fig. 2 (descr. ♀) **syn. nov.**

Pseudodrassus quadridentatus; Caporiacco 1935: 286 (transfer).

Setaphis mollis; Platnick & Murphy 1996: 12, figs 25–28 (descr. ♂); Levy 1998b: 97, figs 6–9 (♂, ♀); Bosmans & Janssen 1999: 86, figs 9–12 (♂, ♀; synonymy).

Type material. Holotype ♀ of *Prosthesima quadridentata* Strand, 1906 from Tunis (36.82°N, 10.15°E) (SMNS); not examined, type material destroyed.

Lectotype ♀ of *Scotophaeus quadridentatus* Caporiacco, 1928 by present designation, from Libya, Giarabub (29.72°N, 24.52°E) (MCSG); examined.

Description. See Platnick & Murphy (1996), Levy (1998b) and Bosmans & Janssen (1999).

Comments. Caporiacco (1928) described the female of *Scotophaeus quadridentatus* as a new species from Libya. Two localities are mentioned in the original description: Porto Bardia and the Giarabub Oasis. The female from the Giarabub Oasis was available for study and selected as the lectotype, and the specimen appears to be identical to *Setaphis mollis* (O. Pickard-Cambridge, 1874), and hence becomes a junior synonym. The name was probably given because of the sclerotic grooves in the epigyne forming a central quadrangle, also seen in Caporiacco's (1928) fig. 2.

It must be pure coincidence that Strand (1906: 613) gave the same species name, *Prosthesima quadridentata*, to a species he described from Tunisia. In Strand (1908: 76) the epigyne was described as follows: "Epigyne bildet in Fluidum gesehen ein abgerundet viereckiges, bräunliches Feld, das ein wenig länger als breit ist". Not only the shape of the epigyne but also the size of the spider and the yellowish brown general colour are similar. Since the type specimens no longer exist, it is our opinion this species should be considered a junior synonym of *Setaphis mollis* as well. Nentwig et al. (2020) suggested the species was a nomen dubium, but we are of the opinion that the arguments above are sufficient to consider it a junior synonym of *Setaphis mollis*.

Distribution. Algeria, Tunisia, Libya and Egypt.

Zelotes tragicus (O. Pickard-Cambridge, 1872) (Fig. 2f)

Melanophora tragica O. Pickard-Cambridge, 1872: 243, pl. 16, fig. 22 (descr. ♂).




Fig. 2: a. *Drassodes obscurus* (Lucas, 1846), holotype female, vulva, dorsal view; b-c. *Megamyrmaekion caudatum* Reuss, 1834, holotype female of *Drassus tarrhunensis* Karsch, 1881; b. Epigyne; c. Vulva, dorsal view; d-e. *Poecilochroa pugnax* (O. Pickard-Cambridge, 1874), holotype female of *Drassus sockniensis* Karsch, 1881; d. Epigyne; e. Vulva, dorsal view; f. *Zelotes tragicus* O. Pickard-Cambridge, 1872, lectotype female of *Prosthesima kerimi* Pavesi, 1880, epigyne

Prosthesima tragicus; Simon 1878: 98 (transfer).

Prosthesima kerimi Pavesi, 1880: 348 (descr. ♂, ♀) **syn. nov.**

Zelotes tragicus; Levy 1998b: 133, figs 82–85 (descr. ♀).

Zelotes tragicus; FitzPatrick 2007: 121, figs 81–84 (♂, ♀).

Type material. Lectotype ♀, paralectotype ♂ (without palps), by present designation, of *Prosthesima kerimi* from Tunisia, Ludien near Tozeur (33.92°N, 10.28°E), Kerim leg. (MCSG); examined.

Description. See Levy (1998b) and FitzPatrick (2007).

Comments. *Prosthesima kerimi* was described more than 100 years ago by Pavesi (1880), but has never been illustrated nor clearly diagnosed. The type series could be examined and it is composed of one male, without palps, and one female, with the epigyne present (Fig. 2f). The female is therefore selected as the lectotype. It appears to be identical to *Zelotes tragicus* (O. Pickard-Cambridge, 1872), a common species in North Africa.

Distribution. Tunisia, Libya, Chad, Ethiopia, Israel.

Family Linyphiidae

Pelecopsis pavesii spec. nov. (Figs 3a-g, 4a-e, Tab. 1)

Erigone parumpunctata; Pavesi 1884: 459 (misidentification).

Type material. ♂ holotype: TUNISIA, surroundings of Tunis ("dintorni di Tunis"), winter 1881–1882, G. & L. Doria leg.; coll. MCSG; misidentified as *Pelecopsis parumpunctata* (Simon, 1881) by Pavesi (1884).



Fig. 3: *Pelecopsis pavesii* spec. nov. **a.** Male, dorsal view; **b.** Idem, lateral view; **c.** Idem, frontal view; **d.** Male palpal tibia, dorsal view; **e.** Male palp, retrolateral view; **f.** Idem, pro-lateral view; **g.** Embolic division, ventral view

Comments. *Pelecopsis parumpunctata* was described by Simon (1881a) from France and recorded in Tunisia by Pavesi (1884). Denis (1964: 345) synonymized it with *Pelecopsis mengei* (Simon, 1884). Although *Pelecopsis parumpunctata* was described before *P. mengei* and should have priority, the name *P. parumpunctata* was apparently suppressed for lack of usage (see WSC 2021). In their revisions of North African *Pelecopsis* species, neither Denis (1962), nor Bosmans & Abrous (1992) mention *P. mengei* from North Africa. Examination of the specimen cited by Pavesi (1884) reveals that it is not *P. mengei* but an unknown species described below.

Diagnosis. By the presence of a basal cymbial tubercle, the species is similar to *Pelecopsis kalaensis* Bosmans, 1992, *P. laptevi* Tanasevitch & Fet, 1986, *P. odontophora* (Kulczyński, 1895), *P. paralleloides* Tanasevitch & Fet, 1986, *P. pavida* (O. Pickard-Cambridge, 1872) and *P. susannae* (Simon, 1915). Of these, only *Pelecopsis kalaensis* occurs in North Africa and this species differs from the new species by the palp with a rectangular dorsal tibial apophysis, which is triangular in *P. pavesii* spec. nov. (Figs 3d, 4c) and also by the elongated palpal femur (Fig. 3b) in *P. pavesii* spec. nov.

Etymology. The species is dedicated to Pietro Pavesi, author of two important contributions to the arachnofauna of Tunisia.

Description. Measurements: Total length 1.5; prosoma 0.85 wide, 0.63 wide. Legs (Tab. 1).

Colour: The specimen is too faded to give information on this feature.

Carapace (Figs 3a-c, 4a-b): With large cephalic elevation carrying PME, a transverse frontal groove, oval postocular sulci, twice the diameter of the PLE and two lines of impressed dots between the PME; AME separated by 2.5 times their diameter, from ALE by 2.5 times their diameter; PME separated by 4 times their diameter.

Tab. 1: Leg measurements of the holotype

	Fe	Pa	Ti	Mt	Ta	Total
I	0.61	0.24	0.49	0.40	0.34	2.08
II	0.60	0.23	0.45	0.39	0.31	1.98
III	0.49	0.21	0.39	0.36	0.30	1.75
IV	0.64	0.23	0.65	0.48	0.31	2.31
Palp	0.42	—	0.37	0.20	0.35	1.34

Opisthosoma: Covered with opaque scutum.

Legs: No spines or trichobothria observed in this old, faded specimen, probably detached and no sign of attachment places.

Palp (Figs 3d-g, 4c-e): Femur elongated, three times longer than wide; tibia with three triangular apophyses, two small retro- and prolateral and a pointed median apophysis. Cymbium with basal tubercle; protegulum protruding; embolus ribbon-like, semi-circular, terminally bent in anterior direction as seen in retrolateral view.

Thaumatoncus indicator Simon, 1884

Thaumatoncus indicator Simon, 1884: 581 (descr. ♂)

Erigone digiticeps Simon, 1881a; Pavesi 1884: 46 (misidentification).

Thaumatoncus indicator; Bosmans 2002: 19 (♂).

Material examined. TUNISIA: Surroundings of Tunis ("dintorni di Tunis"), 2 ♂♂, winter 1881–1882, G. & L. Doria leg. (MCSG); examined.

Comments. Pavesi (1884) cited *Erigone digiticeps* Simon, 1881, now *Dactylopisthes digiticeps* (Simon, 1881), from Tunisia. This species has never been found in North Africa and Bosmans (1996) considered it a misidentification of *Delorrhripis fronticornis* Simon, 1884. Examination of Pavesi's specimen now reveals it is *Thaumatoncus indicator*, a fairly common




Fig. 4: *Pelecopsis pavesii* spec. nov. **a.** Male, dorsal view; **b.** Idem, lateral view; **c.** Male palpal tibia, dorsal view; **d.** Male palp, retralateral view; **e.** Embolic division, ventral view

species in Tunisia (Bosmans 2002). *Delorrhapis fronticornis* should be deleted from the spider species list of Tunisia.

Distribution. Algeria, Tunisia, Spain, S. France (Bosmans 2002, WSC 2021).

Family Theridiidae

***Steatoda erigoniformis* (O. Pickard-Cambridge, 1872)** (Fig. 5a-f)

Theridion erigoniforme O. Pickard-Cambridge, 1872: 284 (descr. ♂, ♀).

Steatoda signata O. Pickard-Cambridge, 1876: 568 (descr. ♀, synonymy by Levy & Amitai, 1982).

Lithyphantes septemmacculatus Keyserling, 1884: 141, pl. 6, fig. 88 (descr. ♀).

Steatoda erigoniformis; Levy & Amitai 1982: 26, figs 63-71 (redescr. ♂, ♀; synonymy).

Euryopis albomaculata Denis, 1951: 313, fig. 1 (descr. ♀) **syn. nov.**

Type material. Holotype ♀ of *Euryopis albomaculata* from Egypt, Sawaleh, 5 km S. Fakous (30.07°N 1.83°E), 7. Sep. 1949, M. B. Condé leg. (MNHN); not examined.

Comments. *Steatoda erigoniformis* has a complex taxonomic history. It was first described in the genus *Theridion* by Pickard-Cambridge (1872). Later, it was redescribed several times as a new species in two different genera: as *Steatoda signata* by Pickard-Cambridge (1876), and as *Lithyphantes septemmacculatus* by Keyserling (1884). It has also been placed in several other genera: in *Asagenella* by Schenkel (1937) and in *Crustullina* by Simon (1881b). *Euryopis albomaculata* Denis, 1951, described from Egypt, appears to be another junior synonym. Comparing Denis' fig. 1 of the epigyne with fig. 70 of Levy & Amitai (1982) (see Fig. 5c-f) and regarding also the small size and the opisthosomal white spots of both species it is evident they are synonyms.



Fig. 5: *Steatoda erigoniformis* O. Pickard-Cambridge, 1872. **a.** Male, dorsal view; **b.** Female, dorsal view; **c.** Epigyne; **d.** Vulva; **e.** Epigyne, drawing by Levy & Amitai (1982) of *Steatoda erigoniformis*; **f.** Epigyne, drawing of *Euryopis albomaculata* by Denis (1951)

Distribution. East Mediterranean to Middle East, Caucasus, China, Korea, Japan. Introduced to the Caribbean (WSC 2021).

Acknowledgements

The authors sincerely thank Maria Tavano, curator of the Museo civico di Storia naturale "G. Doria", for the loan of the Pavesi material, Christine Rollard, curator of the Muséum d'Histoire naturelle de Paris for the loan of the Lucas and Simon material and Jason Dunlop, curator of Zoologisches Museum, Berlin for the loan of the Karsch material. Pierre Oger is thanked for the photos. Finally, Jørgen Lissner, Andrei V. Tanasevitch, an anonymous reviewer, and Theo Blick, Tobias Bauer, Petr Dolejš, Konrad Wiśniewski are thanked for their most valuable comments.

References

- Bosmans R 1996 The genera *Araeoncus* Simon, *Delorrhapis* Simon and *Diplocephalus* Bertkau in northern Africa (Araneae: Linyphiidae: Erigoninae): Studies on North African Linyphiidae VII. – Belgian Journal of Zoology 126: 123-151
- Bosmans R 2002 Les genres *Acartauchenius* Simon et *Thaumatoncus* Simon en Afrique du Nord. Etudes sur les Linyphiidae africaines, no IX (Araneae, Linyphiidae, Erigoninae). – Revue Arachnologique 14: 1-24
- Bosmans R & Abrous O 1992 Studies on North African Linyphiidae VI The genera *Pelecopsis* Simon, *Trichopterna* Kulczynski and *Ouedia* gen. n. (Araneae: Linyphiidae). – Bulletin of the British Arachnological Society 9: 65-85
- Bosmans R & Janssen M 1999 The ground spider genus *Setaphis* in the Maghreb countries (Araneae Gnaphosidae). – Bulletin & Annales de la Société Entomologique de Belgique 155: 82-90

- Caporiacco L di 1928 Resultati zoologici della Missione inviata dalla R. Società Geografica Italiana per l'esplorazione dell'oasi di Giarabub (1926-1927). Aracnidi di Giarabub e di Porto Bardia (Tripolis). – Annali del Museo Civico di Storia Naturale di Genova 53: 77-107
- Caporiacco L di 1933 Araneidi. In: Spedizione scientifica all'oasi di Cufra (Marzo-Luglio 1931). – Annali del Museo Civico di Storia Naturale di Genova 56: 311-340
- Caporiacco L di 1935 Escursione del Prof. Nello Beccari in Anatolia. Aracnidi. – Monitore Zoologico Italiano 46: 283-289
- Caporiacco L di 1936a Aracnidi raccolti durante la primavera 1933 nelle oasi del deserto libico. – Memorie della Società Entomologica Italiana, Genova 15: 93-122
- Caporiacco L di 1936b Aracnidi fezzanesi raccolti dal Prof. G. Scortecci nel 1934 - XII (Missione della R. Società Geografica). – Atti della Società Italiana di Scienze Naturali e del Museo Civico di Storia Naturale di Milano 75: 67-94
- Denis J 1947 Results of the Armstrong College expedition to Siwa Oasis (Libyan desert), 1935. Spiders [Araneae]. – Bulletin de la Société Fouad 1er d'Entomologie 31: 17-103
- Denis J 1951 Quelques araignées de Basse-Égypte recueillies par M. B. Condé. – Bulletin de la Société Zoologique de France 76: 313-315
- Denis J 1962 Notes sur les érigonides XXIV Les *Pelecopsis* nord-africaines. – Bulletin de la Société des Sciences Naturelles du Maroc 42: 279-291
- Denis J 1964 Notes sur les érigonides. XXVII. Remarques sur les genres *Pelecopsis* Simon et *Trichopterna* Kulcz., avec la description du mâle de *Pelecopsis major* (Denis). – Bulletin du Muséum National d'Histoire Naturelle de Paris (2) 36: 339-347
- Denis J 1966 Les araignées du Fezzâne. – Bulletin de la Société d'Histoire Naturelle d'Afrique du Nord 55: 103-144
- Fage L 1929 Mission saharienne Augieras-Draper 1927-1928. Araignées nouvelles. – Bulletin du Muséum National d'Histoire Naturelle de Paris (2) 1: 248-254
- FitzPatrick MJ 2007 A taxonomic revision of the Afrotropical species of *Zelotes* (Arachnida: Araneae: Gnaphosidae). – Bulletin of the British Arachnological Society 14: 97-172 – doi: [10.13156/arac.2011.14.3.97](https://doi.org/10.13156/arac.2011.14.3.97)
- Grasshoff M 1980 Contributions à l'étude de la faune terrestre des îles granitiques de l'archipel des Sénèches (Mission P.L.G. Benoit - JJ. Van Mol 1972). Aranidae-Arhopinae, Aranidae-Araneinae (Araneae). – Revue Zoologique Africaine 94: 387-409
- Karsch F 1881 Verzeichniss der während der Rohlfs'schen Afrikanischen Expedition erbeuteten Myriopoden und Arachniden. – Archiv für Naturgeschichte 47: 1-14
- Keyserling E 1884 Die Spinnen Amerikas II. Theridiidae. I. Hälfte. Bauer & Raspe, Nürnberg. pp. 1-222, pl. 1-10
- Koch CL 1837 Übersicht des Arachnidensystems. C. H. Zeh'sche Buchhandlung, Nürnberg, Heft 1, pp. 1-39. – doi: [10.5962/bhl.title.39561](https://doi.org/10.5962/bhl.title.39561)
- Levy G 1998a Twelve genera of orb-weaver spiders (Araneae, Aranidae) from Israel. – Israel Journal of Zoology 43: 311-365
- Levy G 1998b The ground-spider genera *Setaphis*, *Trachyzelotes*, *Zelotes*, and *Drassyllus* (Araneae: Gnaphosidae) in Israel. – Israel Journal of Zoology 44: 93-158
- Levy G 1999 Spiders of six uncommon drassodine genera (Araneae: Gnaphosidae) from Israel. – Israel Journal of Zoology 45: 427-452
- Levy G 2009 New ground-spider genera and species with annexed checklist of the Gnaphosidae (Araneae) of Israel. – Zootaxa 2066: 1-49 – doi: [10.11646/zootaxa.2066.1.1](https://doi.org/10.11646/zootaxa.2066.1.1)
- Levy G & Amitai P 1982. The cobweb spider genus *Steatoda* (Araneae, Theridiidae) of Israel and Sinai. – Zoologica Scripta 11: 13-30. – doi: [10.1111/j.1463-6409.1982.tb00515.x](https://doi.org/10.1111/j.1463-6409.1982.tb00515.x)
- Lucas H 1846 Histoire naturelle des animaux articulés In: Exploration scientifique de l'Algérie pendant les années 1840, 1841, 1842 publiée par ordre du Gouvernement et avec le concours d'une commission académique Paris, Sciences physiques, Zoologie 1: 89-271 – doi: [10.5962/bhl.title.112444](https://doi.org/10.5962/bhl.title.112444)
- Morano E & Bonal R 2018 *Araneus bonali* sp. n., a novel lichen-patterned species found on oak trunks (Araneae, Araneidae). – ZooKeys 779: 119-145 – doi: [10.3897/zookeys.779.26944](https://doi.org/10.3897/zookeys.779.26944)
- Nentwig W, Blick T, Gloor D, Jäger P & Kropf C 2020 How to deal with destroyed type material? The case of Embrik Strand (Arachnida: Araneae). – Arachnologische Mitteilungen 59: 22-29 – doi: [10.30963/aramit5904](https://doi.org/10.30963/aramit5904)
- Pavesi P 1880 Studi sugli Aracnidi africani. I. Aracnidi di Tunisia. – Annali del Museo Civico di Storia Naturale di Genova 15: 283-388
- Pavesi P 1884 Materiali per lo studio della fauna tunisina raccolti da G. e L. Doria: Aracnidi. – Annali del Museo Civico di Storia Naturale di Genova 20: 446-486
- Pickard-Cambridge O 1872 General list of the spiders of Palestine and Syria, with descriptions of numerous new species, and characters of two new genera. – Proceedings of the Zoological Society of London 40: 212-354, pl. 13-16 – doi: [10.1111/j.1469-7998.1872.tb00489.x](https://doi.org/10.1111/j.1469-7998.1872.tb00489.x)
- Pickard-Cambridge O 1874 On some new species of Drassidae. – Proceedings of the Zoological Society of London 42(3): 370-419 – doi: [10.1111/j.1096-3642.1874.tb02495.x](https://doi.org/10.1111/j.1096-3642.1874.tb02495.x)
- Pickard-Cambridge O 1876 Catalogue of a collection of spiders made in Egypt, with descriptions of new species and characters of a new genus. – Proceedings of the Zoological Society of London 44(3): 541-630, pl. 58-60. – doi: [10.1111/j.1096-3642.1876.tb02595.x](https://doi.org/10.1111/j.1096-3642.1876.tb02595.x)
- Platnick NI & Murphy JA 1996 A review of the zelotine ground spider genus *Setaphis* (Araneae, Gnaphosidae). – American Museum Novitates 3162: 1-23
- Reimoser E 1919 Katalog der echten Spinnen (Araneae) des Paläarktischen Gebietes. – Abhandlungen der Zoologisch-Botanischen Gesellschaft in Wien 10(2): 1-280
- Reuss A 1834 Zoologische Miscellen. – Museum Senckenbergianum, Abhandlungen aus dem Gebiete der beschreibenden Naturgeschichte 1: 195-276
- Roewer CF 1955 Katalog der Araneae von 1758 bis 1940, bzw. 1954. 2. Band, Abt. a (Lycosaeformia, Dionycha [excl Salticiformia]). 2. Band, Abt b (Salticiformia, Cribellata) (Synonyma-Verzeichnis, Gesamtindex). Institut royal des Sciences naturelles de Belgique, Bruxelles. 1751 pp.
- Schenkel E 1937 Beschreibungen einiger afrikanischer Spinnen und Fundortsangaben. – Festschrift Embrik Strand 3: 373-398
- Simon E 1864 Histoire naturelle des araignées (aranéides). Paris. 540 pp. – doi: [10.5962/bhl.title.47654](https://doi.org/10.5962/bhl.title.47654)
- Simon E 1878 Les arachnides de France. Paris 4, 1-334
- Simon E 1881a Description d'espèces nouvelles du genre *Erigone*. – Bulletin de la Société Zoologique de France 6: 233-257
- Simon E 1881b Les arachnides de France. Tome cinquième, première partie. Roret, Paris. 180 pp.
- Simon E 1884 Les arachnides de France. Tome cinquième, deuxième et troisième partie. Roret, Paris. pp. 180-885
- Simon E 1885 Etudes sur les Arachnides recueillis en Tunisie en 1883 et 1884 par MM. A. Letourneau, M. Sébillot et Valéry Mayet, membres de la mission de l'Exploration scientifique de la Tunisie. In: Exploration scientifique de la Tunisie, publiée sous les auspices du Ministère de l'instruction publique. Zoologie – Arachnides. Paris. 55 pp.
- Simon E 1908 Etude sur les arachnides recueillis par M. le Dr. Klaptocz en Tripolitaine. – Zoologische Jahrbücher, Abteilung für Systematik, Geographie und Biologie der Tiere 26: 419-438
- Simon E 1909 Etude sur les arachnides recueillis au Maroc par M. Martinez de la Escalera en 1907. – Memorias de la Real Sociedad Española de Historia Natural (6) 1: 5-43
- Strand E 1906 Diagnosen nordafrikanischer, hauptsächlich von Carlo Freiherr von Erlanger gesammelter Spinnen. – Zoologischer Anzeiger 30: 604-637, 655-690
- Strand E 1908 Nordafrikanische, hauptsächlich von Carlo Freiherr von Erlanger gesammelte Aviculariidae, Drassidae und Theridiidae. – Jahreshefte des Vereins für vaterländische Naturkunde in Württemberg 64: 11-101
- WSC 2021 World Spider Catalog. Version 22.0. Natural History Museum Bern. – Internet: <http://wsc.nmbe.ch> (15. Jan. 2020) – doi: [10.24436/2](https://doi.org/10.24436/2)