

Natural Nest Sites of the Red-Breasted Flycatcher *Ficedula parva* in a Primeval Forest

Authors: Mitrus, Cezary, and Soćko, Beata

Source: *Acta Ornithologica*, 39(1) : 53-57

Published By: Museum and Institute of Zoology, Polish Academy of Sciences

URL: <https://doi.org/10.3161/068.039.0111>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Natural nest sites of the Red-breasted Flycatcher *Ficedula parva* in a primeval forest

Cezary MITRUS & Beata SOĆKO

Department of Zoology, University of Podlasie, Prusa 12, 08–110 Siedlce, POLAND, e-mail: ficedula@ap.siedlce.pl

Mitrus C., Soćko B. 2004. Natural nest sites of the Red-breasted Flycatcher *Ficedula parva* in a primeval forest. *Acta Ornithol.* 39: 53–57.

Abstract. 117 nests of Red-breasted Flycatchers in Białowieża primeval forest (NE Poland) were characterised. Most nests (79%) were built in Hornbeam *Carpinus betulus* and lime *Tilia cordata*. Three types of nest sites were distinguished: chimney shaped (26.4%), half cavities (46.4%) or shelves — the nest was wedged under a piece of bark against the main trunk (27.3%). Most cavity entrances were exposed to the south. Compared to other species of secondary cavity nesters in Białowieża National Park, Red-breasted Flycatchers used cavities of a different shape, with a smaller bottom area and at a relatively low height above the ground ($\bar{x} = 4.9 \pm 3.13$ m). Nesting trees had a smaller diameter ($\bar{x} = 31.2 \pm 21.4$ cm) and were more often dead (29.8%) than trees used by the other secondary cavity nesters. Only four nest sites were used in consecutive seasons.

Key words: Red-breasted Flycatcher, *Ficedula parva*, nest sites, cavity nesters, holes, Białowieża National Park

Received — Jan. 2004, accepted — March 2004

INTRODUCTION

Nest site features can affect reproductive success by limiting clutch size and cause partial or complete losses of clutches and broods (e.g. van Balen 1973, Slagsvold 1987, Walankiewicz 1991, Rendell & Robertson 1993). Long-term intensive forest management often damages natural habitats and restricts suitable places for breeding (e.g. Bruns 1960, Haapanen 1965). Shortage of suitable nest sites may increase inter- and intra-specific competition and consequently restrict population size (Newton 1998). Knowledge of nest site characteristics can help to formulate management guidelines in order to protect a population.

The Red-breasted Flycatcher is a small passerine that breeds in “half cavities” in forests from Kamchatka to Central Europe (Cramp 1993). The species occurs mainly in mature or young, dense deciduous and mixed forest (Flade 1997). In the oak-hornbeam stands of Białowieża National Park, the Red-breasted Flycatcher reaches breeding densities of up to 2.0 pairs/10 ha (Wesołowski et al. 2002).

Characteristic of natural nest sites of the most secondary cavity nesters breeding in primeval forest

has been presented in few papers so far (Wesołowski 1989, 1996, Mitrus et al. 1996, Walankiewicz 1991, Czeszczewik & Walankiewicz 2003) and only incidental data exist about nest sites of the Red-breasted Flycatcher (Glutz von Blotzheim & Bauer 1993).

The aim of this study was to fill this gap and to describe natural nests sites of the Red-breasted Flycatcher in primeval stands of the Białowieża National Park.

STUDY AREA AND METHODS

Data were collected in 1977–2003 in the Białowieża National Park (BNP) (52°41'N, 23°52'E, NE-Poland). It is the best preserved and strictly protected area of the Białowieża Forest. Almost all nests were located in oak-hornbeam *Tilio-Carpinetum* stands characterised mainly by Hornbeam *Carpinus betulus*, Small-leaved Lime *Tilia cordata*, Pendunculate Oak *Quercus robur*, Norway Maple *Acer platanoides* and Norway Spruce *Picea abies* (Tomiałojć et al. 1984, Tomiałojć 1991).

Most nests (88 out of 117) were found during the last four breeding seasons (2000–2003).

Nests were located following females during nest construction, searching sites where birds were frequently seen, and following adults when they were feeding nestlings. For almost all nests the following parameters were described: nesting tree species, height of the nest above the ground, diameter of the trunk at breast height (DBH), stage of tree (dead or living), type of nest site and for some of the nests depth and bottom area of the cavity. Bottom area was calculated used formula:

$$V = \pi * \text{radius}_1 * \text{radius}_2$$

only for half cavities and chimneys. To analysis were included multiple nests which were used more than once.

The data presented in this paper may possess some bias caused by searching methods. The Red-breasted Flycatcher is a very shy species, especially during incubation period and some nests are very difficult to find. Due to area size and spatial distribution of the species we were able to find only a part of nests of the entire population breeding in Białowieża Forest.

RESULTS

During the study period 117 nests sites occupied by Red-breasted Flycatcher were found. Only four of them were used more than one time (out of 88 sites checked). Most of them were in living trees (70.2%) and mostly situated in Hornbeam and Lime (Table 1). An average height above ground was 4.9 m, but more than 70% were located lower than 6.0 m (Table 1, Fig. 1). The highest nests were situated in hornbeams, but differences in nest height between tree species were not significant ($F_{779} = 0.91$, $p = 0.90$, Table 1).




Fig. 1. Distribution of the height above ground of the Red-breasted Flycatcher's nests.




Fig. 2. Entrance orientation of the nests.

Table 1. Features of nesting trees used by Red-breasted Flycatcher in Białowieża 1977–2003.

| Tree species | Height above ground (m) | | | Diameter at breast height (cm) | | |
|-------------------------|-------------------------|----------|----------|--------------------------------|------------|------------|
| | N | Range | Mean±SD | N | Range | Mean±SD |
| <i>Carpinus betulus</i> | 64 | 1.3-16.0 | 5.4±3.34 | 41 | 13.0-75.0 | 33.1±12.96 |
| <i>Tilia cordata</i> | 27 | 1.5-13.0 | 4.7±2.89 | 27 | 10.0-103.0 | 20.1±22.25 |
| <i>Pinu silvestris</i> | 11 | 1.6-13.0 | 4.9±3.13 | 8 | 47.0-106.0 | 67.9±21.27 |
| <i>Picea abies</i> | 6 | 0.7-6.0 | 2.9±1.94 | 4 | 11.0-46.0 | 25.5±15.37 |
| <i>Ulmus sp.</i> | 3 | 2.5-4.5 | 3.2±1.15 | 3 | 12.0-22.0 | 15.3±5.78 |
| <i>Quercus robur</i> | 2 | 2.0-6.0 | 4.0±2.83 | 2 | 10.0-53.0 | 31.5±30.41 |
| Others | 4 | 1.9-2.5 | 2.2±0.32 | 2 | 30.0-30.0 | 30.0±0.0 |
| Total | 117 | 0.7-16.0 | 4.9±3.13 | 87 | 10.0-106.0 | 31.2±21.44 |




Fig. 3. Types of nest sites: A – half holes, B – chimneys, C – shelves.

The diameter of the nesting tree varied from 10 to 106 cm ($\bar{x} = 31.2 \pm 21.4$, $n = 87$) and depended on the species ($F_{7,79} = 6.97$, $p < 0.001$), pines being the thickest and elms the thinnest (Table 1). Nests were built in shallow cavities ($\bar{x} = 6.1 \pm 2.05$, $n = 12$) with a small bottom area ($\bar{x} = 43.9 \pm 16.99$, $n = 26$). The bottom area was not related to tree diameter ($r = 0.07$, $p = 0.74$, $n = 26$) and height above ground ($r = -0.07$, $p = 0.74$, $n = 26$). Most of holes entrances were directed south, south-east and east (Fig. 2). Distribution differed significantly than normal curve (K-S $d = 0.14$, $p < 0.05$).

Three types of nest sites were distinguished: half cavity (shallow cavity with big entrance), chimney (usually in broken and rotten top of thin tree), shelf (place outside the main trunk, usually under bark, Fig. 3). Most nests were located in half cavities (46.4%), 26.4% were built in chimneys and 27.3% on shelves. The type of nest site was strictly related to tree species. Most chimneys were found in limes (72.5%), whereas half holes and shelves (76.5% and 40.0%, respectively) were located mainly in hornbeams, which is connected with different features of those tree species. Types of nests sites were not related to height above the ground ($F_{2,84} = 2.3$, $p = 0.10$) but were so to tree diameter ($F_{2,84} = 34.15$, $p < 0.001$, Fig. 4).




Fig. 4. Types of nests sites and tree diameter.

DISCUSSION

Only few authors have described nests sites of the Red-breasted Flycatcher so far. The most numerous data presented by Aleknonis (1976) from Lithuania, much less was known from Germany (Weber 1958, Miera 1978), Belarus and Russia (Peklo 1987, Byshnev & Stavrovsky 1998) and other countries (Glutz von Blotzheim & Bauer 1993). The Red-Breasted Flycatcher is usually described as a half-cavity nester. We, however, divided nest sites into tree groups: half cavities, chimneys and shelves. Nest site types were connected with tree species and diameter. Using different nest sites is probably a result of the diversity of conditions and tree stands in Białowieża National Park. The choice of nesting tree species differed from that described in other studies in Europe, probably reflecting the species composition in the nesting

habitat. In northern Russia and Lithuania most birds bred in Alder *Alnus glutinosa*, in Beskidy Mountains in Beech *Fagus sylvatica*, Silver Fir *Abies alba* and Norway Spruce (Aleknonis 1976, Peklo 1987, Byshnev & Stavrovsky 1998). Similarly to Red-breasted Flycatcher, other hole-nesters in Białowieża national Park often use hornbeam and lime as a nesting tree (Wesołowski 1989).

Only four cases of using the same place in subsequent seasons for breeding could be anti-predator adaptation or result of changing features and quality of nest site after winter. Nests sites of Red-breasted Flycatcher are more open, often in dead trees and seems to be less durable than holes using by other cavity-nesting birds (Wesołowski 1995)

An average height of nests above the ground in Białowieża Forest (4.9 m) was higher than in Germany and Russia (Muller 1970, Cramp 1993, Glutz von Blotzheim & Bauer 1993) and similar to Lithuania (Aleknonis 1976). Other hole-nesters from Białowieża Forest, except Marsh Tit *Parus palustris*, build nests usually much higher than Red-breasted Flycatcher (Wesołowski 1989, 2002, Mitrus et al. 1996, Walankiewicz 1991, Czeszczewik & Walankiewicz 2003). There is lack of data about diameter of tree trunk used by Red-breasted Flycatcher in other parts of the Europe. In BNP on average this species breeds in much thinner trees than other hole nesters (Wesołowski 1989, 1996, Czeszczewik & Walankiewicz 2003). This is probably connected with the type of holes used by the species.

In the Beskidy Mountains, as in Białowieża, 30% of the nests were located in dead trees (Glutz von Blotzheim & Bauer 1993). Other sources do not mention the percentage of nests in dead or living trees. Other secondary hole-nesters breeding in BNP use dead trees in various rates, but other flycatcher species breed in dead trees less frequently (Wesołowski 1989, Czeszczewik & Walankiewicz 2003).

So far no data were published about hole-entrance exposure in Red-breasted Flycatcher nests. Similar to others hole-nesters from BNP most nest were directed to the south, southeast, southwest and east (Wesołowski 1989, 1996, Mitrus et al. 1996, Walankiewicz 1991, Czeszczewik & Walankiewicz 2003).

Usually the rims of the nests were visible from the ground and they were build in the shallow cavities with small bottom areas. This type of cavity distinctly differs from cavities preferred by other secondary hole-nesters in BNP (Walankiewicz 1991, Mitrus et al. 1996, Wesołowski 1989, Czeszczewik & Walankiewicz 2003). It could

be result of late arrival of this species, in the time when other *Ficedula* flycatchers and tits take many of holes and start to breed (Mitrus 2003, Czeszczewik & Walankiewicz 2003). On the other hand surplus of free cavities in Białowieża National Park was reported and low competition for nests sites were suggested (Wesołowski 1989, Walankiewicz 1991, Mitrus et al. 1996).

ACKNOWLEDGMENTS

We thank everybody who has helped to collect the data and handed them in Polish Nest Record Scheme during the past 27 years, especially Tomasz Wesołowski, Ludwik Tomiałojć, Wiesław Walankiewicz, Patryk Rowiński, Arkadiusz Szymura and Willem van Mannen. The research was granted by the University of Podlasie and supported by the Białowieża National Park authorities.

REFERENCES

- Aleknonis A. 1976. Biology of Red-breasted Flycatcher in SW part of Lithuania. Mat. IX Pribalt. Orn. Konf. Vilnius, 3–6.
- Balen van J. H. 1973. A comparative study of the breeding ecology of the Great Tit *Parus major* in different habitats. Ardea 61: 1–93.
- Bruns H. 1960. The economic importance of birds in forests. Bird Study 7: 193–208.
- Byshnev I., Stavrovsky K. D. 1998. On the biology of Red-breasted Flycatcher (*Ficedula parva*) in Berezinsky Nature Reserve. Subbuteo 1: 25–28.
- Cramp S. 1993. Birds of Europe the Middle East ad North Africa. The Birds of the Western Palearctic Vol. VII. Oxford Univ. Press.
- Czeszczewik D., Walankiewicz W. 2003. Natural nest sites of the Pied Flycatcher in a primeval forest. Ardea 91: 221–230.
- Flade M. 1997. Red-breasted Flycatcher. In: Hagemeyer W. J. M, Blair M J. 1997. The EBCC Atlas of European Birds. Poyser, London.
- Glutz von Blotzheim U. N., Bauer K. M. 1993. Handbuch der Vogel Mitteleuropas. Passeriformes. Aula Verlag, Wiesbaden.
- Haapanen A. 1965. Bird fauna of the Finish forests in relation to forest succession. Ann. Zool. Fenn. 2: 153–196.
- Miera C. 1978. Zur Brutbiologie des Zwergschnäppers. Der Falke 25: 120–127.
- Mitrus C. 2003. Temperature dependence of breeding phenology of the Collared Flycatcher *Ficedula albicollis* in the Białowieża Forest (NE Poland). Acta Ornithol. 38: 73–76.
- Mitrus C., Walankiewicz W., Czeszczewik D., Jabłoński P. M. 1996. Age and arrival date of Collared Flycatcher *Ficedula albicollis* males do not influence quality of natural cavities used. Acta Ornithol. 31: 101–106.
- Müller S. 1970. Das Brutvorkommen des Zwergschnäppers bei Warnemünde 1964–1968. Der Falke 17: 76–82.
- Newton I. 1998. Limitation population in birds. Academic Press, London.
- Peklo A. M. 1987. [Fauna of the flycatchers of the USSR]. Naukova Dumka, Kyiv.

- Rendell W. B., Robertson R. J. 1993. Cavity size, clutch-size and the breeding ecology of Tree Swallows *Tachycineta bicolor*. *Ibis* 135: 305–310.
- Slagsvold T. 1987. Nest site preference and clutch size in the Pied Flycatcher *Ficedula hypoleuca*. *Ornis Scand.* 18: 189–197.
- Tomiałojć L., Wesołowski T., Walankiewicz W. 1984. Breeding bird community of a primeval temperate forest (Białowieża National Park, Poland). *Acta Ornithol.* 20: 241–310.
- Tomiałojć L. 1991. Characteristics of old growth in the Białowieża Forest, Poland. *Nat. Areas J.* 11: 7–18.
- Walankiewicz W. 1991. Do secondary cavity nesting birds suffer more from competition for cavities or from predation in a primeval deciduous forest? *Nat. Areas J.* 11: 203–211.
- Weber H. 1958. Beobachtungen am Nest des Zwergschnappers (*Muscicapa parva*). *J. Ornithol.* 99: 160–172.
- Wesołowski T. 1989. Nest-sites of hole-nesters in primeval temperate forest (Białowieża National Park, Poland). *Acta Ornithol.* 25: 321–351.
- Wesołowski T. 1995. The loss of avian cavities by injury compartmentalization in a primeval European forest. *Condor* 97: 256–257.
- Wesołowski T. 1996. Natural nest sites of Marsh Tit (*Parus palustris*) in a primeval forest (Białowieża National Park, Poland). *Vogelwarte* 38: 235–249.
- Wesołowski T. 2002. Anti-predator adaptations in nesting Marsh Tits *Parus palustris*: the role of nest-site security. *Ibis* 144: 593–601.
- Wesołowski T., Tomiałojć L., Mitrus C., Rowiński P., Czeszczewik D. 2002. The breeding birds community of a primeval temperate forest (Białowieża National Park, Poland) at the end of 20th century. *Acta Ornithol.* 37: 27–45.

STRESZCZENIE

[Miejsca lęgowe muchołówki małej w naturalnym lesie (Białowieski Park Narodowy)]

Obserwacje prowadzono w latach 1977–2003 na terenie Białowieskiego Parku Narodowego. Większość gniazd (88 z 117) opisano w ostatnich czterech sezonach lęgowych (2000–2003), nato-

miast wcześniejsze dane pochodzą z Kartoteki Gniazd i Lęgów. Gniazda wyszukiwano obserwując samice podczas budowy, przeszukując miejsca wcześniej stwierdzonych par lub śledząc dorosłe ptaki w okresie karmienia piskląt. Dla większości gniazd opisano: gatunek drzewa, wysokość nad ziemią, pierśnicę, kierunek wlotu do dziupli, stan drzewa (martwe lub żywe), typ lokalizacji gniazda i dla niektórych średnicę dna i głębokość dziupli. Większość gniazd znajdowała się w drzewach żywych (70.2%), najwięcej w grabach *Carpinus betulus* i lipach *Tilia cordata* (Tab. 1). Wyróżniono następujące typy miejsc gniazdowych: półdziuple (płytkie dziuple z dużym otworem wejściowym – 46.4%), kominy (płytkie zagłębienia, zwykle w złamanym szczycie cienkiego drzewa – 26.4%) i półki (miejsca na zewnątrz pnia głównego, zwykle pod odstającym fragmentem kory – 27.3%, Fig. 3). Rodzaj miejsca gniazdowego zależał od gatunku drzewa. Większość kominów zlokalizowano w lipach (72.5%), podczas gdy półdziuple (76.5%) i półki (40.0%) charakterystyczne były dla grabów. Wlot do dziupli znajdował się najczęściej od strony południowej (Fig. 2). Gniazda były budowane stosunkowo nisko nad ziemią (średnio 4.9 ± 3.12 m). Najwyżej umiejscowione gniazda znaleziono w grabach (Tab. 1), ale różnice w wysokości gniazd w zależności od gatunku drzew nie były znaczące ($F_{7,79} = 0.91$, $p = 0.90$). Drzewa gniazdowe miały małą średnicę (31.2 ± 21.4 cm). Pierśnica zależała od gatunku drzewa ($F_{7,79} = 6.97$, $p < 0.001$), najgrubsze były sosny, a najcieńsze wiązy. Stwierdzono tylko cztery przypadki powtórnego wykorzystania tego samego miejsca lęgowego.