

Recent Literature on Bryophytes—108(4)

WILLIAM R. BUCK

New York Botanical Garden, Bronx, NY 10458-5126, U.S.A. e-mail: bbuck@nybg.org

BRUCE H. ALLEN

Missouri Botanical Garden, P.O. Box 299, St. Louis, MO 63166-0299, U.S.A. e-mail: bruce.allen@mobot.org

RONALD A. PURSELL

Department of Biology, 208 Mueller Laboratory, The Pennsylvania State University, University Park, PA 16802-5301, U.S.A. e-mail: rap10@psu.edu

- ADAMS, K. 2005. Changes in the bryophyte flora of eastern England. *Field Bryology* 85: 21–23, 1 tab.
- AICARDI, O. 2004. Contribution à l'inventaire de la bryoflore française (année 2003). *Bulletin de la Société Botanique du Centre-Ouest Nouvelle Série* 35: 337–342.
- AKIYAMA, H. 2005. Accidental resemblance in bryophytes. *Bryological Research* 8: 360–368. [Entirely in Japanese.]
- AKIYAMA, H. & T. MATSUI. 2005. *Distichophyllum yakumontanum* (Musci, Daltoniaceae), a new species from Yakushima Island, Japan. *Bryological Research* 8: 343–348.
- ALBERTOS, B. & R. GARILLETI. 2005. Breve panorama de los estudios de conservación de briófitos en Europa. *Conservación Vegetal* 9: 3–4.
- ALLEN, B., W. R. BUCK & R. A. PURSELL. 2005. Recent literature on bryophytes—108(2). *THE BRYOLOGIST* 108: 324–337.
- Anonymous. 2005. Zur Befruchtung von *Conocephalum conicum*. *Bryologische Rundbriefe* 89: 3.
- AONO, N., R. KOFUJI & M. HASEBE. 2004. [Abstract] Functional analysis of MADS-box genes in *Physcomitrella patens*. *Journal of Plant Research* 117(Supplement): 142.
- ARIKAWA, T. & M. HIGUCHI. 2004. [Abstract] The nucleotide sequences of *rbcL* gene of the genus *Takakia*—inter- and intraspecific variation and RNA editing. *Journal of Plant Research* 117(Supplement): 42.
- ASHIDA, Y. & M. KIMURA. 2005. *Neckeropsis obtusata* in Mie Prefecture, Japan. *Bryological Research* 8: 373–374. [In Japanese with locality data in English.]
- ASHIDA, Y., Y. TATEISHI & N. NISHIMURA. 2005. *Aulacomnium trichophyllum* found in Mie Prefecture, Japan. *Bryological Research* 8: 372–373. [In Japanese with locality data in English.]
- AUSTRHEIM, G., K. HASSEL & A. MYSTERUD. 2005. The role of life history traits for bryophyte community patterns in two contrasting alpine regions. *THE BRYOLOGIST* 108: 259–271.
- BALDWIN, L. K. & G. E. BRADFIELD. 2005. Bryophyte community differences between edge and interior environments in temperate rain-forest fragments of coastal British Columbia. *Canadian Journal of Forest Research* 35: 580–592.
- BARLOW, A. J., B. J. COMPTON, U. HERTEWICH, S. D. LORIMER & R. T. WEAVERS. 2005. Sesquiterpenes from the New Zealand liverwort *Lepidolaena hodgsoniae*. *Journal of Natural Products* 68: 825–831.
- BATES, J. 2005. Should we have a recording scheme for autecological attributes? *Field Bryology* 85: 30–31.
- BECKETT, R. P., M. MARSHALL & Z. LAUFER. 2005. Hardening enhances photoprotection in the moss *Atrichum androgynum* during rehydration by increasing fast- rather than slow-relaxing quenching. *Journal of Bryology* 27: 7–12.
- BELLAND, R. J. 1981. Ecology and Phytogeography of the Mosses of the Bonne Bay Region, Western Newfoundland. xii + 133 pp. Memorial University of Newfoundland, St. John's, Newfoundland, Canada. [Master's thesis.]
- BELLAND, R. J. 1984. The Disjunct Bryophyte Element of the Gulf of St. Lawrence Region: Glacial and Postglacial Dispersal and Migrational Histories. xvii + 269 pp. Memorial University of Newfoundland, St. John's, Newfoundland, Canada. [Ph.D. thesis.]
- BELLAND, R. J. 2005. A multivariate study of moss distributions in relation to environment in the Gulf of St. Lawrence region, Canada. *Canadian Journal of Botany* 83: 243–263.
- BERGAMINI, A. & N. SCHNYDER. 2005. *Funaria attenuata* (Dicks.) Lindb. und *F. obtusa* (Hedw.) Lindb. im Tessin wieder gefunden. *Meylania* 31: 32–34.
- BEZANILLA, M., P.-F. PERROUD, A. PAN, P. KLUEH & R. S. QUATRANO. 2005. An RNAi system in *Physcomitrella patens* with an internal marker for silencing allows for rapid identification of loss of function phenotypes. *Plant Biology* 7: 251–257.
- BOBORADJABOV, B. 2003. [Biological characteristics of the Pamir region (Tajikistan)]. *Izvestiya Akademii Nauk Tadzhikskoi SSR: Otdelenie Biologicheskikh Nauk* 150(3): 34–40. [In Russian with Tatar summary.]
- BOSANQUET, S. D. S. 2005. The changing bryophyte flora of south Wales. *Field Bryology* 85: 27–28.
- BOUDIER, P. 2004. Observations en Auvergne de quelques bryophytes épiphytes. *Bulletin de la Société Botanique du Centre-Ouest Nouvelle Série* 35: 347–352.
- BRAGGINS, J. 2005. Dame Ella Orr Campbell DNZM, Dip Tchng, MA (Hons), DSc (1910–2003). *Journal of Bryology* 27: 75–76.
- BRASSARD, G. R. 1970. The Mosses of Northern Ellesmere Island: Floristics and Bryogeography. xii + 246 pp. University of Ottawa, Ottawa, Canada. [Ph.D. thesis.]
- BRIDGLAND, J. P. 1986. The Flora and Vegetation of Cape Herschel, Ellesmere Island, N.W.T. xi + 149 pp. Memorial University of Newfoundland, St. John's, Newfoundland, Canada. [Master's thesis; includes bryophytes and lichens.]
- BRÜCKER, G., F. MITTMANN, E. HARTMANN & T. LAMPARTER. 2005. Targeted site-directed mutagenesis of a heme oxygenase locus by gene replacement in the moss *Ceratodon purpureus*. *Planta* 220: 864–874.
- BRYAN, V. S. 2005. Underlying specimens for the systematic treatment of the Ephemeraceae in the Bryophyte Flora of North America. *EVANSIA* 22: 13–29.
- BURYOVÁ, B. & A. J. SHAW. 2005. Phenotypic plasticity