



Seed Mix Performance and Cheatgrass Suppression on Arid Rangelands

Authors: Clements, Charlie D., Harmon, Dan N., and Blank, Robert R.

Source: Rangelands, 44(2) : 129-135

Published By: Society for Range Management

URL: <https://doi.org/10.1016/j.rala.2022.02.003>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Research Paper



Seed mix performance and cheatgrass suppression on arid rangelands

By Charlie D. Clements, Dan N. Harmon and Robert R. Blank

On the Ground

- The accidental and subsequent invasion of cheatgrass throughout millions of hectares of Intermountain West rangelands has truncated secondary succession by providing a fine-textured, early maturing fuel that has increased the chance, rate, spread, and season of wildfire.
- The restoration or rehabilitation of degraded rangelands throughout the Intermountain West is very challenging due to annual invasive species that exhibit high growth rates and seed production.
- The use of the pre-emergent herbicide, *Imazapic*, decreased cheatgrass densities >95% during the fallow year and before sowing seed the following fall during this study, which significantly reduced the cheatgrass competition for seedlings of seeded species.
- Seed mix performances were significantly higher in herbicide-treated plots than control plots for both sites for both years. Native, introduced, and native/introduced seed mixes were significantly more successful in the treated plots at the Bedell Flat site compared with the Antelope site for both years.
- Cheatgrass densities were significantly higher in the control plots at both sites for both years compared with herbicide/seed mix-treated plots.
- Success and failure of establishing perennial grasses in restoration or rehabilitation practices is highly dependent on proper seed and seed mix selections, seeding methodologies, and rates as well as favorable precipitation.

Keywords: *Bromus tectorum*, wildfire, restoration, herbicide, perennial grasses.

Rangelands 44():129–135

doi 10.1016/j.rala.2022.02.003

Published by Elsevier Inc. on behalf of The Society for Range Management. This is an open access article under the CC BY-NC-ND license (<http://creativecommons.org/licenses/by-nc-nd/4.0/>)

The restoration or rehabilitation of degraded rangelands throughout the Intermountain West is challenging due to annual invasive species that exhibit high growth rates and seed production.^{1–6} Invasive annual grasses, such as cheatgrass (*Bromus tectorum* L.), outcompete native perennial species for limited resources, especially in arid environments.^{4,7–10} The accidental and subsequent invasion of cheatgrass throughout millions of hectares of Intermountain West rangelands has truncated secondary succession by providing a fine-textured early maturing fuel that has increased the probability, rate, spread, and season of wildfire.^{11–13} The increase in wildfire size and frequency can significantly reduce recovery of native species and increase annual grass dominance.^{5,6,10,13–15} After wildfire, the reduction in perennial grass densities and increased abundance of cheatgrass can significantly influence fire behavior through increased cheatgrass densities and associated fuel loads.^{5,10,13,16} The best known method at suppressing cheatgrass and associated fuels is through the establishment of perennial grasses.^{5–7,10,17–22} Rehabilitation of cheatgrass-infested rangelands is needed to reduce wildfire threats, reduce soil erosion, allow succession to proceed, and improve grazing and wildlife resources.^{6,12} Aggressive and effective weed control methods are essential in reducing cheatgrass densities to improve seedling survivability of seeded species, as well as selecting seed mixes that will improve seedling success.^{5,6,19,20,23} Our study was initiated to test native, introduced, and native/introduced seed mix performance and the ability of these seed mixes to suppress cheatgrass densities.

Cheatgrass control and seeding

In the fall of 2014 and 2015, we applied *Imazapic* (Plateau) at 105 g ai/ha (6 oz/acre) to 24 m x 45 m plots (80 feet x 150 feet), with two replications and two control plot replications that did not receive herbicide application (Fig. 1), each year. Our two study sites were the Bedell Flat study site located 48 km (30 miles) north of Reno, Nevada and the Antelope study site located 210 km (130 miles) east of Reno, Nevada. There were eight plots total at each site. Both study sites burned in 1999 wildfires and the following seeding efforts failed (Fig. 2).

The Bedell Flat study site is dominated by cheatgrass with a sparse density of bottlebrush squirreltail (*Elymus ely-*




Figure 1. Schematic diagram of treated and control plots with seed mixes and randomized monitoring quadrats. I indicates introduced seed mix plots; N, native seed mix plots; M, introduced/native seed mix plots. The * represents an example of the randomized quadrats per plot.



Figure 2. Bedell Flat study site, August 2014, northern Nevada failed 1999 post wildfire seeding.

moides [Raf.] Swezey), Indian ricegrass (*Achnatherum hymenoides* [Roemer and Schultes] Barkworth), needle and thread grass (*Hesperostipa comata* [Trin. and Rupr.] Barkworth), bluegrass (*Poa secunda* J.S. Presl.), and Wyoming big sagebrush (*Artemisia tridentata* ssp. *wyomingensis* Beetle and A. Young). The site receives an average of 220 mm (8.7 inches) of annual precipitation (USDA-ARS unpublished data) and

the soil is mapped as a Graufels series, mixed mesic, Torripsammetic Haploxeroll, and developed in granitic residuum with an elevation of 1,545 m (5,098 feet).²⁴ The Antelope study site is dominated by cheatgrass with a sparse bluegrass presence. The site receives an average annual precipitation of 178 mm (7 inches; USDA-ARS unpublished data) and the soils are mapped as Hessing Wholan series, mixed

Table 1

Seed mixes and seeding rates of perennial grasses for the Bedell Flat and Antelope study sites in northern Nevada

Site	Seed mix	Rate	Seeds/area
Native mix			
Bedell Flat	'Anatone' bluebunch wheatgrass [<i>Pseudoroegneria spicata</i> (Pursh) A. Love ssp. <i>spicata</i>]	8 kg/ha (7 lb/acre)	257 seeds/m ² (24 seeds/feet ²)
Antelope	Bottlebrush squirreltail (<i>Elymus elymoides</i> [Raf.] Swezey)	8 kg/ha (7 lb/acre)	220 seeds/m ² (20 seeds/feet ²)
Bedell Flat	Sherman big blue grass (<i>Poa secunda</i> [J. Presl] formerly <i>Poa ampla</i>)	2.4 kg/ha (2 lb/acre)	300 seeds/m ² (28 seeds/feet ²)
Bedell Flat	Sandberg bluegrass (<i>Poa secunda</i> [J. Presl] formerly <i>Poa sandbergii</i>)	2.4 kg/ha (2 lb/acre)	330 seeds/m ² (31 seeds/feet ²)
Introduced mix			
Bedell Flat	'Hycres' crested wheatgrass (<i>Agropyron cristatum</i> L.)	4 kg/ha (3.5 lb/acre)	113 seeds/m ² (10 seeds/feet ²)
Antelope	Siberian wheatgrass (<i>Agropyron fragile</i> [Roth])	4 kg/ha (3.5 lb/acre)	105 seeds/m ² (9 seeds/feet ²)
Native/introduced mix			
Bedell Flat	'Anatone' bluebunch wheatgrass	4 kg/ha (3.5 lb/acre)	129 seeds/m ² (12 seeds/feet ²)
Antelope	Bottlebrush squirreltail	4 kg/ha (3.5 lb/acre)	110 seeds/m ² (10 seeds/feet ²)
Bedell Flat	Siberian wheatgrass	4 kg/ha (3.5 lb/acre)	105 seeds/m ² (9 seeds/feet ²)
Antelope	Sherman big bluegrass	2.4 kg/ha (2 lb/acre)	300 seeds/m ² (28 seeds/feet ²)
Bedell Flat	Sandberg bluegrass	2.4 kg/ha (2 lb/acre)	330 seeds/m ² (31 seeds/feet ²)

xeric, and silty to sandy loam with an elevation of 1,620 m (5,346 feet).²⁴

Treated plots were followed for 1 year after herbicide treatment, separated into thirds (i.e., 8 m x 45 m [26.2 feet x 150 feet]), and then seeded the following October with 3 separate seed mixes (native, introduced and native/introduced) using a Kincaid no-till drill (Kincaid Model 70, Kincaid Equipment, Haven, KS). Seeds were commercially acquired, and seed mixes and seeding rates at each study site included a native, introduced, and native/introduced seed mixes (Table 1). Each seed mix was also seeded on two replicated control plots for each year with the same dimensions that did not receive herbicide treatments (Fig. 1). Each plot received 10 randomly selected 1-m² (10.76 feet²) quadrats that were monitored for 2 years (Fig. 1). Two years post seeding the seeded species plant densities and cheatgrass plant densities were recorded. Data were analyzed using SAS JMP²⁵ with treatment, year, and interaction as fixed effects and the site, site x treatment, and year x site x treatment as random effects.

Results and Discussion

The application of imazapic at both study sites was highly successful at decreasing cheatgrass densities (Fig. 3). Seed mix performances were significantly ($P \leq 0.05$) higher in herbicide treated plots than control plots for both sites for both years ($n = 40$; Fig. 4). Native, introduced, and native/introduced seed mixes were significantly ($P \leq 0.01$) more successful in the treated plots at the Bedell Flat site compared

with the Antelope site in both years (Fig. 4). There were no significant differences in control plots between sites and years.

The Bedell Flat site recorded 7.9 (0.73/feet²), 7.6 (0.70/feet²), and 7.2 (0.67/feet²) individual perennial grasses/m² in the native, introduced, and native/introduced treated plots in year 1, respectfully, compared with 1.1/m² (0.10/feet²), 2.3/m² (0.21/feet²), and 2.8/m² (0.26/feet²) in year 1 at the Antelope site. In year 2, the Bedell Flat site recorded 7.9 (0.73/feet²), 10.7 (0.99/feet²) and 8.1 (0.75/feet²) individual perennial grasses/m² in the native, introduced, and native/introduced seed mixes, respectfully, compared with 1.7/m² (0.16/feet²), 4.2/m² (0.39/feet²), and 3.4/m² (0.32/feet²) at the Antelope site (Fig. 4). There was no significant difference in seed mix performance of herbicide treated plots at the Bedell Flat site in year 1, whereas in year 2 the introduced seed mix performed significantly ($P \leq 0.05$) better than the native and native/introduced seed mixes. The introduced seed mix was significantly ($P \leq 0.05$) more successful in year 2 than year 1 at the Bedell Flat site. The introduced and native/introduced seed mixes performed significantly ($P \leq 0.05$) higher in herbicide treated plots both years at the Antelope site (Fig. 4).

Amount and periodicity of precipitation plays a crucial role in the survivability and establishment of perennial bunchgrasses.^{4,5,18,27,33} The Bedell Flat study site received 270 mm (10.6 inches) of precipitation in year 1 and 327 mm (12.9 inches) in year 2, both were above average years. The Antelope study site received 223 mm (8.8 inches) of precipitation in year 1 and 216 mm (8.5 inches) in year 2, which was also above average for both years (USDA-ARS unpublished data).



Figure 3. Effective cheatgrass control using a pre-emergent herbicide, *Imazapic*, at the Antelope study site, June 2015, northern Nevada.




Figure 4. Perennial grass establishment for each treatment and year at our Antelope and Bedell Flat study sites in northern Nevada. Corresponding letters represent no significant difference ($P < 0.05$). N/I is Native/Introduced mix.

Cheatgrass densities were significantly ($P \leq 0.05$) higher in the control plots at both sites for both years compared with herbicide/seed mix treated plots (Table 2). The Antelope site had significantly ($P \leq 0.05$) more above-ground cheatgrass densities in the control plots compared with the Bedell Flat site for both years (Table 2). There were no significant differences in cheatgrass above-ground densities among the three seed mixes at the Bedell Flat site, while at the Antelope site

the introduced and native/introduced seed mixes had significantly ($P \leq 0.05$) less cheatgrass than the native seed mix plots (Table 2). Even though it is well reported that increased densities of perennial bunchgrasses reduce cheatgrass dominance, our study quantifies this reduction.

Restoration or rehabilitation of degraded big sagebrush communities in arid environments, especially those receiving ≤ 254 mm (10 inches) of annual precipitation, are very limited



Figure 5. Native seed mix plot 5 years post seeding, June 2020, Bedell Flat study site, northern Nevada.

Table 2

Cheatgrass (*Bromus tectorum*) densities after seed mix treatments at Bedell Flat and Antelope study sites in northern Nevada

Year	Site	Cheatgrass (<i>Bromus tectorum</i>) plants/m ²			
		Control	Native mix	Introduced mix	N/I mix
2014-15	Antelope	667 (a)	359 (b)	243 (c)	139 (c)
2014-15	Bedell Flat	161 (a)	25 (b)	18 (b)	22 (b)
2015-16	Antelope	640 (a)	338 (b)	140 (c)	160 (c)
2015-16	Bedell Flat	257 (a)	27 (b)	12 (b)	15 (b)

Note. Corresponding letters represent no significant difference ($P < 0.05$). N/I indicates native/introduced mix.

due to the lack of adequate precipitation.^{5,10,26} Near complete control of cheatgrass is necessary if desirable perennial grasses are to be successfully seeded and established.^{5,6,10,17,20,27,28} The use of pre-emergent herbicides to aid in the establishment of perennial grasses is an effective cheatgrass control practice.^{5,6,29} The use of the pre-emergent herbicide, *Imazapic*, decreased cheatgrass densities >95% during the fallow year and before sowing seed the following fall during our study, which significantly reduced the cheatgrass competition for seedlings of seeded species. This level of cheatgrass control is essential as cheatgrass outcompetes perennial seedlings at the seedling stage for limited moisture and nutrients as well as contributing to significant rangeland seeding failures.^{5,9,17,30-32}

Restoring or rehabilitating degraded Great Basin rangelands by re-establishing perennial grasses is extremely difficult because annuals such as cheatgrass maintain dominance on disturbed sites by exhibiting high growth rates and seed production.^{1,19,33-35} At our Bedell Flat study site, over the 2-

year period of our study, the level of perennial grass seedling germination, emergence, and establishment significantly reduced cheatgrass densities by as much as 95% in the introduced seed mix in year 2 with a low cheatgrass suppression level of 85% in the native seed mix plot in year 1. The native/introduced seed mix reduced cheatgrass by 86% in year 1 and 94% in year 2 (Table 2). The Antelope site, which is more arid, experienced less perennial grass establishment and therefore was less successful at suppressing cheatgrass. The native seed mix only reduced above-ground cheatgrass densities 47% in each of the 2 years of our study, and native/introduced seed mix reduced above-ground cheatgrass densities by 79% in year 1 and 75% in year 2. The introduced seed mix reduced above-ground cheatgrass densities by 64% and 78% in year 1 and 2, respectively (Table 2).

We started experimenting with native/introduced seed mixes nearly 3 decades ago to hedge our bets on above and below average precipitation years,³⁶⁻⁴⁰ and others have also reported on this approach to restore or rehabilitate degraded rangelands.^{7,41-45} Climate patterns throughout the Great Basin are highly variable and erratic,⁴⁶ and this erratic unpredictable pattern has significant impact on germination, emergence, and seedling establishment of seeded and nonseeded plant species.^{5,47,48} The two sites in our study received less than average precipitation >60% of the time over the 30+ years we have actively been recording precipitation (USDA-ARS unpublished data). Therefore, our study was designed to use introduced plant material that has been reported to be more successful in reseeding efforts if our sites received submarginal precipitation.^{1,17,33,43,49,50} Throughout our study the native seed mix performed very well at the Bedell Flat site (Fig. 5), while performing far less at the more arid Antelope site. As previously mentioned, perennial grasses

have a difficult time establishing when annual precipitation is ≤ 254 mm (10 inches), and especially true for native perennial grasses.^{5,10,26,45} Managers should be aware of such limitations when seeding arid rangelands. Prior research has reported that 2.5 to 3.0 individual perennial grasses/m² (0.23–0.28/feet²) is a desired density to maintain perennial vegetation and prevent cheatgrass dominance,^{27,51} which we achieved at the Bedell Flat site for all mixes for both years. This level of density was not achieved with the native mix at the Antelope site and struggled to meet the desired density levels with the introduced and native/introduced seed mixes. Our data suggests that to actively suppress cheatgrass densities and reduce its dominance, >4 individual perennial grasses/m² (0.37/feet²) are necessary to achieve this goal.

The use of introduced perennial grasses in restoration/rehabilitation efforts continues to be a focus of considerable debate. A wide range of concerns exists ranging from introduced perennial grasses significantly inhibiting the return of native species back to the site to the use of more adapted introduced perennial grasses that are essential in protecting watersheds, soil resources, and reducing cheatgrass dominance and further site degradation on the other edge of the spectrum.^{4,33,49,52–56} Success and failure of establishing perennial grasses in restoration or rehabilitation practices is highly dependent on proper seed and seed mix selections and rates, seeding methodologies, and favorable precipitation.^{5,20,27,56,57} A first step in restoring or rehabilitating degraded rangelands throughout the Great Basin is providing resource managers with effective weed control practices and proper seeding methodologies to combat such aggressive annual weeds as cheatgrass, therefore reducing cheatgrass densities and associated fuels that add further risk to watersheds, soil properties, grazing resources, and wildlife habitats.

Declaration of competing interest

The authors declare the following financial interests/personal relationships which may be considered as potential competing interests. The authors certify that they have no financial interest in the subject matter discussed in the manuscript. C. D. C., D. N. H., and R. R. B. are employees of USDA Agricultural Research Service and were associated with management decisions regarding the topic of this manuscript

References

- HARRIS G. Some competitive relationships between *Agropyron spicatum* and *Bromus tectorum*. *Ecol Monogr*. 1967; 37:89–111. doi:10.2307/2937337.
- WILSON AM, WONDERCHECK DE, GOEBEL CJ. Response of range grass seeds to winter environments. *J Range Manage*. 1974; 27:120–122. doi:10.2307/3896746.
- AGUIRRE L, JOHNSON DA. Influence of temperature and cheatgrass competition on seedling development of two bunchgrasses. *J Range Manage*. 1991; 44:347–354.
- HUMPHREY LD, SCHUPP EW. Competition as a barrier to establishment of a native perennial grass (*Elymus elymoides*) in alien annual grass (*Bromus tectorum*) communities. *J Arid Environ*. 2004; 58:405–422. doi:10.1016/j.aridenv.2003.11.008.
- YOUNG JA, CLEMENTS CD. *Cheatgrass: Fire and Forage on the Range*. University Nevada Press; 2009:348.
- CLEMENTS CD, HARMON DN, BLANK RR, WELTZ M. Improving seeding success on cheatgrass infested rangelands in northern Nevada. *Rangelands*. 2017; 39:174–181.
- ROBERTSON JH, PEARCE CK. Artificial reseeding and closed communities. *Northwest Sci*. 1945; 19:58–66.
- EVANS RA, HOBO HR, ECKERT RE, YOUNG JA. Functional environment of downy brome communities in relation to weed control and revegetation. *Weed Sci*. 1970; 18:154–162.
- MELGOZA AG, NOWAK RS, TAUSCH RJ. Soil water exploitation after fire: competition between *Bromus tectorum* (cheatgrass) and two native species. *Oecologia*. 1990; 83:7–13.
- CHAMBERS JC, ROUNDY BA, BLANK RR, MEYER SE, WHITTAKER A. What makes Great Basin sagebrush ecosystems invulnerable by *Bromus tectorum*? *Ecol Monogr*. 2007; 77:117–145.
- PECHANEC JF, HULL AC. Spring forage loss through cheatgrass fires. *National Wool Grower*. 1945; 35:13.
- CLEMENTS CD, YOUNG JA. A viewpoint: rangeland health and mule deer habitat. *J Range Manage*. 1997; 60:129–138. doi:10.2307/4002369.
- PELLANT M, ABBEY B, KARL S. Restoring the Great Basin Desert, U.S.A.: integrating science, management, and people. *Environ Monit Assess*. 2004; 99:169–179.
- MILLER RF, ROSE JA. Fire history and western juniper encroachment in sagebrush steppe. *J Range Manage*. 1999; 52:550–559.
- BROOKS ML, D'ANTONIO CM, RICHARDSON DM, GRACE JB, KEELEY JE. Effects of invasive alien plants on fire regimes. *BioScience*. 2004; 54:677–688.
- YOUNG JA, EVANS RA, ECKERT RE, CHEATGRASS KAY BL. *Rangelands*. 1987; 9:266–270.
- HULL AC, STEWART G. Replacing cheatgrass by reseeding with perennial grasses on southern Idaho ranges. *J Agron*. 1948; 40:226–262.
- CHAMBERS JC, MILLER RF, BOARD DI, ET AL. Resilience and resistance of sagebrush ecosystems: implications for state and transition models and management treatments. *Rangeland Ecol Manag*. 2014; 67:440–454.
- DAVIES KW, NAFUS AM, SHELEY RL. Non-native competitive perennial grass impedes the spread of an invasive annual grass. *Biol Invasions*. 2010; 12:3187–3194.
- DAVIES KW, JOHNSON DD, NAFUS AM. Restoration of exotic annual grass-invaded rangelands: Importance of seed mix composition. *Invasive Plant Sci Manage*. 2014; 7:247–256.
- BLANK RR, MORGAN T. Suppression of *Bromus tectorum* L. by established perennial grasses: potential mechanisms—part one. *Applied Environ Soil Sci*. 2012; 9.
- BLANK RR, CLEMENTS CD, MORGAN TA, HARMON DN, ALLEN F. Suppression of cheatgrass by perennial bunchgrasses. *Rangeland Ecol Manag*. 2020; 73(6):766–771.
- DAVIES KW, BOYD CS, JOHNSON DD, MADSEN MD. Success of seeding native compared with introduced perennial vegetation for revegetating medusahead-invaded sagebrush rangelands. *Rangeland Ecol Manag*. 2015; 68:224–230. doi:10.1016/j.rama.2015.03.004.
- SOIL SS, Natural Resources Conservation Service, United States Department of Agriculture. Web Soil Survey. Accessed January 4, 2022. <http://websoilsurvey.sc.egov.usda.gov/>.
- SAS INSTITUTE. *JMP Version 12.0.1*. Cary, NC, USA: SAS Institute Inc; 2017.

26. MONSON SB. The competitive influences of cheatgrass (*Bromus tectorum*) on site restoration. In: Monsen SB, Kitchen SG, eds. *Proceedings Ecology and Management of Annual Rangelands*. Ogden, UT: USDA-Forest Service Gen; 1994:43–50.
27. EVANS RA, YOUNG JA. Weed control-revegetation systems for big sagebrush-downy brome rangelands. *J Range Manage.* 1977; 30:331–336. doi:10.2307/3897714.
28. JORDAN GL. Planting limitations for arid, semi-arid and salt desert shrublands. *Proceedings of Managing Intermountain Rangelands-Improvement of Range and Wildlife Habitats*; 1983:11–16.
29. ECKERT RE, ASHER JE, CHRISTENSEN MD, EVANS RA. Evaluation of the atrazine-fallow technique for weed control and seedling establishment. *J Range Manage.* 1974; 27:288–292.
30. EVANS RA. Effects of different densities of downy brome (*Bromus tectorum*) on growth and survival of crested wheatgrass (*Agropyron desertorum*) in the greenhouse. *Weeds.* 1961; 9:216–223. doi:10.2307/4040408.
31. ROBERTSON JH, ECKERT JR RE, BLEAK AT. Responses of grasses seeded in an *Artemisia tridentata* habitat in Nevada. *Ecology.* 1966; 47:187–194. doi:10.2307/1933764.
32. RAFFERTY DL, YOUNG JA. Cheatgrass competition and establishment of desert needlegrass seedlings. *J Range Manage.* 2002; 55:70–72. doi:10.2307/4003265.
33. SVEJCAR T, BOYD C, DAVIES K, HAMERLYNCK E, SVEJCAR L. Challenges and limitations to native species restoration in the Great Basin, USA. *Plant Ecol.* 2017; 218:81–94.
34. HARRIS AG, WILSON AM. Competition for moisture among seedlings of annual and perennial grasses as influenced by root elongation and low temperature. *Ecology.* 1970; 51:530–534.
35. PLATT K, JACKMAN ER. The cheatgrass problem in Oregon. *Oregon State University Extension Bulletin.* 1946; 668.
36. YOUNG JA, CLEMENTS CD. Cheatgrass control and seeding. *Rangelands.* 2000; 22:3–7.
37. CLEMENTS CD, GREY K, YOUNG JA. Forage kochia: to seed or not to seed. *Rangelands.* 1997; 19:29–31.
38. CLEMENTS CD, YOUNG JA, GRAY K. Cheatgrass invasion and mule deer habitat. *African J Range For Sci.* 2003; 20:161.
39. CLEMENTS CD, MCCUIN G, SHANE RS, McADOO K, HARMON DN. Wildfire rehabilitation and restoration: triage in the pursuit of resilience. *Rangelands.* 2009; 31:30–35.
40. CLEMENTS CD. The need to improve mule deer populations: habitat conversion. *Progressive Rancher.* 2021; 21:18–20.
41. DAVIES KW, BOYD CS, MADSEN MD, HULET KA. Evaluating a seed technology for sagebrush restoration across an elevation gradient: support for bet hedging. *Rangeland Ecol Manag.* 2018; 71:19–24. doi:10.1016/j.rama.2017.07.006.
42. ASAY KH, CHATTERTON NJ, JENSEN KB, JONES TA, WALDRON BL, HORTON WH. Breeding improved grasses for semi-arid rangelands. *Arid Land Res Manag.* 2003; 17:469–478.
43. PLUMMER AP, HURD RM, PEARSE CK. *How to reseed Utah range lands.* USDA, Forest Service Research Paper INT-1. Ogden, UT: Intermountain Forest and Range Experiment Station; 1943:1–22.
44. PEARSE CK, PLUMMER AP, SAVAGE D. Restoring the range by reseeding. In: *Yearbook of Grass.* U. S. Department of Agriculture; 1948:227–233.
45. ASAY K H, HORTON WH, JENSEN KB, PALAZZO AJ. Merits of native and introduced Triticeae grasses on semiarid rangelands. *Can J Plant Sci.* 2001; 81:45–52. doi:10.4141/P99-131.
46. BARBOUR MG, BILLINGS WD. *North American Terrestrial Vegetation.* Cambridge University Press; 1988:1–434.
47. CALL CA, ROUNDY BA. Perspectives and processes in revegetation of arid and semiarid rangelands. *Rangeland Ecol Manag.* 1991; 44:543–549. doi:10.2307/4003034.
48. HARDEGREE SP, FLERCHINGER GN, VAN VACTOR SS. Hypothermal germination response and the development of probabilistic germination profiles. *Ecol Mod.* 2003; 167:305–322.
49. ASAY KH, CHATTERTON NJ, JENSEN KB, JONES TA, WALDRON BL, HORTON WH. Breeding improved grasses for semi-arid rangelands. *Arid Land Res Manag.* 2003; 17:469–478.
50. WALDRON BL, MONACO TA, JENSEN KB, HARRISON RD, PALAZZO AJ, KULBETH JD. Coexistence of native and introduced perennial grasses following simultaneous seeding. *Agron J.* 2005; 97:990–996. doi:10.2134/agronj2004.0265.
51. EVANS RE, YOUNG JA. Effectiveness of rehabilitation practices following wildfire in a degraded big sagebrush downy brome community. *J Range Manage.* 1978; 31:185–188.
52. YOUNG JA, McLAIN JL. Grazing livestock as a management tool in the intermountain west. *Proc Am Soc Anim Sci. Western Section. San Luis Obispo, CA.* 1997; 48:3–7.
53. PELLANT M. The cheatgrass-wildfire cycle-are there any solutions?. *Proceedings Cheatgrass Invasions, Shrub Die-off and Other Aspects of Shrub Biology and Management*; 1990:11–18.
54. PYKE DA. Rangeland seedings and plantings: exotics or natives? Edge WD, Olson-Edge SL, eds. *Proceedings of a Symposium on Sustaining Rangeland Ecosystems* Oregon State University; 1996:32–44.
55. RICHARDS RT, CHAMBERS JC, ROSS C. Use of native plants on federal lands: policy and practice. *J Range Manage.* 1998; 51:625–632. doi:10.2307/4003603.
56. PILLIOD DS, WELTY J, TOEVS GR. Seventy-five years of vegetation treatments on public rangelands in the Great Basin of North America. *Rangelands.* 2017; 39:1–9.
57. DAVIES KW, JOHNSON DD. Established perennial vegetation provides high resistance to re-invasion by exotic annual grasses. *Rangel Ecol Manag.* 2017; 70:754–784.

Authors are from: Rangeland Scientist, Agricultural Research Technician, and Soil Scientist (retired), USDA, Agricultural Research Service, Great Basin Rangelands Research Unit, Reno, NV, 89512 USA