



Smutgrass (*Sporobolus indicus*) control in bahiagrass is improved with applications of herbicide and fertilizer

Authors: Shay, Nicholas J., Baxter, Lisa L., Basinger, Nicholas T., Schwartz, Brian M., and Belcher, Jason

Source: Weed Technology, 36(5) : 700-707

Published By: Weed Science Society of America

URL: <https://doi.org/10.1017/wet.2022.73>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Research Article

Cite this article: Shay NJ, Baxter LL, Basinger NT, Schwartz BM, Belcher J (2022) Smutgrass (*Sporobolus indicus*) control in bahiagrass is improved with applications of herbicide and fertilizer. *Weed Technol.* **36**: 700–707. doi: [10.1017/wet.2022.73](https://doi.org/10.1017/wet.2022.73)

Received: 4 July 2022

Revised: 12 September 2022

Accepted: 13 September 2022

First published online: 30 September 2022

Associate Editor:

Drew Lyon, Washington State University

Nomenclature:

Hexazinone; indaziflam; bahiagrass; *Paspalum notatum* Alain ex Fluggé; smutgrass; *Sporobolus indicus* (L.) R. Br.






Keywords:

Integrated weed management plan; forages; pasture; agricultural pesticides; fertility; economics

Author for Correspondence:

Lisa L. Baxter, State Forage Extension Specialist, Department of Crop and Soil Sciences, The University of Georgia, 2360 Rainwater Road, Tifton, GA 31794.
Email: baxterl@uga.edu

Smutgrass (*Sporobolus indicus*) control in bahiagrass is improved with applications of herbicide and fertilizer

Nicholas J. Shay¹ , Lisa L. Baxter² , Nicholas T. Basinger³ ,
Brian M. Schwartz⁴  and Jason Belcher⁵ 

¹Graduate Student, Department of Crop and Soil Sciences, The University of Georgia, Tifton, GA, USA; ²State Forage Extension Specialist, Department of Crop and Soil Sciences, The University of Georgia, Tifton, GA, USA; ³Assistant Professor, Department of Crop and Soil Sciences, The University of Georgia, Athens, GA, USA; ⁴Professor, Department of Crop and Soil Sciences, The University of Georgia, Tifton, GA, USA and ⁵Stewardship and Development Manager, Eastern US, Envue, Vegetation Management, Forestry, Range and Pasture, Cary, NC, USA

Abstract

Smutgrass is an invasive weed species that can quickly outcompete bahiagrass because of its aggressive growth, prolific seed production, and rhizomatous nature. Total renovation of bahiagrass pastures or hayfields is generally not a feasible or economically viable option for most producers. Therefore, controlling the continual spread of smutgrass will require an integrated weed management (IWM) plan that incorporates multiple strategies. The objective of this study was to test the interactions of herbicides and fertilizers on smutgrass control in bahiagrass and determine the most efficacious and economical IWM plan for low-input bahiagrass systems. This research was conducted on a mixture of ‘Tifton 9’ and ‘Pensacola’ bahiagrass at the Alapaha Beef Station in Alapaha, GA. The study design was a randomized complete block with a three-by-four factorial treatment arrangement with six replications. Fertility treatments included 56 kg N ha⁻¹ (ammonium nitrate, 34% N) + 56 kg K₂O ha⁻¹, 56 kg N ha⁻¹, and an unfertilized control. Smutgrass was reduced to <15% ground coverage when a postemergent herbicide was applied. The addition of a preemergent herbicide and/or fertilizer further reduced the coverage of smutgrass ($P < 0.01$). As smutgrass declined, the bahiagrass ground coverage increased; other vegetation and dead material did not differ by treatment. Generally, herbage accumulation and crude protein were only affected following the second N application ($P < 0.01$). Treatments that included preemergent (indaziflam) and postemergent (hexazinone) herbicides in addition to N and K₂O resulted in an improved bahiagrass stand as timely weed suppression removed competition, while fertilizer provided essential nutrients for optimum growth to fill in the gaps. Combining herbicide and fertilizer is a more economical option for producers when compared to a complete bahiagrass renovation.

Introduction

Bahiagrass is a popular low-input forage option for livestock producers in the Coastal Plains Region of the southeastern United States because of its low fertility and input requirements for sustained growth in sandy soils (Beaty et al. 1960). It is a reliable, warm-season perennial forage with above-average persistence in adverse climatic conditions as well as resistance to most diseases and pests (Chambliss and Sollenberger 1991; Dias et al. 2018). Bahiagrass is well adapted to the southern half of Georgia, outperforming other warm-season forages where extreme temperatures as well as intermittent flooding or drought can otherwise limit biomass and nutritive value. One disadvantage of bahiagrass production systems is the lack of herbicide options to control highly competitive grass weeds and non-native invasive species that can easily overtake the stand if left unmanaged. Mitigating performance losses to bahiagrass, such as herbage accumulation (HA) and groundcover, will require effectively utilizing novel pest management strategies to deter encroachment of these opportunistic weeds.

Smutgrass is one of the invasive, non-native weeds that has become a major pest in perennial grasslands throughout the US Southeast (Rana et al. 2012). This bunch-type, warm-season perennial grass invades around 1.6 million ha of permanent and temporary pastures across the southeastern United States (Wallau et al. 2010). The name is derived from the dark-colored fungus [*Drechslera ravenelii* (M.A. Curtis ex Berk.)] present on the inflorescence that closely resembles smut from a fireplace or chimney (Mislevy et al. 1999). Previous research has identified two varieties of smutgrass populations: small smutgrass [*S. indicus* (L.) R. Br. var. *indicus*] and giant smutgrass [*S. indicus* var. *pyramidalis* (P. Beauv.) Veldkamp]. Small smutgrass is the dominant variety and has been found in 23 states, whereas giant smutgrass is most predominant in Florida. Regardless of the population,

© The Author(s), 2022. Published by Cambridge University Press on behalf of the Weed Science Society of America. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution and reproduction, provided the original article is properly cited.



smutgrass grows well in a wide array of environmental conditions, grows rapidly in vegetative and reproductive stages, continuously produces seed, and displays discontinuous germination.

Previous research has been conducted in Florida on controlling smutgrass in bahiagrass pasture systems. Initially, viable control options were focused on isolating responses to single variables such as mowing, fertility management, or intensive rotational grazing (Ferrell and Mullahey 2006; Kemp and King 2001). McCaleb et al. (1963) first recommended that dalapon be applied at 2.27 kg ai ha⁻¹ in 379 L of water followed by mowing to 7.62 cm every week starting in August and continuing for 13 consecutive weeks. This controlled smutgrass up to 85% without serious injury to the bahiagrass. Later studies discouraged the use of mowing for smutgrass control, as it can broadcast smutgrass seeds and may not be economically feasible with rising fuel costs (Ferrell and Mullahey 2006; Mislevy et al. 1999; Sellers and Ferrell 2011; Sellers et al. 2020). Dalapon is no longer recommended and was federally deregistered for use in pastures in the 1980s. Following this decision, DuPont filed for a federal label for hexazinone (Velpar L[®]; Bayer Crop Science, Whippany, NJ) for smutgrass control in pastures and hayfields (Mislevy et al. 1999).

The efficacy of hexazinone on smutgrass in bahiagrass and bermudagrass [*Cynodon dactylon* (L.) Pers.] has been researched thoroughly in Florida (Ferrell et al. 2006; Mislevy et al. 2002; Nolte 2017; Sellers and Ferrell 2011; Sellers et al. 2020; Wilder et al. 2008). Research indicates that the best timing for application is during conditions that are both warm and wet. The mode of action for hexazinone requires rainfall to allow the chemical to infiltrate the soil where it can be absorbed via root uptake. However, precipitation events exceeding 76 mm may lead to lower rates of efficacy as well as possible soil leaching. To achieve >90% control, Sellers and Ferrell (2011) recommended an application within 7 d before a rainfall event. It is also important that applications be made during the growing season when grasses are actively growing, as hexazinone requires movement into the xylem for translocation as a photosynthesis disruptor (photosystem II inhibitor) so as to achieve maximum herbicidal efficacy. Herbicide options for selective smutgrass control in bahiagrass are currently limited to hexazinone. Additionally, hexazinone presents potential challenges as an herbicide option because of cost to low-input producers as well as initial injury to bahiagrass; these challenges could have an impact on its competitiveness with other opportunistic weeds. In response to these disadvantages, research should explore new technologies such as indaziflam (Rezilon[®]; Bayer Crop Science, Whippany, NJ). Although a common preemergence herbicide in the turf industry, little is known about the use of indaziflam on smutgrass or other perennial grass weed species in bahiagrass. Previously, pendimethalin (Prowl H₂O; BASF Plant Science, Raleigh, NC) was the only recommended preemergence herbicide for use in bahiagrass in Georgia, but Prowl H₂O had little impact on smutgrass control (Sellers et al. 2020). Rezilon[®] was recently registered by Bayer Crop Science for use in warm-season grass pastures and hayfields. Hurdle et al. (2020) investigated the impact of indaziflam on bermudagrass forage production. The bermudagrass tolerated the indaziflam applications at the recommended rates, and no crop injury was reported. However, the impact of indaziflam on bahiagrass has yet to be determined, and more research will be required.

An IWM plan for removing smutgrass from existing bahiagrass stands must combat rhizomatous growth, as well as the continuous and prolific seed production throughout the growing season (Mislevy et al. 2002). Isolated treatments with singular uses of

herbicide or fertilizer may not be adequate for controlling smutgrass in the attempt to improve bahiagrass forage systems. Many nontargeted species could emerge unexpectedly with a broad range of responses outside the scope of targeted treatments (Kemp and King 2001). Thus, the importance of a preemptive plan that expands the range of management in complex forage systems cannot be overemphasized. Few studies have addressed the importance of fertilizing with nitrogen (N) and potassium (K) as part of an IWM plan to improve herbage accumulation and increase bahiagrass vigor by giving it a competitive advantage over weed species (Beaty et al. 1974; Silveira et al. 2017; Yarborough et al. 2017). Although species differ in their resource needs for space, light, water, and nutrients, changes in these resources can increase the competition between these species (Kemp and King 2001). Therefore, combining herbicides and fertilizer should improve productivity of bahiagrass by eliminating weed competition and increasing plant density of bahiagrass. However, there is much to learn about the interactive effects of an integrated management system. There is currently a paucity of research on how bahiagrass forage systems respond to collective herbicide and fertilizer treatments. The objectives of this experiment were to (1) evaluate the interactive effects of herbicide and fertilizer applications for controlling smutgrass and (2) determine the most efficacious and economical IWM plan for low-input bahiagrass systems in the Coastal Plains region of Georgia.

Materials and Methods

Description of Research Site

This research was conducted at the University of Georgia Alapaha Beef Station in Alapaha, GA, (31.58° N, 83.58° W; 81 m elev) from April through October in 2020 and 2021. The experimental site (Figure 1) was located in a previously established Tifton-9 and Pensacola bahiagrass pasture with a preexisting population of small smutgrass (percent visual groundcover of smutgrass in location 1 (initiated 2020): average = 42%; range = 20% to 80%; location 2 (initiated 2021): average = 27%; range = 2% to 100%). The experimental areas were fenced off to exclude grazing. The research site is nearly level (<2% slope) and primarily composed of Alapaha loamy sand (loamy, siliceous, subactive, thermic Arenic Plinthic Paleaquults) and Rutledge loamy sand (sandy, siliceous, thermic Typic Humaquepts), with an average soil pH of 5.0 (Web Soil Survey 2022).

Experimental Design and Treatments

The experiment was arranged in a randomized complete block design with a four-by-three factorial arrangement of treatments and six replications. Treatments included 4 herbicide (factor a) and 3 fertilizer (factor b) combinations, totaling 12 treatment combinations. Treatment combinations were randomly assigned to plots within each replicate. Each 2-m by 5-m plot was surrounded by 1-m alleyways on all sides for distinction.

Herbicide treatment levels consisted of an unsprayed control, preemergence, postemergence, and a combination of both preemergence and postemergence (PRE+POST). Indaziflam was applied preemergence at 0.058 kg ai ha⁻¹ on April 7, 2020 and March 15, 2021. Hexazinone was applied postemergence at 0.98 kg ai ha⁻¹ on August 7, 2020 and August 30, 2021 following harvest 4 (details below in forage sampling section). The combination (PRE+POST) herbicide treatment received both indaziflam and hexazinone applications as previously described. All herbicide treatments were applied using a tractor-mounted, 1.83-m boom sprayer with shield and TeeJet TP8003VS




Figure 1. Experimental site location 1 and location 2 at the UGA Alapaha Beef Unit near Alapaha, GA (31.58° N, 83.58° W). Research initiated: location 1, 2020; location 2, 2021.




Figure 2. Cumulative monthly rainfall from January to December for 2020 and 2021 in Alapaha, GA. The 100-yr historical average and data were collected from Georgia Weather Network (<http://www.georgiaweather.net>).

nozzles (TeeJet Technologies Inc., Glendale Heights, IL) calibrated to deliver 205.7 L ha⁻¹.

Ideally, the herbicide applications would have been made earlier each season. The preemergence applications were delayed in 2020 because university regulations initially prohibited any research

activities following the onset of COVID-19 restrictions. The post-emergence application should have been made in June or July; however, in 2020 there was insignificant precipitation forecasted to activate the hexazinone, and in 2021 the plots were flooded and inaccessible in 2021; therefore, the applications were delayed (Figure 2).

Fertilizer treatment levels consisted of an unfertilized control, nitrogen only (N), and nitrogen plus potassium (N+K). Fertilizers were hand-applied following green-up (April 7, 2020; April 23, 2021), and following harvest 3 (July 12, 2020; July 16, 2021). Each fertilizer application included 56.04 kg N ha⁻¹ (applied as ammonium nitrate, 34% N) or 56 kg N ha⁻¹ (applied as ammonium nitrate, 34% N) + 56 kg K₂O ha⁻¹ (applied as muriate of potash; N+K). The fertilizer treatments were below the recommendations provided by the University of Georgia Feed and Environmental Water Laboratory in Athens, GA, but are typical of what most bahiagrass fields would receive in South Georgia (Kissel and Sonon 2008).

Forage Sampling and Nutritive Value Analyses

Plots were harvested every 4–6 wk from April until winter dormancy (October). Plot borders were mowed to 8 cm prior to each harvest. All plots were visually evaluated for groundcover of bahiagrass, smutgrass, and other plant species before they were harvested to 8 cm with a zero-turn mower with a bagger attachment. The collected material from each plot was weighed using a tarp and tripod before a subsample was collected for dry-matter (DM) determination and nutritive value analysis. Post-harvest, residual forage was removed with the same mower (and to the same height) used for harvest. Subsamples were dried in a forced-air dryer at 55 C for 7 d before grinding to pass through a 1-mm screen in a Thomas Model 4 Wiley Mill (Thomas Scientific, Philadelphia, PA) followed by a Foss CT-293 Cyclotec Cylcone Mill (Foss Analytics, Eden Prairie, MS) with 1-mm screen (McIntosh et al. 2022). Ground samples were scanned for nutritive value using the 2020 grass hay calibration provided by the NIRS Forage and Feed Testing Consortium (NIRSC 2020). Samples were scanned on a Foss DS2500 near-infrared spectrometer (Metrohm USA Inc., Riverview, FL) that was standardized to the NIRSC master instrument to ensure prediction accuracy. Nutritive value data were reported with predictions fitting the allowable global $H < 3.0$ statistical comparison with the overall calibration population (Murray and Cowe 2004). Total digestible nutrients (TDN) were calculated using the grass equation provided in Moore and Undersander (2002) as follows using data obtained from NIRS analyses:

$$TDN = (NFC \times .98) + (CP \times .87) + (FA \times .97 \times 2.25) + NDFn \times NDFDp/100 - 10$$

where *NFC* is nonfibrous carbohydrate (% of DM) = 100 – (*NDFn* + *CP* + *EE* + ash), *CP* is crude protein (% of DM), *FA* is fatty acids (% of DM) = ether extract (*EE*) – 1, *NDFn* is nitrogen free *NDF* = *NDF* – *NDFCP*, otherwise estimated as *NDFn* = *NDF* × .93, and *NDFDp* is 22.7 + .664 *NDFD*, where *NDFD* is 48-h in vitro *NDF* digestibility (% of *NDF*).

Statistical Analyses

Data were analyzed by restricted maximum likelihood using PROC MIXED in SAS 9.4 (Littell et al. 2006). A Kenward–Rodgers adjustment was applied to correct the denominator degrees of freedom, ensuring appropriate standard errors and F statistics for each model. Multiple covariance structures were tested, and the Bayesian's Information Criterion indicated that Autoregressive (1) was the best fit. Differences in HA, nutritive value, and botanical composition were examined within harvest. Fixed effects included treatment,

year, and treatment × year. Replicates were the random effect. Means were compared using the LSMEANS procedure with Tukey–Kramer adjustment ($P \leq 0.05$). Differences were considered significant at $P \leq 0.05$.

Economic Analyses

An economic analysis was made with respect to market costs of the examined treatments and compared to a total bahiagrass renovation. All fertilizer prices were collected from DTN in January 2022. All herbicide prices were collected from Chemical Warehouse herbicide costs in the spring of 2022 (Chemical Warehouse 2022; Quinn 2022). The costs associated with each treatment were calculated by multiplying the quantities of inputs used by the market prices for the region.

Bahiagrass renovations were calculated by modifying the University of Georgia Extension, College and Agricultural and Environmental Sciences–Applied Economics, 2018 hybrid bermudagrass hay–nonirrigated establishment budget to reflect fertilizer rates and seed costs recommended by UGA bahiagrass management bulletin (Hancock et al. 2010). This budget included market costs for a glyphosate burndown, 2,4-D for postemergence application, 'TifQuik' bahiagrass seed costs (Hancock Seed Company, Dade City, FL), fertilizer at planting and after first mowing, and fuel. In addition, the budget included estimated costs for repairs, maintenance, labor, and interest (Lacy et al. 2016).

Results and Discussion

Weather

Daily air temperatures followed 100-yr historical monthly average temperatures for both 2020 and 2021. Minimum monthly average temperatures for April, July, and October, representing the beginning, mid-point, and the end of the growing season were 12 C, 22 C, and 16 C, respectively. Maximum monthly average temperatures for April, July, and October were 25 C, 34 C, and 28 C respectively (Georgia Weather Network 2021). Rainfall varied year to year, with precipitation following historical average rainfall of 715 mm beginning in April through October for 2020 and above average for 2021 with 1,103 mm (Figure 2) (Georgia Weather Network 2021).

Groundcover Percentage

Year × treatment did not interact to affect smutgrass, bahiagrass, other, or dead groundcover, except at harvest 4 where the interaction was a difference of magnitude ($P < 0.01$). Again, the data were pooled over both years and analyzed within harvest. Initial observations were recorded at harvest 0 before preemergence herbicide application where no differences were reported ($P = 0.59$; Table 1; March). Similarly, no differences were reported for harvest 1 ($P = 0.82$; Table 1; April) following preemergence and fertilizer application. Indaziflam has been well established as a preemergence herbicide in turf, where it reduces seedling emergence, and thus it is not surprising that it did not affect preexisting smutgrass groundcover. No differences in smutgrass cover were found at harvest 2 (June) ($P = 0.84$) and harvest 3 (July) ($P = 0.12$). Despite the fertilizer application following harvest 3, harvest 4 (August) also resulted in no differences among treatments ($P = 0.28$). The postemergence herbicide was applied following harvest 4, and the effects were again seen in subsequent harvests (September and October). Smutgrass groundcover numerically

Table 1. Effect of treatment on smutgrass groundcover percentage of bahiagrass-dominant pasture (includes smutgrass, bahiagrass, other weeds) grown at the Alapaha Beef Unit near Alapaha, GA, during 2020 and 2021.

Treatment ^a	Harvest 0 ^b	Harvest 1	Harvest 2	Harvest 3	Harvest 4	Harvest 5	Harvest 6
% Groundcover							
Control	42	21	27	13	20	27 ab ^c	55 a
PRE	38	27	30	9	25	32 a	51 a
POST	39	34	27	18	26	4 cd	13 bc
PRE+POST	37	29	20	20	23	12 abcd	3 c
N	32	28	25	18	35	26 ab	46 a
PRE, N	33	32	23	14	32	13 abcd	30 ab
POST, N	35	31	30	21	22	9 bcd	0 c
PRE+POST, N	30	34	29	23	17	5 cd	4 c
N+K	30	28	19	12	25	24 abc	46 a
PRE, N+K	36	35	23	22	23	27 ab	45 a
POST, N+K	26	24	29	25	23	4 cd	8 bc
PRE+POST, N+K	33	37	24	18	13	2 d	3 c
SEM ^d	10.69	6.44	5.71	7.80	8.70	4.73	5.91

^aPRE (preemergence; indaziflam, 0.28 kg ai ha⁻¹). POST (postemergence; hexazinone, 4.82 kg ai ha⁻¹). N (nitrogen, 56.04 kg N ha⁻¹, applied as ammonium nitrate, 34% N). N+K, nitrogen, 56.04 kg N ha⁻¹ (applied as ammonium nitrate, 34% N) + 56.04 kg K₂O ha⁻¹ (applied as muriate of potash).

^bHarvest periods refer to 2020 and 2021 dates, respectively, as follows: (0, refers to initial observation) April 7 and March 15, (1) May 11 and 21, (2) June 8 and 18, (3) July 12 and 16, (4) August 7 and 27, (5) September 27 and October 1, (6) October 26. Data are pooled across replications and years.

^cLeast square means within each harvest not sharing a common letter differ according to Tukey-Kramer test ($P \leq 0.05$).

^dSEM, standard error of the mean.

Table 2. Effect of treatment on bahiagrass groundcover of bahiagrass-dominant pasture (includes smutgrass, bahiagrass, other weeds) grown at the Alapaha Beef Unit near Alapaha, GA, during 2020 and 2021.

Treatment ^a	Harvest 0 ^b	Harvest 1	Harvest 2	Harvest 3	Harvest 4	Harvest 5	Harvest 6
% Groundcover							
Control	58	65	68	84	77	68 cd ^c	45 b
PRE	62	62	69	88	70	64 d	49 b
POST	60	54	71	77	72	87 abc	73 ab
PRE+POST	62	63	79	78	75	81 abcd	90 a
N	65	58	69	78	60	71 bcd	54 b
PRE, N	65	60	73	83	66	86 abcd	70 ab
POST, N	64	53	65	76	75	88 abc	97 a
PRE+POST, N	69	56	66	75	81	91 ab	92 a
N+K	70	62	75	84	74	74 abcd	54 b
PRE, N+K	63	53	75	75	75	71 bcd	55 b
POST, N+K	73	60	68	73	76	93 ab	91 a
PRE+POST, N+K	65	53	73	79	86	95 a	92 a
SEM ^d	9.78	7.03	6.58	6.22	8.28	5.58	7.44

^aPRE (preemergence; indaziflam, 0.28 kg ai ha⁻¹). POST (postemergence; hexazinone, 4.82 kg ai ha⁻¹). N (nitrogen, 56.04 kg N ha⁻¹, applied as ammonium nitrate, 34% N). N+K, nitrogen, 56.04 kg N ha⁻¹ (applied as ammonium nitrate, 34% N) + 56.04 kg K₂O ha⁻¹ (applied as muriate of potash).

^bHarvest periods refer to 2020 and 2021 dates, respectively, as follows: (0, refers to initial observation) April 7 and March 15, (1) May 11 and 21, (2) June 8 and 18, (3) July 12 and 16, (4) August 7 and 27, (5) September 27 and October 1, (6) October 26. Data are pooled across replications and years.

^cLeast square means within each harvest not sharing a common letter differ according to Tukey-Kramer test ($P \leq 0.05$).

^dSEM, standard error of the mean.

declined for all plots that received postemergence herbicide compared to the unsprayed plots and most plots receiving only preemergence ($P < 0.01$). Unfortunately, the large variation in the plots at the final harvest event resulted in a large standard error of the mean, and this decline was not always significant among certain pairwise comparisons.

Similar trends were observed with respect to bahiagrass groundcover (Table 2). Treatment responses did not indicate any differences from the nontreated control for bahiagrass groundcover in harvest 0 through harvest 4 (April–August; Table 2; $P > 0.05$). Following postemergence application of hexazinone at harvest 4 (August) bahiagrass groundcover generally increased in the respective treatments for the following harvests (September and October; $P < 0.01$). Other weeds' groundcover followed

similar trends to bahiagrass cover from the initial observation at harvest 0 through harvest 4 (April–August; $P > 0.05$). Following the application of hexazinone, the PRE+POST+N, POST+N+K, and PRE+POST+N+K treatments reduced other weeds' groundcover when compared to the nontreated control at harvest 5 ($P < 0.01$). Data could not be analyzed at harvest 6, because other weeds were not found in any of the plots.

Data for dead plant material could not be analyzed for harvests 0 through harvest 4 (April–August), because none were found in any of the plots before postemergence application of hexazinone. However, following hexazinone treatment applications, dead plants were observed for all treatments that included some level of postemergence. Witnessing necrosis and terminated plants following postemergence application was expected

(Coffman et al. 1993). Observations of dead plant material did not negatively affect bahiagrass and were limited to targeted weeds. It should be noted that even though the *P* value indicated treatment differences at harvest 5 (September) and 6 (October), the results from the Tukey-Kramer tests did not find differences in the pairwise comparisons.

When botanical composition is coupled with the results from herbage accumulation (HA), it is evident that an IWM plan including preemergence, postemergence, and fertilizer produces a more favorable result than the current practice (postemergence alone). An IWM plan would help to reduce the introduction of other weedy species and provide essential nutrients for sustaining bahiagrass long-term that could not be accounted for with a singular application of hexazinone. The results of this research support previous literature on the efficacy of hexazinone as a management tool for controlling smutgrass (Hancock et al. 2010; Mislevy et al. 1999; Sellers et al. 2020). However, their research had a singular focus on postemergence alone or postemergence and fertilizer interactions. One of the many challenges that result from removing smutgrass from bahiagrass forage systems is the disturbance to the preexisting canopy. Reduced groundcover leads to increased light penetration and accessibility to space and nutrients for many opportunistic annual and perennial weedy species. This is a well-known ecological principle among species dynamics and plant-plant interactions (Kemp and King 2001). Integrating indaziflam can benefit bahiagrass by limiting weed competition while it grows to fill the voids left by the mature smutgrass plants. Results indicated that a preemergence application of indaziflam before harvest 1 (May) did not negatively affect bahiagrass cover, supporting similar research conducted by Hurdle et al. (2020) in bermudagrass. This is not surprising, as morphological characteristics between bermudagrass and bahiagrass are similar. Continued research is needed to determine the impact of indaziflam on future smutgrass populations in upcoming seasons.

The benefits of fertilizer to bahiagrass have been well established. However, the lack of postemergence activity may permit smutgrass to outcompete bahiagrass, even when fertilizer is applied without subsequent herbicide. Consequently, a singular focus on fertilizer to increase the competitive advantage of bahiagrass over smutgrass and other weeds may not be the best approach. Vengris et al. (1953) suggested that weed species can better utilize plant nutrients and are more aggressive than most desired crop species, regardless of the stage of growth, soil fertility relationship, and seasonal weather conditions. Much of the region is prone to acidic soils ($\text{pH} = 4.5\text{--}5.5$), and climatic challenges coupled with a high water table disallows forage producers from maintaining a neutral pH . It has been well established that bahiagrass grows best at a pH of 5.5. Rana et al. (2013) concluded that a pH range that is either too high (6.5) or too low (4.5) provided a distinct competitive advantage for giant smutgrass over bahiagrass. Furthermore, Beaty et al. (1960) highlighted that the peak performance of bahiagrass is during the hottest months of the year in June and July. Holding to the advantageous characteristics of many other weedy species, smutgrass productivity is not limited by these seasonal patterns and benefits greatly as a result of declining bahiagrass later in the season. Even though bahiagrass is known as a somewhat aggressive warm-season perennial species, forage systems are dynamic, with extreme environmental exposures and antagonistic pests competing for the same resources. As a result, continual shifts in species abundance and distribution is commonplace, making it challenging to manage.

Herbage Accumulation

Year \times treatment did not interact to affect HA, so data were pooled over both years ($P = 0.75$). Data were analyzed within harvest to better isolate the treatment responses ($P < 0.01$). No differences in HA were reported in harvest 1 ($P = 0.15$; Table 3; May). Early-season fertilizer generally resulted in an increase in HA of the fertilized treatments at the second harvest (June); however, the increase was not always significantly different from unfertilized treatments. Again, no differences in HA were found at harvest 3 (July) despite the previous fertilizer treatment ($P < 0.01$). It should be noted that even though the *P* value indicated treatment differences at harvest 3, the results from the Tukey-Kramer tests did not find differences in the pairwise comparisons. The second fertilizer application was made following harvest 3, so it was not surprising that the treatments receiving N had greater HA at harvest 4 (August) than the unfertilized treatments ($P < 0.01$). The postemergence herbicide was applied following harvest 4, and the effects are seen in subsequent harvests. In general, HA declined at harvests 5 and 6 (September–October) in treatments including postemergence herbicide ($P < 0.01$). However, these differences were not always different from the treatments not receiving the postemergence herbicide treatment. This is likely a consequence of biomass differences between bahiagrass, smutgrass, and other weeds. Overall, HA illustrated seasonal trends following low yields at the beginning of the growing season, peaking during the mid-point when daily temperatures were most extreme, and gradually declining into the fall, representing a typical growth pattern for warm-season perennial forages (Beaty et al. 1960). Overall, HA was not a sufficient metric to determine treatment effectiveness in the study.

Forage Nutritive Values

No interactions between year \times treatment for CP and TDN were observed; therefore, data were pooled over years ($P = 0.75$). Data for CP and TDN were also analyzed within harvest. In harvest 1, CP increased from 7.6 to 9.2 mg g^{-1} with the addition of N compared to the nontreated control, supporting experimental evidence by Silveira et al. (2017). No differences in CP concentrations were observed with K and herbicide treatments. Furthermore, the relatively low pH found in bahiagrass forage systems in this region (4.5–5.5) is similar to those found in Florida (5.5), where Yarborough et al. (2017) expressed the impact of this lower pH range to the potential leaching of K. Regardless of treatment combinations, there were no practical differences in TDN throughout the experimental period.

Economics

Costs among the treatments ranged from US\$145.00 to 610.00 ha^{-1} , whereas the estimated cost for complete bahiagrass renovation was US\$1,079.00 ha^{-1} (based on Lacy et al. 2016; Table 4). The most common practice for producers in the southeastern United States is to utilize postemergence alone, keeping costs low at US\$145.00 ha^{-1} . However, work done by Sollenberger (2019) highlighted the negative impact of neglecting proper fertilization that leads to declining bahiagrass forage systems in Florida. Therefore, the addition of N is highly recommended, with an associated cost of US\$415.00 ha^{-1} for POST + N. Yarborough et al. (2017) highlighted the benefit of adding K in addition to N for increasing root mass and the importance of this management practice for promoting persistence in

Table 3. Effect of treatment on forage accumulation (given as dry matter, DM) of bahiagrass-dominant pasture (includes smutgrass, bahiagrass, other weeds) grown at the Alapaha Beef Unit near Alapaha, GA, during 2020 and 2021.

Treatment ^a	Harvest 1 ^b	Harvest 2	Harvest 3 ^d	Harvest 4	Harvest 5	Harvest 6
	kg DM ha ⁻¹					
Control	889	552 bc ^c	729	935 b	970 abc	380 abc
PRE	767	475 c	864	974 b	969 abc	415 ab
POST	889	541 bc	883	830 b	609 d	229 d
PRE+POST	1,048	593 abc	874	984 b	717 cd	274 cd
N	1,058	831 a	693	1,378 a	1,248 a	413 ab
PRE, N	980	685 abc	701	1,442 a	1,291 a	444 a
POST, N	1,130	733 abc	726	1,385 a	988 abc	312 bcd
PRE+POST, N	1,097	705 abc	848	1,312 a	828 bcd	268 d
N+K	1,011	734 abc	849	1,381 a	1,154 ab	422 ab
PRE, N+K	1,168	780 ab	826	1,448 a	1,269 a	455 a
POST, N+K	1,095	829 a	862	1,372 a	877 bcd	254 d
PRE+POST, N+K	1,144	784 ab	826	1,427 a	845 bcd	278 cd
SEM ^e	262.84	62.53	399.85	91.37	613.79	28.10

^aPRE (preemergence; indaziflam, 0.28 kg ai ha⁻¹). POST (postemergence; hexazinone, 4.82 kg ai ha⁻¹). N (nitrogen, 56.04 kg N ha⁻¹, applied as ammonium nitrate, 34% N). N+K, nitrogen, 56.04 kg N ha⁻¹ (applied as ammonium nitrate, 34% N) + 56.04 kg K₂O ha⁻¹ (applied as muriate of potash).

^bHarvest periods refer to 2020 and 2021 dates, respectively, as follows: (0) April 7 and March 15, (1) May 11 and 21, (2) June 8 and 18, (3) July 12 and 16, (4) August 7 and 27, (5) September 27 and October 1, (6) October 26. Data are pooled across replications and years.

^cLeast square means within each harvest not sharing a common letter differ according to Tukey-Kramer test ($P \leq 0.05$).

^dNote: P-value indicated treatment differences at harvest 3, although results from the Tukey-Kramer tests did not find differences in the pairwise comparisons.

^eSEM, standard error of the mean.

Table 4. Market costs for selected integrated management strategies for controlling smutgrass in bahiagrass forage systems compared to a complete bahiagrass renovation following University of Georgia recommendations.

Treatment	US\$	Unit	US\$ ha ⁻¹
Indaziflam	0.34	mL	95.00
Hexazinone	0.03	mL	145.00
Nitrogen (UAN32) ^a	1.02	L	270.00
Potassium (muriate of potash) ^a	0.90	kg	100.00
Selected integrated plans ^b			US\$ ha ⁻¹
POST			145.00
PRE+POST			240.00
POST+N			415.00
PRE+POST+N			510.00
POST+N+K			515.00
PRE+POST+N+K			610.00
Bahiagrass renovation ^c			
2018 hybrid Bermuda hay-non irrigated establishment			1,079.00

^aFertilizer prices were collected from DTN in January 2022.

^bPRE (preemergence; indaziflam, 0.28 kg ai ha⁻¹). POST (postemergence; hexazinone, 4.82 kg ai ha⁻¹). N (nitrogen, 56.04 kg N ha⁻¹, applied as ammonium nitrate, 34% N). N+K, nitrogen, 56.04 kg N ha⁻¹ (applied as ammonium nitrate, 34% N) + 56.04 kg K₂O ha⁻¹ (applied as muriate of potash).

^cBahiagrass renovations were calculated by modifying the University of Georgia Extension, College and Agricultural and Environmental Sciences–Applied Economics, 2018 hybrid bermuda hay-nonirrigated establishment budget. Market costs for integrated management strategies were selected based on smutgrass management in bahiagrass forage systems, with the addition of indaziflam for current research.

extensive grazing systems. Costs for a regime that includes POST+N+K are US\$515.00 ha⁻¹.

The greatest return on investment for controlling smutgrass and other invasive weeds should include a preemergence treatment for controlling annual weed emergence, following a postemergence application for controlling preexisting weeds. Incorporating a full-spectrum fertilizer plan that includes at a minimum N+K for boosting bahiagrass growth over competitive weeds increases its overall longevity. The IWM plan (POST+PRE+N+K) costs were US\$610.00 ha⁻¹. Many producers will find this costly; however, when compared to the costs of complete bahiagrass renovation

at US\$1,079.00 ha⁻¹, there are significant savings that can be utilized elsewhere in the operation—not to mention the loss of production time for reestablishment. Because of the significant expense, Sellers et al. (2020) recommend only treating fields infested with more than 50% smutgrass. Regardless, rising costs and a potentially limiting supply of both fertilizer and herbicides could negatively affect management decisions. Fortunately, several treatment options (e.g., POST+N; US\$415.00 ha⁻¹) (Table 4) are available for reducing smutgrass and limiting weed emergence, giving producers additional options when difficult circumstances arise.

Future Work

Producers should implement an integrated management plan that utilizes a timely application of indaziflam to reduce the introduction and emergence of weed species and hexazinone when preexisting smutgrass infestations are at least 50%. This should be combined with fertilizer applications for enriching bahiagrass productivity that increases both HA and groundcover, providing a competitive advantage in disturbed canopies. This research concluded that treatments that included preemergent (indaziflam) and postemergent (hexazinone) herbicides in addition to N and K₂O resulted in an improved bahiagrass stand, as timely weed suppression removed competition, whereas fertilizer provided essential nutrients for optimum growth to fill in the gaps. Combining herbicide and fertilizer is a more economical option for producers when compared to a complete bahiagrass renovation (US\$610.00 ha⁻¹ vs US\$1079.00 ha⁻¹). Smutgrass was reduced to <15% ground coverage when a postemergent herbicide was applied. The addition of a preemergent herbicide and/or fertilizer further reduced the coverage of smutgrass ($P < 0.01$). It is important to note that bahiagrass ground coverage increased, whereas other and dead material did not differ by treatment.

Future work will continue to look at preventing further smutgrass infestations utilizing this novel strategy. Research is ongoing to improve the effectiveness of herbicide and fertilizer treatments and screen new herbicide technologies for preventing the introduction of other noxious weeds. Although the need for more research

is understood, the scope of this research by expanding novel management strategies will deliver agronomic and economic stability to producers for improving bahiagrass forage utilization and preventing the introduction of other weeds in the southeastern United States.

Acknowledgments. Funding for this research was provided by Georgia Beef Cattle Commodity Commission.

No conflict of interest has been declared.

References

- Beatty ER, McCreery RA, Powell JD (1960) Response of Pensacola bahiagrass to nitrogen fertilization. *J Agron* 52:453–455
- Beatty ER, Smith YC, Powell JD (1974) Response of Pensacola bahiagrass to irrigation and time of N fertilization. *J Range Manag* 27:394–396
- Chambliss CG, Sollenberger LE (1991) Bahiagrass: The foundation of cow–calf nutrition in Florida. Pages 74–80 in *Proceedings of the 40th Annual Beef Cattle Short Course*. Gainesville, FL: IFAS, University of Florida
- Chemical Warehouse (2022). Herbicide costs. <https://chemicalwarehouse.com/>. Accessed: May 1, 2022
- Coffman CB, Frank JR, Potts WE (1993) Crop responses to hexazinone, imazapyr, tebuthiuron, and triclopyr. *Weed Technol* 7:140–145
- Dias JLC, Sellers BA, Ferrell JA, Silveira ML, Vendramini JM (2018) Herbage responses to dogfennel cover and limited nitrogen fertilization in bahiagrass pastures. *J Agron* 110:2507–2512
- Ferrell JA, Mullahey JJ (2006) Effect of mowing and hexazinone application on giant smutgrass (*Sporobolus indicus* var. *pyramidalis*) control. *Weed Technol* 20:90–94
- Ferrell JA, Mullahey JJ, Dusky JA, Roka FM (2006) Competition of giant smutgrass (*Sporobolus indicus*) in a bahiagrass pasture. *Weed Sci* 54:100–105
- Georgia Weather Network (2022) University of Georgia, College of Agricultural and Environmental Sciences, Athens, GA. <http://www.georgiaweather.net/>. Accessed: May 1, 2022
- Hancock DW, Lacy RC, Stewart L, Tubbs RS, Kichler JM, Green TW, Hicks R (2010) The management and use of bahiagrass. University of Georgia. Bulletin 1362. extension.uga.edu/publications. Accessed: May 1, 2022
- Hurdle NL, Grey TL, McCullough PE, Shilling D, Belcher J (2020) Bermudagrass tolerance of indaziflam PRE applications in forage production. *Weed Technol* 34:125–128
- Kemp DR, King WM (2001) Plant competition in pastures—implications for management. Pages 85–102 in Tow PG, Lazenby A, eds, *Competition and Succession in Pastures*. Wallingford, UK: CABI Publishing
- Kissel DE, Sonon LS (2008) Soil test handbook for Georgia (Bulletin 62). Athens, GA: University of Georgia Extension
- Lacy RC, Morgan RN, Russell LA (2016) Tifton 85 Bermuda Grass Hay Budget. University of Georgia: Extension Ag Economics and Crop & Soil Science Departments. <https://agecon.uga.edu/extension/budgets.html>. Accessed: May 1, 2022
- Littell R, Milliken G, Stroup W, Wolfinger R, Schabenberger O (2006) SAS for Mixed Models, 2nd Edn. Cary, NC: SAS Institute Inc.
- McCaleb JE, Hodges EM, Kirk WG (1963) Smutgrass control. University of Florida IFAS Extension, Ona, FL. Circular S-149. 10 p.
- McIntosh D, Anderson-Husmoen BJ, Kern-Lunbery R, Goldblatt R, Lemus P, Griggs T, Bauman L, Boone S, Shewmaker G, Teutsch C (2022) Guidelines for Optimal Use of NIRSC Forage and Feed Calibrations in Membership Laboratories, 2nd Edn. Knoxville, TN: The University of Tennessee Press
- Mislevy P, Shilling DG, Martin FG, Hatch SL (1999) Smutgrass (*Sporobolus indicus*) control in bahiagrass (*Paspalum notatum*) pastures. *Weed Technol* 13:571–575
- Mislevy P, Martin FG, Hall DW (2002) West Indian dropseed/giant smutgrass (*Sporobolus indicus* var. *pyramidalis*) control in bahiagrass (*Paspalum notatum*) pastures. *Weed Technol* 16:707–711
- Moore JE, Undersander DJ (2002) Relative forage quality: an alternative to relative feed value and quality index. Pages 16–28 in *Proceedings of the 13th Annual Florida Ruminant Nutrition Symposium*, January 2002, Gainesville, FL.
- Murray I, Cowe I (2004) Sample preparation. *Near-infrared Spectroscopy in Agriculture* 44:75–112
- NIRSC (2020) NIRSC Forage and Feed Testing Consortium. <https://www.nirconsortium.com/>. Accessed: May 1, 2022.
- Nolte S (2017) Smutgrass management in bermudagrass and bahiagrass pastures. Texas A&M AgriLife Extension Service, College Station, TX: Bulletin SCS-2017-18. <http://publications.tamu.edu>. Accessed: May 1, 2022
- Quinn R (2022) DTN retail fertilizer trends. <https://www.dtnpf.com/agriculture/web/ag/crops/article/2022/03/02/average-retail-fertilizer-prices>. Accessed: May 1, 2022
- Rana N, Sellers B, Ferrell J, MacDonald G, Silveira M, Vendramini J (2013) Impact of soil pH on bahiagrass competition with giant smutgrass (*Sporobolus indicus* var. *pyramidalis*) and small smutgrass (*Sporobolus indicus* var. *indicus*). *Weed Sci* 61:109–116
- Rana N, Wilder B, Sellers B, Ferrell J, MacDonald G (2012) Effects of environmental factors on seed germination and emergence of smutgrass (*Sporobolus indicus*) varieties. *Weed Sci* 60:558–563
- Sellers BA, Ferrell JA (2011) The impact of mowing smutgrass control. The Florida Cattlemen and Livestock Journal. <https://rcrec-ona.ifas.ufl.edu/media/rcrec-onaifasufledu/pdf/or6-2012.pdf>. Accessed: October 24, 2022
- Sellers BA, Rana N, Dias JLC, Devkota P (2020) Smutgrass control in perennial grass pastures. University of Florida IFAS Extension: Bulletin SS-AGR-18/AA261, Ona, FL.
- Silveira ML, Vendramini JM, Sellers BA (2017) Nitrogen fertilizer source effects on bahiagrass responses. *Crops Soils* 50:4–9
- Sollenberger LE (2019) Finding a solution to declining bahiagrass pastures. Hay & Forage Grower. <https://hayandforage.com/article-2364-finding-a-solution-to-declining-bahiagrass-pastures.html>. Accessed: May 1, 2022
- Vengris J, Drake M, Colby W, Bart J (1953) Chemical composition of weeds and accompanying crop plants. *J Agron* 45:213–218
- Wallau M, Vendramini J, Dubeux J, Blount A (2010) Bahiagrass (*Paspalum notatum* Fluegge): overview and pasture management. University of Florida, IFAS Extension: Bulletin SS-AGR-332, Ona, FL. <https://edis.ifas.ufl.edu/ag342>. Accessed: May 1, 2022
- Web Soil Survey (2022) Natural Resources Conservation Service, United States Department of Agriculture. <http://websoilsurvey.sc.egov.usda.gov/>. Accessed: May 1, 2022
- Wilder B, Ferrell JA, Sellers BA, MacDonald GE (2008) Influence of hexazinone on ‘Tifton 85’ bermudagrass growth and forage quality. *Weed Technol* 22:499–501
- Yarborough JK, Vendramini JM, Silveira ML, Sollenberger LE, Leon, RG, Sanchez JM, Filho CS (2017) Impact of potassium and nitrogen fertilization on bahiagrass herbage accumulation and nutrient concentration. *J Agron* 109:1099–1105