



Sensitivity to Salinity at the Emergence and Seedling Stages of Barnyardgrass (*Echinochloa crus-galli*), Weedy Rice (*Oryza sativa*), and Rice with Different Tolerances to ALS-Inhibiting Herbicides

Authors: Fogliatto, Silvia, Patrucco, Lorenzo, Milan, Marco, and Vidotto, Francesco

Source: Weed Science, 69(1) : 39-51

Published By: Weed Science Society of America

URL: <https://doi.org/10.1017/wsc.2020.72>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Research Article

Cite this article: Fogliatto S, Patrucco L, Milan M, Vidotto F (2021) Sensitivity to salinity at the emergence and seedling stages of barnyardgrass (*Echinochloa crus-galli*), weedy rice (*Oryza sativa*), and rice with different tolerances to ALS-inhibiting herbicides. *Weed Sci.* **69**: 39–51. doi: [10.1017/wsc.2020.72](https://doi.org/10.1017/wsc.2020.72)

Received: 19 June 2020

Revised: 15 September 2020

Accepted: 16 September 2020

First published online: 12 October 2020

Associate Editor:

Franck E. Dayan, Colorado State University

Keywords:

Climate change; herbicide resistance; weed growth

Author for correspondence:

Silvia Fogliatto, Dipartimento di Scienze Agrarie, Forestali e Alimentari, Università degli Studi di Torino, Largo P. Braccini 2, 10095 Grugliasco, Torino, Italy.
Email: silvia.fogliatto@unito.it

Sensitivity to salinity at the emergence and seedling stages of barnyardgrass (*Echinochloa crus-galli*), weedy rice (*Oryza sativa*), and rice with different tolerances to ALS-inhibiting herbicides

Silvia Fogliatto¹ , Lorenzo Patrucco², Marco Milan¹  and Francesco Vidotto³ 

¹Postdoc Fellow, Dipartimento di Scienze Agrarie, Forestali e Alimentari, Università degli Studi di Torino, Grugliasco, Torino, Italy; ²Postgraduate Fellow, Dipartimento di Scienze Agrarie, Forestali e Alimentari, Università degli Studi di Torino, Grugliasco, Torino, Italy and ³Associate Professor, Dipartimento di Scienze Agrarie, Forestali e Alimentari, Università degli Studi di Torino, Grugliasco, Torino, Italy

Abstract

Salinization is affecting many rice (*Oryza sativa* L.) areas worldwide and weed infestation, together with the occurrence of herbicide-resistant populations, is further limiting rice yield. This study aimed at evaluating the effect of water salinity on the emergence and seedling growth of five Italian barnyardgrass [*Echinochloa crus-galli* (L.) P. Beauv.] populations (three sensitive and two resistant to acetolactate synthase-inhibiting herbicides), three Italian weedy rice (*Oryza sativa* L.) populations (all sensitive to imazamox), and two rice varieties (the conventional ‘Baldo’ variety and the imazamox-tolerant ‘CL80’ one). In 2017, seeds were sown in alveolar trays filled with sand, a nutrient solution, and water with the following salt concentrations: 0, 50, 100, 150, 200, and 250 mM NaCl. Plant emergence (after 15 d), plant height, shoot and root weight, chlorophyll a, chlorophyll b, and carotenoid content were measured at 40 d after sowing. *Echinochloa crus-galli* showed a higher tolerance to salinity than *O. sativa* and rice. All species were affected more at the seedling stage than at emergence. A variable behavior of the herbicide-resistant populations was shown; one resistant *E. crus-galli* population was affected more by salinity and showed a lower emergence rate (about 20% against 40% emergence of the other populations at the highest salt concentration) and reduced seedling growth, while the other resistant population’s response was similar to that of the sensitive populations. The chlorophyll content increased as the salt content increased in all *E. crus-galli* populations. The highest emergence and growth reduction in *O. sativa* were recorded in the imazamox-tolerant rice. Rice and *O. sativa* were able to grow only up to 50 mM. *Echinochloa crus-galli* populations are probably favored under saline conditions, while lower infestation by *O. sativa* can be expected.

Introduction

Environmental stresses, such as high and low temperatures, drought, salinity, alkalinity, and high CO₂ concentrations, are able to influence the growth of plants (Ahuja et al. 2010).

Salinity represents one of the main environmental stresses that can affect plant growth and cause crop yield reductions (Negrão et al. 2017; Parihar et al. 2015). All the primary metabolic processes (photosynthesis, protein synthesis, and lipid metabolism) of a plant are involved in the set of responses to salt stress. Currently, the effects of salinity on plant growth are gaining importance because salinization areas are increasing throughout the world, mainly as a consequence of the overexploitation of water resources and climate change (Akbari et al. 2020; Kibria et al. 2017; Singh 2018). It has been estimated that about 20% of irrigated lands have suffered from an increase in this phenomenon over the last few decades (Machado and Serralheiro 2017). The presence of salt in these areas can be an obstacle to the growth of various crops, such as rice (*Oryza sativa* L.), because of its toxicity and because of osmotic stress that can alter the water uptake potential/dynamics of plants (Kibria et al. 2017).

Rice is known to be sensitive to salt stress and, even though the plant is able to germinate in moderately saline conditions, it is more susceptible during the seedling emergence and early growth stages (Bertazzini et al. 2018). Previous studies found that seedling growth was reduced at a salinity level of 1.9 dS m⁻¹ for some rice varieties and that values of 3 dS m⁻¹ can reduce rice survival (Bertazzini et al. 2018; Zeng and Shannon 2000). Rice is usually better at tolerating saline conditions in the subsequent vegetative stages, but it still becomes sensitive during the reproductive stage, when high-saline conditions can hinder grain formation (Bertazzini et al. 2018; Hoang et al. 2016). Moderate salinity levels in water and soil are thus able to cause serious

© The Author(s), 2020. Published by Cambridge University Press on behalf of Weed Science Society of America. This is an Open Access article, distributed under the terms of the Creative Commons Attribution licence (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted re-use, distribution, and reproduction in any medium, provided the original work is properly cited.



reduction in yield, while very high salinity ($>11.5 \text{ dS m}^{-1}$) leads to total failure of a crop (Zeng and Shannon 2000).

The effects of salinity on cultivated plants can be intensified by the presence of other limiting factors, such as competition due to the presence of weeds, which are in general less susceptible to environmental stresses than crops, as they are more adaptable to changes (Fogliatto et al. 2020; Korres et al. 2016). The higher tolerance of weeds to environmental changes is mainly related to the fact that wild species have not undergone the same level of selection to obtain advantageous traits (Korres et al. 2016). A previous review (Cirillo et al. 2018) highlighted that the majority of the weeds considered in the study showed a higher tolerance to salt, in terms of germination and plant growth, than most of the crops; in fact, it has been found that, for the majority of weeds throughout the world, the level of salinity that can cause a growth reduction of 50% (EC_{50}) is higher than that of crops, with an average EC_{50} of 16.4 and 10.6 dS m^{-1} for weeds and crops, respectively (Cirillo et al. 2018; Tanji and Kielen 2002).

Barnyardgrass [*Echinochloa crus-galli* (L.) P. Beauv.] and weedy rice (*Oryza sativa* L.) are among the principal weeds in rice fields throughout the world (Andres et al. 2015; Vidotto et al. 2020). According to some studies, *E. crus-galli* is considered to be able to cope with high-saline conditions, probably because this species is able to take up potassium more selectively than sodium (Wilson and Read 2006). The salt tolerance of *E. crus-galli* is generally higher than that of many other crops, such as rice, and other weeds, like *O. sativa* (Fogliatto et al. 2019; Hakim et al. 2011; Serra et al. 2018). *Echinochloa* is also one of the genera most prone to developing herbicide resistance; it has been demonstrated that different populations of *E. crus-galli* have become resistant to eight site-of-action herbicides used in rice throughout the world and, in particular, to acetolactate synthase (ALS) and acetyl-CoA carboxylase (ACCase) inhibitors (Heap 2020). *Oryza sativa* populations have also developed resistance to ALS inhibitors, even though this has occurred more recently, after the introduction of Clearfield® rice varieties tolerant to imidazolinone herbicides (Goulart et al. 2014; Sudianto et al. 2013). *Oryza sativa* resistance to imidazolinone herbicides is caused by one or different point mutations in the ALS gene that result in amino acid substitution; the introduction of Clearfield® varieties harboring these mutations has increased the risk of resistant gene flow from these varieties to *O. sativa*; gene flow has been found to be the predominant source of resistance in *O. sativa*, while independent mutation is a minor source (Fogliatto et al. 2020; Goulart et al. 2014).

Many studies have been conducted to determine the level of rice's tolerance to salt, although only a few studies have been carried out to establish the effect of salinity on weed growth (Chauhan et al. 2013; Fogliatto et al. 2019; Hakim et al. 2011; Serra et al. 2018). Moreover, the growth behavior of herbicide-resistant and herbicide-sensitive weeds under saline conditions has been investigated even less (Fogliatto et al. 2019; Serra et al. 2018). Herbicide resistance can induce in some resistant populations a fitness cost, as some resources in the resistant plants are diverted to protect the plants from the effects of the herbicides instead of being used for growth, reproduction, and countering abiotic stresses (Goulart et al. 2012; Vila-Aiub et al. 2015). Resistant individuals can exhibit different behavior than susceptible individuals in different environments as a consequence of the fitness penalty (reduced germination, growth, seed production, etc.), and it has been established that fitness costs are usually more evident under adverse environmental conditions (i.e., salinity) (Wu et al. 2016).

The present study has been conducted with the aim of evaluating the effect of salinity on the emergence and plant growth of some Italian populations of *O. sativa* and *E. crus-galli*.

Limited information is available on weed emergence under saline conditions, as many studies only considered the effect of salt on germination in tests that were often conducted in petri dishes, while in studies focused on weed growth, salt was usually added a few days after emergence in order to have a higher number of emerged plants. As one of the aims of this study was to evaluate the effects of salt on emergence, weeds and rice species were left to germinate and emerge from sand in saline solutions.

This study was also aimed at establishing whether two populations of *E. crus-galli* known to be resistant to ALS-inhibiting herbicides would respond differently to saline conditions compared with three herbicide-sensitive populations. The study also compared the growth behavior in response to salinity of the conventional rice variety 'Baldo' and the imazamox-tolerant rice variety 'CL80'.

Materials and Methods

Weed Populations, Rice Varieties, and Salt Concentrations

The study was conducted on five *E. crus-galli* populations, three *O. sativa* populations, and two rice varieties. Seeds from *E. crus-galli* populations were collected in August 2016 from Italian rice fields located in different zones of the main area of rice cultivation in Italy (between Piedmont and Lombardy) that have similar climatic conditions; the fields had a history of repeated applications of the ALS-inhibiting herbicide penoxsulam. Three *E. crus-galli* populations (named s1, s2, and s3) were found to be sensitive to penoxsulam in previous greenhouse and laboratory tests (unpublished data), while two (named r1 and r2) were found to be resistant as a result of a target-site mutation (Trp-574-Leu). The same populations showed the following germination levels without salt in petri dishes: 90%, 95%, 88%, and 85%, respectively for populations s1, s2, s3, and r. Three *O. sativa* populations (wr1, wr2, and wr3) were also collected in 2016 from different Italian paddy fields, also located in the main Italian rice area, and proved sensitive to the main ALS-inhibiting herbicide imazamox used in rice cultivation (data not shown). The same *O. sativa* populations, in a study carried out in petri dishes, showed the following germination levels without salt: 93%, 92%, and 95%, respectively for wr1, wr2, and wr3.

The present study was also conducted on the Baldo rice variety, one of the most frequently cultivated varieties in the Mediterranean Basin, which is characterized by high commercial importance, and on the imidazolinone (imazamox)-tolerant rice variety CL80 as a representative of the Clearfield® varieties cultivated in Italy. Germination level without salt assessed in petri dishes was 97% in Baldo and 89% in CL80.

The following salt (NaCl) concentrations were considered: 0 (control), 50, 100, 150, 200, and 250 mM, corresponding to an Electrical Conductivity (EC) of about 0.5, 5.9, 10.6, 15.9, 21.3, and 26.6 dS m^{-1} , a range that comprises some of the possible concentrations that can be found in European rice-cultivation areas (Badia 2000; Gay et al. 2010; Isla et al. 2003).

The following salt (NaCl) concentrations were considered: 0 (control), 50, 100, 150, 200, and 250 mM, corresponding to an Electrical Conductivity (EC) of about 0.5, 5.9, 10.6, 15.9, 21.3, and 26.6 dS m^{-1} , a range that comprises some of the possible concentrations that can be found in European rice-cultivation areas (Badia 2000; Gay et al. 2010; Isla et al. 2003).

Whole-Plant Bioassay

The trials were carried out during the spring of 2017 (March to June) in a greenhouse at the Department of Agricultural, Forest and Food Sciences of the University of Turin (Italy).

The weed populations and the rice varieties were sown in alveolar trays with 20 holes (2.7 by 2.7 by 5.8 cm each) arranged in five columns by four rows. The trays were kept in the greenhouse at an average daily temperature of 25 C and a photoperiod of 12 h. The trays were filled with fine river sand, and three seeds were sown per hole.

One tray hosted the *E. crus-galli* populations for each salt concentration and another tray hosted the *O. sativa* populations and the rice varieties. Each column of the alveolar tray hosted either one weed population or a rice variety. The column occupied by a weed population or a rice variety was attributed randomly to each tray. Each alveolar tray was considered as a replicate, and three replications were conducted for each salt concentration. Plants were kept in the same trays for the duration of the test.

After seeding, the trays were placed into rectangular plastic containers, each filled with 2 L of solution characterized by different saline concentrations. The solutions were prepared using tap water and different proportions of NaCl to obtain the desired salt concentrations. The following NaCl quantities per liter of water were added: 2.9, 5.8, 8.8, 11.7, and 14.6 g to obtain 50, 100, 150, 200, and 250 mM NaCl concentrations, respectively. The nutrient solution, Hoagland's No. 2 Basal Salt Mixture (powder for tissue culture, product no. H2395, Sigma-Aldrich, 3050 Spruce Street, St Louis, MO 63103, USA), was added to the 2 L of solution in each tray at a dose of 3.2 g. An aliquot of 250 ml of tap water was added to the solution in the trays three times a week, starting from 7 d after the beginning of the test, to compensate for losses in the solution due to evaporation and/or adsorption and to restore the correct EC values. The pH was measured every day for the duration of the study. At the beginning of the test, the pH was equal to 6.6, while the value increased and stabilized at 7.75 during the trial. The entire study was repeated twice, with the second study conducted about 20 d after the first one ended. The average daily temperature of the greenhouse for both studies was about 25 C with a 12-h daylength.

Seedling Development

Plant Emergence

The number of *E. crus-galli*, *O. sativa*, and rice plants that emerged was recorded for each salt concentration at 15 d after sowing. No other seedlings emerged from the substrate after this date. The number of plants that had emerged on the 15th day was then expressed as percentage of emergence.

Plant Height, Shoot, and Root Weight

Plant height was measured from the ground to the apex of the longest leaf at about 40 d after sowing, when plants had reached the 5-leaf stage. The measurements for each population/cultivar, treatment, and replicate were carried out on four plants randomly selected from each column in the alveolar tray. The shoot and root fresh weight of all the plants from each weed population and rice variety were then measured at the same growth stage (5 leaves) by removing the plants from the trays and cutting the shoots from the roots. Sand residues were carefully removed before the roots were weighed.

Chlorophyll and Carotenoid Extraction

On the same day as the plant growth assessment, chlorophyll *a*, chlorophyll *b*, and carotenoid were extracted from each weed population and rice variety for each salt concentration. The extraction was conducted on 0.2-g samples of leaves collected from all the

replicates of each population/variety, to which 10 ml of 80% acetone had been added, and the samples thus obtained were finely ground for a minute using a homogenizer. The samples were then stored for 24 h in darkness at 4 C (Guo et al. 2017). Spectrophotometric measurements of chlorophyll *a*, chlorophyll *b*, and carotenoid content were performed on supernatant samples obtained after centrifugation for 10 min at $2,000 \times g$. Wavelengths of 662, 645, and 470 nm were used for chlorophyll *a*, chlorophyll *b*, and carotenoids, respectively. The amount of these three photosynthetic pigments was expressed as milligrams per gram of seedling fresh weight and determined using the Lichtenthaler and Wellburn formulas (Zhan et al. 2009).

Statistical Analyses

The analyses of the *E. crus-galli* populations were conducted separately, while *O. sativa* and rice were instead analyzed together, as they belong to the same species.

Plant Emergence

The emergence data were first analyzed to check whether they could be represented by a single model (the three-parameter log-logistic regression model in Equation 1) after pooling the data of the repeated experiments. The data were first fit as a pooled data set and then as two separate data sets. ANOVA was conducted to verify the presence of significant differences between the two analyses ($P \leq 0.05$). Given that no differences were found, the emergence data of the two subsets were analyzed as a single data set.

The percentage of emerged plants at 15 d after sowing was fit against the salt concentration, according to a three-parameter log-logistic regression model (Equation 1):

$$Y = \frac{d}{1 + \exp\{b[\log(x) - \log(e)]\}} \quad [1]$$

where *Y* is the plant emergence, *x* is the NaCl concentration expressed in mM, *d* is the upper limit, and *b* is the relative slope at the point of inflection, *e*. Model fitting was performed using the *drm* function of the DRC add-on package of the open-source program R (R Core Team 2019; Ritz et al. 2015).

The salt concentrations required to reduce emergence by 50% (*EC*₅₀) compared with the values observed at 0 mM NaCl were calculated from the fitted model using the *ED* function of the DRC package.

These *EC*₅₀ values were used to carry out pairwise comparisons between the populations and varieties through calculation of the sensitivity index (SI) (Ritz et al. 2006) (Equation 2):

$$SI = \frac{EC_{50}(A)}{EC_{50}(B)} \quad [2]$$

where *A* and *B* indicate the comparison of two different populations/varieties. The significance of SI ($P \leq 0.05$) of each comparison was obtained using the *EDcomp* function of the DRC package. The function *EDcomp* compared the relative potencies for arbitrary doses (in this case the *EC*₅₀) between fitted dose-response curves (Ritz and Streibig 2016). Moreover, to check similarity between curves, a pairwise comparison ($P \leq 0.05$) between curve parameters was carried out using the function *CompParm* of the DRC package. This function tests the null hypothesis that the ratio between parameters is 1, meaning that the two parameters are not significantly different.

Seedling Development and Chlorophyll and Carotenoid Contents

An ANOVA series was conducted separately for each growth parameter to determine the significance of the effect of population, salt concentration, and their interaction on each considered growth parameter and on chlorophyll and carotenoid contents. The interaction was significant for the majority of the considered parameters for both *Echinochloa* spp. and *O. sativa*, the only exception was plant height for both species and shoot weight for *O. sativa* ($P \leq 0.05$). Because of the significance interactions, a further ANOVA series was conducted separately for each weed population and rice variety to evaluate any differences in growth and in the chlorophyll, chlorophyll *a/b* ratio, and carotenoid contents for the different NaCl concentrations. No differences were found between the two experiments, and the data were pooled ($P \leq 0.05$).

Each measured parameter of each population/variety was also expressed as a percentage difference relative to the control (0 mM concentration). A further ANOVA series was then carried out on the percentage differences of each measured parameter for each NaCl concentration to assess the differences among populations/varieties ($P \leq 0.05$). The Levene test was performed to check the homogeneity of variances, and the percentage data were arcsin transformed whenever the ANOVA assumptions were violated. ANOVA was performed using the R statistical program; the means were then separated through the Tukey honest significant difference (HSD) post hoc test included in the *agricolae* package of R.

Results and Discussion

Echinochloa crus-galli

Seedling Development

Plant Emergence. The regression analysis carried out on plant emergence for different salt concentrations established that emergence decreased in all *E. crus-galli* populations for increasing salt levels, albeit at different magnitudes (Figure 1). The percentages of emerged seedlings were already different at 0 mM, with population s2 showing the highest emergence rate, that is, about 86%, and population r2 the lowest, with an average emergence value of about 27%. The emergence rates of populations s1, s3, and r1 in the control (0 mM) were 50%, 55.5%, and 55.6%, respectively. The different initial rate of emergence between populations was also visible when parameter *d* (upper asymptote) was compared across the curves (Table 1). In fact, parameter *d* for population s2, which had the highest emergence rate, was significantly different compared with all the other populations; moreover, populations r1 and r2 also differed in their initial emergences. The *CompParm* function also highlighted that parameter *b*, which is the curve slope, was different between all the resistant populations and populations s2 and s3. Parameter *e*, the point of inflection of the curve, was instead similar for all curves (Table 1). The most similar curves were those of populations s1 and s3 and those of populations s1 and r1, as they showed nonsignificantly different curve parameters and SI values, which is also a comparison between emergence in a specific point of the curve (Table 1).

All populations showed an increase in emergence, compared with the control, for 50 mM, probably due to a hormetic effect of the salt; this increase was moderate in some populations, such as in s2, which only showed a 3% increase (88.8% emergence vs. 86.1% at 0 mM), while it was more pronounced in others, such as in population r2, which displayed a 70% increase in emergence (47.2% emergence vs. 27.7% at 0 mM). However, at 100 mM, only




Figure 1. Dose-response emergence curve with the average data points of the different *Echinochloa crus-galli* populations against the salt concentration. Curve parameter estimates (Equation 1): s1 ($b = 1.94$, $d = 55.85$, $e = 287.76$), s2 ($b = 1.52$, $d = 89.91$, $e = 222.71$), s3 ($b = 1.13$, $d = 89.91$, $e = 282.58$), r1 ($b = 3.96$, $d = 67.67$, $e = 196.55$), r2 ($b = 6.85$, $d = 94.64$, $e = 123.58$). The salt concentration required to reduce emergence by 50% (EC₅₀) is shown below the graph. Only the significant pairwise comparisons between EC₅₀ (SI index) are shown (Equation 2).

population r1 showed an increase in emergence, that is, of about 20% compared with the control (66.7% emergence vs. 55.5% at 0 mM), while all others showed a reduction in emergence. The decrease in emergence was even more notable for higher salt rates; in fact, about 30% emergence reduction or higher was observed in populations s1 and s3 compared with the control, while population r2 showed the greatest reduction (80%).

The NaCl concentration required to reduce emergence by 50% ranged from 123.58 mM NaCl (population r2) to 287.76 mM (population s1). It was observed, from a pairwise comparison of the SIs, that population r2 was affected significantly more than the sensitive populations (s1, s2, and s3) (Table 1).

Echinochloa crus-galli is one of the most common and problematic weeds for rice production throughout the world, and previous studies demonstrated that the species is quite tolerant to salinity (Chauhan et al. 2013; Cirillo et al. 2018; Hakim et al. 2011; Serra et al. 2018). Although most plants can cope with salinity during germination, they are generally more sensitive at the emergence and early growth stages (Läuchli and Grattan 2007). A previous study conducted on Italian ALS inhibitor-resistant and ALS inhibitor-sensitive *E. crus-galli* populations found that germination occurred up to 400 mM NaCl, although with lower values, that is, about 13% (Serra et al. 2018). Moreover, in the same study, values of up to 90% were recorded for the germination of sensitive populations for up to 250 mM NaCl, while an average germination of 70% was observed for resistant populations (Serra et al. 2018). Previous studies also found a high tolerance of the *Echinochloa* genus during germination for two populations of *Echinochloa glabrescens* Munro ex Hook f. collected in the Philippines, which

Table 1. Curve parameter (Equation 1) and sensitivity index (SI) (Equation 2) comparison between *Echinochloa* spp. populations (estimate and P-values).

Population comparison	Parameter <i>b</i> comparison		Parameter <i>d</i> comparison		Parameter <i>e</i> comparison		SI	
	Estimate	P-value	Estimate	P-value	Estimate	P-value	Estimate	P-value
s1/s2	1.28	0.772	0.62	0.000*	1.29	0.581	1.29	0.581
s1/s3	1.72	0.649	0.96	0.843	1.01	0.977	1.02	0.977
s1/r1	0.49	0.184	0.82	0.165	1.46	0.387	0.68	0.207
s1/r2	0.54	0.350	1.49	0.130	2.31	0.177	0.43	0.002*
s2/s3	1.34	0.726	1.55	0.034*	0.79	0.641	0.79	0.641
s2/r1	0.38	0.003*	1.33	0.049*	1.13	0.623	0.88	0.577
s2/r2	0.42	0.050*	2.40	0.004*	1.80	0.162	0.56	0.013*
s3/r1	0.28	0.001*	0.85	0.310	1.43	0.577	0.69	0.424
s3/r2	0.31	0.014*	1.55	0.121	2.28	0.337	0.44	0.031*
r1/r2	1.10	0.898	1.81	0.027*	1.59	0.151	1.59	0.151

*Significant comparisons at $P \leq 0.05$.

showed a germination of more than 60% for 200 mM NaCl, with an EC_{50} value of between 250 and more than 300 mM (Opeña et al. 2014). These values were similar to those found by Serra et al. (2018) and had the same magnitude as the values found in this study for the emergence of sensitive populations.

In this study, values of between 36% and 44% were observed for the emergence of sensitive populations for the highest salt level, while percentages lower than 20% were recorded for the resistant populations. Thus, emergence seems to be affected more by salinity than germination, as observed in the germination test conducted previously on the same populations; in addition, the emergence of herbicide-resistant populations was reduced to a great extent by high salt concentrations compared with the sensitive ones, as already observed for germination in a previous study (Serra et al. 2018).

Plant Height, Shoot, and Root Weight. *Echinochloa crus-galli* seedling development was hampered by the salinity conditions, as seedling emergence occurred for NaCl concentrations above 150 mM, but this was followed by the total desiccation of the plants. However, population r2 was already affected at 150 mM NaCl, as the seedlings were able to emerge at this concentration but were unable to grow.

Variability in plant heights among the populations was already observed in the control, in which the height ranged between about 20 cm (population s1) and almost 43 cm (population r1) (Table 2). Increasing salt concentrations did not significantly change the plant height in population s1 or r2, while a general decrease in height was observed in the other populations. Increase in plant height was observed at 50 mM, compared with 0 mM, in all the sensitive populations, with increased values that ranged between about 5% (populations s2 and s3) and about 49% (population s1). The resistant population r2 showed a 38% plant height decrease at 50 mM, and thus an absence of hormesis can be hypothesized; plant heights in population r1 were only slightly reduced (decreased by about 10%). Only population s1 still showed an increase in plant height at 100 mM compared with the control (about 38%), while all the other populations behaved similarly to one another, that is, their height was reduced. Apart from r2, all the other populations showed a more marked height reduction for the highest salt concentration (150 mM), which was more moderate in populations s1 and s2.

Shoot weight was affected more than plant height by increasing salt concentrations, as weight reductions occurred (Table 2). Population s1 showed a higher tolerance to salt, and it was the only one to show an increase in shoot weight at 50 mM (63% increase compared with the control); population r1 did not show any

reduction in weight for this salt concentration. All populations showed similar weight reduction at 100 mM, that is, above 70% compared with the control, apart from population s1, which only showed a reduction of about 14%. Population r1 showed an intermediate behavior, with a weight reduction of about 46%. Apart from population r2, which failed to grow, no significant differences in weight reduction were observed among the populations for 150 mM, with reduction values between 62% in population s1 and 88% in population s3, compared with the control.

Root weight also reflected the higher salt tolerance observed for population s1 and the lower salt tolerance observed for population r2. A probable hormetic effect was observed for 50 mM in population s1, whose root weight increased by about 54% compared with 0 mM, while the other populations showed a moderate (s2, r1, and r2) or a more marked (s2) root weight decrease. All populations showed a root weight reduction at 100 mM, which was moderate in populations s1 (about 36%) and r1 (about 46%), but more evident in all the others, reaching even higher reduction values than 80% (populations s3 and r2). As already observed for the shoot weight, the root weight decrease was similar among all populations for 150 mM NaCl.

In general, seedling growth (plant height, shoot and root weight) was affected more by salinity at 150 mM in the ALS inhibitor-resistant population r2, which was not able to mature, compared with the sensitive populations, as already observed for emergence; in fact, even though no significant differences were found between resistant and sensitive populations at 50 mM and 100 mM, growth was completely hampered in r2 at the high salt concentration. Among the sensitive populations, s1 was the most tolerant to salt, showing the smallest reduction for all parameters. However, these differences were only detectable up to 100 mM NaCl.

Reductions in plant height and in the biomass of *E. crus-galli* for increasing salinity levels were observed in previous studies (Chauhan et al. 2013; Hakim et al. 2011; Serra et al. 2018). A study carried out on *E. crus-galli* seeds, collected in environments with different degrees of salinity, showed that the seeds were able to germinate up to 1.5% NaCl, while once they emerged, they only survived about 20 d at a concentration of about 1% NaCl (Rahman and Ungar 1990). A previous study of ours confirmed these results and highlighted that *E. crus-galli* demonstrates a lower tolerance at the early seedling stage than at the emergence and germination stages (Serra et al. 2018). Plant weight was affected more than height, as a generally more marked reduction was observed in the case of shoot and root weight for increasing salt levels; this was already observed in the same species in a previous study in which plant height was found to be more tolerant to salt than

Table 2. Plant height and shoot and root weight of the different *Echinochloa crus-galli* populations and the percentage difference of these growth parameters for each salt concentration compared with the control.

Population	NaCl concentration	Plant height ^a	Plant height difference compared with the control ^b	Shoot weight ^a	Shoot weight difference compared with the control ^b	Root weight ^a	Root weight difference compared with the control ^b
	mM	cm	%	g	%	g	%
s1	0	20.80	—	2.73	—	2.62 ab	—
	50	31.04	49.23 <i>c</i>	4.46	63.29 <i>b</i>	4.04 <i>b</i>	54.20 <i>b</i>
	100	28.85	38.70 b	2.36	−13.65 b	1.69 <i>a</i>	−35.69 b
	150	19.25	−7.45 <i>B</i>	1.02	−62.50	0.68 <i>a</i>	−74.04
s2	0	37.89 ab	—	15.79 <i>c</i>	—	15.35 <i>b</i>	—
	50	40.31 <i>b</i>	6.38 <i>b</i>	10.14 <i>bc</i>	−35.79 <i>a</i>	10.57 <i>ab</i>	−31.15 <i>ab</i>
	100	30.28 <i>ab</i>	−20.09 a	4.16 <i>ab</i>	−73.64 a	4.29 <i>a</i>	−72.05 a
	150	27.85 <i>a</i>	−26.50 <i>B</i>	2.73 <i>a</i>	−82.71	5.29 <i>a</i>	−65.53
s3	0	41.81 <i>b</i>	—	11.84 <i>b</i>	—	12.30 <i>b</i>	—
	50	44.14 <i>b</i>	5.58 <i>b</i>	10.35 <i>b</i>	−12.58 <i>a</i>	9.46 <i>ab</i>	−40.73 <i>a</i>
	100	32.19 <i>ab</i>	−23.01 a	2.75 <i>a</i>	−76.80 a	3.12 <i>a</i>	−80.48 a
	150	21.64 <i>a</i>	−48.25 <i>A</i>	1.34 <i>a</i>	−88.68	1.89 <i>a</i>	−88.14
r1	0	44.92 <i>b</i>	—	12.77 <i>b</i>	—	13.08 <i>b</i>	—
	50	40.45 <i>b</i>	−9.95 <i>ab</i>	12.74 <i>b</i>	−0.26 <i>ab</i>	16.81 <i>b</i>	−8.67 <i>ab</i>
	100	35.45 <i>ab</i>	−21.07 a	6.83 <i>ab</i>	−46.51 ab	9.84 <i>ab</i>	−46.53 ab
	150	25.36 <i>a</i>	−43.55 <i>A</i>	2.02 <i>a</i>	−84.18	3.04 <i>a</i>	−83.47
r2	0	32.86	—	3.64 <i>c</i>	—	3.20 <i>b</i>	—
	50	20.65	−37.16 <i>a</i>	2.12 <i>b</i>	−41.84 <i>a</i>	2.73 <i>ab</i>	−14.60 <i>ab</i>
	100	22.00	−33.05 a	0.73 <i>a</i>	−80.02 a	0.56 <i>a</i>	−82.38 a
	150	—	—	—	—	—	—

^aLowercase letters refer to a comparison of the plant height and shoot and root weight of *E. crus-galli* for NaCl concentrations within the same population.

^bThe plant height and shoot and root weight variations of *E. crus-galli* among populations are indicated in lowercase letters in italics, lowercase letters in bold, and uppercase letters for 50, 100, and 150 mM concentrations, respectively. Different letters indicate significantly different values, according to Tukey's honest significant difference test. Where no letters are present, the comparison was not significant.

biomass (Wilson and Read 2006). A moderate reduction in plant height in response to salt allows *E. crus-galli* to be competitive in saline environments, even when the plant biomass is reduced, and it has been demonstrated that most of the competitive ability of this species is due to its fast growth and ability to shade other plants as a result of its height (Percy et al. 1981; Wilson and Read 2006). Moreover, the allocation of resources toward plant height instead of biomass may be an adaptation of this species to abiotic stresses that permits the species to succeed in difficult environments (Wilson and Read 2006).

In this study, the root weight reduction in all populations showed a similar trend to that of the shoot weight reduction. Such a similarity was also observed in another study on the same species; however, in that study, *E. crus-galli* was still able to produce roots at 24 dS m⁻¹ EC, while in present experiment, no plants survived after 150 mM NaCl, which could approximately correspond to 15 dS m⁻¹ EC (Chauhan et al. 2013; Government of South Australia 2015).

Chlorophyll and Carotenoid Content

The chlorophyll *a* content increased in all populations as the salt level increased, except in population r2, which showed a moderate reduction, with a maximum of 3%, compared with the control (Table 3). The greatest increase in chlorophyll *a* (about 40% compared with the control) was observed in population s1 for all salt concentrations, followed by population r1, while populations s2 and r1 showed intermediate values with a more moderate increase. The chlorophyll *a* content increased slightly in population s3, reaching a maximum of 5% for 150 mM NaCl. The chlorophyll *b* content displayed an even greater increase for higher salt concentrations compared with the control. The only exception was population s3, which showed reduction in the chlorophyll *b* content of about 20% for 50 and 100 mM and about 40% for 150 mM NaCl compared with the control. All the other populations showed a marked increase in chlorophyll *b* content for 50 mM compared with the control, with values above 90% in populations s1 and s2. Populations s1 and r1 showed a further increase for 100 mM, that is, above 100%, compared with the control, while populations s2 and r2 showed lower values. Apart from populations s3 and r2, the chlorophyll *b* content was still higher for 150 mM in all the other populations than that observed at 0 mM NaCl, albeit with lower values than observed with the other salt concentrations.

Carotenoid content showed a more variable behavior among the populations; decrease in carotenoid content was in fact observed for increasing salt rates for populations s2, r1, and r2, while an increase was detected for populations s1 and s3 compared with the control. A more pronounced decrease was displayed by the ALS inhibitor-resistant populations for all salt concentrations, while population s1 showed the highest increase in carotenoid content, that is, an increase of more than 100% for 150 mM.

Salinity can affect photosynthesis by inducing stomatal closure and a reduction in carbon assimilation, thereby resulting in growth limitation in the short term (Acosta-Motos et al. 2017; Parida and Das 2005). In the long term, salinity can cause the accumulation of salt in the leaves and a reduction in the chlorophyll and carotenoid contents (Acosta-Motos et al. 2017). In our study, the photosynthetic pigments increased in the majority of the populations under salt stress, so it is probable that the salinity stress did not last long enough to cause a reduction in the pigment content, as this is generally a long-term response to salt (Acosta-Motos et al. 2017; Stepien and Johnson 2009). However, not only were the chlorophyll and carotenoid contents not reduced under saline conditions,

they also showed an increase, thus indicating the tolerance of *E. crus-galli* to salt; a rise in chlorophyll under saline conditions is considered a marker of salt tolerance in plants, as observed in other salt-tolerant species (Acosta-Motos et al. 2015; Ashraf and Harris 2013; Stepien and Johnson 2009).

In our study, population r2 displayed a reduction in both chlorophyll *a* and carotenoid contents, thus confirming it was the most sensitive to salt. Another indicator of salt tolerance is the high chlorophyll *a/b* ratio, which is known to be an indicator of health and nonstressed plants (Duarte et al. 2013). In our study, most populations showed a decreasing value for the chlorophyll *a/b* ratio as the salt concentration increased up to 100 mM (Table 3); only population s3 showed an increase in this ratio at increasing salt concentrations, while a more variable behavior was shown by population r2. This could indicate that even though such plants have a certain degree of salt tolerance, salinity induced stress and slowed down photosynthesis, as also demonstrated by the reduction in seedling weight.

Carotenoid content was reduced more compared with chlorophyll content in presence of salt; a reduction in carotenoids has been found to be caused by both oxidation due to ROS and destruction of carotenoids due to salt (Abdel Latef et al. 2020). However, populations s1 and s3 showed an increase in carotenoid content for increasing levels of salt, thus demonstrating they were less sensitive to salinity; the high carotenoid content in these populations could have protected the chlorophyll, thereby preventing its photo-oxidation (Toscano et al. 2019).

Oryza sativa (Weedy Rice and Rice)

Seedling Development

Plant Emergence. Seedling emergence of *O. sativa* and rice was affected more by salinity than *E. crus-galli* emergence was. Plant emergence for 0 mM NaCl was quite similar for all the *O. sativa* populations and Baldo, with values of about 90% or above; only CL80, the variety tolerant to imazamox, displayed values of about 60% (Figure 2). The comparison of parameter *d* of the curve showed an initial different emergence among all rice and *O. sativa* populations, including Baldo, and the CL80 resistant variety (Table 4). Slope was instead more similar between curves, and the only detected differences were between CL80 and both populations wr2 and wr3. The point of inflection *e* of the curves was different between populations, as differences were found between population wr1 and all the other populations and rice varieties and between wr3 and Baldo. The most similar curves, showing nonsignificant differences between parameters and SI, were wr2 versus wr3 and wr2 versus Baldo (Table 4).

Emergence was quite similar to that of the control for 50 mM NaCl. A more marked emergence reduction occurred for 100 mM compared with the control, particularly in the CL80 variety, which showed a 30% decrease in emergence, and an even more marked decrease occurred for 150 mM. Population wr1 displayed the highest emergence percentage for 200 mM, with an average emergence value of about 43%, while the lowest rate of emergence was recorded for Baldo (about 17%), CL80, and wr2, with about 19% of germination, which corresponds to a reduction of about 80% compared with the control. All the *O. sativa* populations and rice varieties showed an emergence reduction of at least 90% for the highest NaCl concentration, compared with the control. A slightly higher emergence was recorded in the Baldo variety (8% emergence) for this salt rate, while emergence did not occur in the wr2 *O. sativa* population.


Table 3. Chlorophyll *a*, chlorophyll *b*, chlorophyll *a/b* ratio, and carotenoid content of the different *Echinochloa crus-galli* populations and the percentage difference of these values for each salt concentration compared with the control.^a

Population	NaCl concentration	Chl <i>a</i> ^b	Chl <i>a</i> difference compared with the control ^c	Chl <i>b</i> ^b	Chl <i>b</i> difference compared with the control ^c	Chl <i>a/b</i> ratio ^b	Carotenoid ^b	Carotenoid difference compared with the control ^c
	mM	mg g ⁻¹ SW	%	mg g ⁻¹ SW	%		mg g ⁻¹ SW	%
s1	0	0.92 a	—	0.41 a	—	3.59 d	0.15 a	—
	50	1.32 c	42.96 e	0.89 c	114.99 d	1.49 b	0.23 c	54.57 e
	100	1.31 b	41.29 e	1.01 d	144.69 e	1.29 a	0.21 b	42.99 e
	150	1.33 d	44.27 D	0.57 b	37.22 B	2.36 c	0.34 d	129.56 C
s2	0	1.25 a	—	0.47 a	—	2.65 c	0.31 b	—
	50	1.33 c	6.54 c	0.91 d	91.78 c	1.68 b	0.27 a	-12.16 c
	100	1.32 b	5.79 c	0.84 c	77.75 c	1.57 a	0.28 a	-10.60 c
	150	1.39 d	11.33 B	0.82 b	73.87 D	1.60 a	0.34 c	9.44 A
s3	0	1.29 a	—	1.17 c	—	1.11 a	0.19 a	—
	50	1.33 c	2.67 b	0.92 b	-21.30 a	1.45 b	0.25 b	32.76 d
	100	1.32 b	2.09 b	0.92 b	-21.05 a	1.47 b	0.25 b	35.65 d
	150	1.36 d	5.33 A	0.69 a	-40.69 A	1.97 c	0.34 c	82.64 B
r1	0	1.14 a	—	0.63 a	—	2.13 d	0.28 c	—
	50	1.32 c	15.77 d	0.95 b	50.90 b	1.39 b	0.23 b	-19.53 b
	100	1.28 b	12.88 d	1.30 c	106.34 d	0.98 a	0.14 a	-51.47 b
	150	1.38 d	21.33 C	0.95 b	50.98 C	1.44 c	0.31 d	-8.92 A
r2	0	1.31 c	—	0.70 a	—	1.88 c	0.33 c	—
	50	1.26 a	-3.46 a	1.49 c	113.51 d	0.85 a	0.06 a	-80.39 a
	100	1.30 b	-0.41 a	1.14 b	63.01 b	1.15 b	0.20 b	-39.36 a
	150	—	—	—	—	—	—	—

^aAbbreviations: Chl, chlorophyll; SW, shoot fresh weight.
^bLowercase letters refer to the comparison of the *E. crus-galli* chlorophyll *a*, chlorophyll *b*, chlorophyll *a/b* ratio, and carotenoid content for the NaCl concentrations within the same population.
^cThe *E. crus-galli* chlorophyll *a*, chlorophyll *b*, and carotenoid content variations among populations are indicated in lowercase letters in italics, lowercase letters in bold, and uppercase letters for 50, 100, and 150 mM concentrations, respectively. Different letters indicate significantly different values according to Tukey's honest significant difference test. Where no letters are present, the comparison was not significant.

Table 4. Curve parameter (Equation 1) and sensitivity index (SI) (Equation 2) comparison between *Oryza sativa* populations/varieties (estimate and P-values).

Population comparison	Parameter <i>b</i> comparison		Parameter <i>d</i> comparison		Parameter <i>e</i> comparison		SI	
	Estimate	P-value	Estimate	P-value	Estimate	P-value	Estimate	P-value
wr1/wr2	1.37	0.450	0.92	0.080	1.22	0.000*	1.22	0.000*
wr1/wr3	1.11	0.790	0.97	0.544	1.13	0.008*	1.13	0.008*
wr1/CL80	3.05	0.086	1.43	0.000*	1.40	0.005*	0.71	0.000*
wr1/Baldo	1.95	0.230	1.00	0.997	1.34	0.000*	0.748	0.000*
wr2/wr3	0.81	0.330	1.05	0.268	0.92	0.056	0.926	0.056
wr2/CL80	2.22	0.050*	1.55	0.000*	1.14	0.206	0.87	0.148
wr2/Baldo	1.42	0.330	1.09	0.224	1.09	0.215	0.91	0.175
wr3/CL80	2.75	0.028*	1.47	0.000*	1.23	0.055	0.81	0.018*
wr3/Baldo	1.76	0.163	1.03	0.656	1.18	0.025*	0.85	0.008*
CL80/Baldo	0.64	0.099	0.70	0.000*	0.95	0.683	1.05	0.696

*Significant comparisons at $P \leq 0.05$.**Figure 2.** Dose-response emergence curve with the average data points of the different *Oryza sativa* (weedy rice) populations and rice varieties against the salt concentration. Curve parameter estimates (Equation 1): wr1 ($b=9.40$, $d=87.14$, $e=195.80$), wr2 ($b=9.40$, $d=87.14$, $e=160.19$), wr3 ($b=8.49$, $d=89.73$, $e=173.01$), Baldo ($b=4.81$, $d=87.12$, $e=146.49$), CL80 ($b=3.08$, $d=60.89$, $e=140.04$). The salt concentration required to reduce the emergence by 50% (EC₅₀) and the significant pairwise comparisons between EC₅₀ (SI index) are shown below the graph (Equation 2).

The NaCl concentration required to reduce emergence by 50% (EC₅₀) was between 140 mM for CL80 and 195 mM for the wr1 population. A pairwise comparison between EC₅₀ values (SI index) showed that wr1 had a significantly higher EC₅₀ than all the tested *O. sativa* populations and rice varieties, thus suggesting a higher tolerance to salt. Moreover, both rice varieties showed a significantly lower value of EC₅₀ than the wr3 *O. sativa* population.

Oryza sativa and rice showed a higher sensitivity to salt than *E. crus-galli*, as demonstrated by the lower EC₅₀ values. Moreover, similar values for emergence reduction were recorded for a salt concentration of 250 mM in *E. crus-galli*, but for lower

concentrations in rice and *O. sativa*, that is, for values between 150 and 200 mM (Figures 1 and 2). Previous studies also demonstrated that a moderate level of soil salinity (between 40 and 100 mM) still permits rice to grow, but some weeds, such as *E. crus-galli*, can germinate and even grow better than rice under these conditions (Chauhan and Johnson 2009). Another study graded *O. sativa* as moderately tolerant to salinity, while *E. crus-galli* was classified within the salt-tolerant species, as its germination was about 30% for salt concentrations of 240 mM (Hakim et al. 2011). A previous study on the sensitivity of Italian *O. sativa* populations and rice varieties to salt showed that CL80 was affected more by salt, in terms of germination, than other herbicide-sensitive *O. sativa* populations and the conventional Baldo rice variety, which was the least affected by salt (Fogliatto et al. 2019). In this study, the Baldo variety was less tolerant to salt, in terms of emergence, compared with germination, as the wr1 *O. sativa* population showed a higher emergence rate for increasing salt levels. This was also demonstrated by the higher EC₅₀ displayed by wr1 compared with all other *O. sativa* and rice varieties, including Baldo. The CL80 variety was confirmed to be salt sensitive, even at emergence, as it showed the highest emergence reduction across all salt concentrations. A certain variability in salt tolerance among rice varieties was demonstrated in previous studies, and the Baldo variety was classified as moderately sensitive or tolerant to salt according to the varieties to which it was compared (Bertazzini et al. 2018). Apart from the abovementioned study on germination, which highlighted a high level of sensitivity to saline conditions in the CL80 variety, no information regarding the sensitivity of this variety to salt is available (Fogliatto et al. 2019).

Plant Height, Shoot, and Root Weight. *Oryza sativa* and rice were observed to be sensitive to salt, as they were only able to develop seedlings for a concentration of 50 mM NaCl. After emergence, seedlings were not able to survive at higher salt levels, thus demonstrating the high sensitivity of *O. sativa* to salt at early seedling stages compared with emergence (Table 5).

The difference in plant height between the control and 50 mM NaCl was only significant for the CL80 variety, for which salt reduced the height by about 35% compared with the control (Table 5). Population wr2 showed a slight reduction in height (about 3%) with salt, while all the other populations were slightly stimulated, with Baldo and wr3 recording the largest increases. The reduction in shoot weight due to salt was again only significant for CL80 (about 66% reduction compared with the control). All the other populations and Baldo showed a more moderate weight

Table 5. Plant height and shoot and root weight of the different *Oryza sativa* populations (weedy rice) and rice varieties and the percentage difference of these growth parameters for each salt concentration compared with the control.

Population	NaCl concentration	Plant height	Plant height variation compared with the control ^a	Shoot weight	Shoot weight variation compared with the control ^a	Root weight	Root weight variation compared with the control ^a
	mM	cm	%	g	%	g	%
wr1	0	16.97	—	1.93	—	1.55	—
	50	17.14	1.03 ab	1.56	−19.14 b	1.04	−33.05 ab
wr2	0	19.08	—	1.94	—	1.86	—
	50	18.43	−3.41 ab	1.41	−27.32 ab	0.96*	−48.57 a
wr3	0	18.09	—	1.51	—	0.73	—
	50	20.25	11.95 b	1.51	0.00 b	0.67	−8.64 b
Baldo	0	20.04	—	2.72	—	2.92	—
	50	23.23	15.92 b	2.06	−24.36 ab	1.80*	−38.43 ab
CL80	0	12.61	—	0.29	—	0.27	—
	50	8.20*	−34.97 a	0.10*	−65.91 a	0.13*	−51.23 a

^aLowercase letters refer to the growth parameter variations among populations for 50 mM. Different letters indicate significantly different values, according to Tukey's honest significant difference (HSD) test. Where no letters are present, the comparison was not significant.

*Significant differences of the growth parameters, according to Tukey's HSD test, between the two NaCl concentrations (0 mM and 50 mM) for each population/rice variety.

reduction, while population wr3 did not display any shoot weight reduction compared with the control. Root weight was affected more by salt than shoot weight, as a significant root weight decrease was observed for both rice varieties and wr2. The highest root weight reduction was observed for CL80 (about 51% reduction) and wr2 (about 48% reduction), while population wr3 was affected the least (about 8% reduction). In general, at the seedling stage, *O. sativa* population wr3 seemed to be able to better tolerate saline conditions when compared with the resistant CL80, as it showed a significantly greater plant height and lower root and shoot weight reduction. Population wr3 also showed a lower root weight reduction compared with wr2, in contrast to what was observed for emergence, in which population wr1 was the most tolerant. As already observed for emergence, among all populations and varieties, CL80 was the most sensitive to salt.

A lower tolerance of rice to salinity, compared with *E. crus-galli*, was found in previous studies in which rice was not able to develop seedlings at a high salt level, as rice is more sensitive to salt at emergence and in the early growth stages (Aslam et al. 1993; Chauhan et al. 2013; Korres et al. 2019). When comparing the differences in salt tolerance between rice and *O. sativa*, it was found that the latter required higher salt levels than rice to achieve a 50% reduction in dry weight, thus highlighting an advantage of the weed over the crop in saline environments (Cirillo et al. 2018; Tanji and Kielen 2002). In our study, regarding growth parameters, we found more variable behavior: CL80 was found to be more sensitive than the *O. sativa* population wr3 and Baldo, which showed an intermediate tolerance, and was similar to that of some *O. sativa* populations.

Plant height of rice and *O. sativa* was not significantly reduced by salt, except for CL80, which appeared to be highly sensitive to salinity. Previous studies conducted on different rice varieties with low salt concentrations showed either a reduction in plant height for the most sensitive varieties or no change in plant height in the case of more tolerant varieties, thus highlighting a wide variability among genotypes (Bertazzini et al. 2018; Hakim et al. 2014; Motamed et al. 2008). In our study, we confirmed both results, as only the most salt-sensitive variety showed a height decrease.

The shoot and root weight were reduced more in the presence of salt, and this can be attributed to reduced photosynthesis and turgor, salt toxicity, and a disruption in mineral nutrition (Bertazzini et al. 2018; Hakim et al. 2014c). Some studies found that root and shoot weight were reduced by salt to a similar extent; in our study, we found a greater reduction in root weight. A previous study

found that the rate of Na⁺ accumulation in rice was higher in roots than in shoots, and this resulted in a reduction in root biomass (Hakim et al. 2014c).

Chlorophyll and Carotenoid Contents

Chlorophyll content showed variable behavior in the presence of salt, depending on *O. sativa* population and rice variety (Table 6). In fact, chlorophyll *a* content increased at 50 mM in populations wr2, wr3, and Baldo compared with the control, while it was reduced at a greater extent in wr1 and CL80. The wr1 *O. sativa* population showed the greatest reduction in chlorophyll, about 80%, followed by CL80, which recorded a reduction of about 50%. Baldo showed the greatest increase in the presence of salt (about 68%). A similar trend was noted for chlorophyll *b* content for all populations and rice varieties, with reduced or increased values similar to those for chlorophyll *a*; the only exception was population wr3, which had a reduction (about 16% decrease) in chlorophyll *b* content. Carotenoid content decreased to a great extent in population wr1, that is, a reduction of more than 70% was observed in the presence of salt, and this was followed by CL80, with a reduction above 67%. Carotenoid content in populations wr2, wr3, and Baldo increased as salinity increased, with the highest value being recorded for Baldo (about 78% increase, compared with the control).

The higher sensitivity to saline conditions of *O. sativa* and rice compared with *E. crus-galli* was also confirmed by the chlorophyll and carotenoid contents, which often showed a decrease in the presence of salt, in particular in the most sensitive *O. sativa* population (wr1) and rice variety (CL80); similar results were found in a previous study that showed a greater reduction of photosynthetic pigments in *O. sativa* than in *E. crus-galli* (Hakim et al. 2014b). Concerning the growth parameters (plant height, shoot and root weight), CL80 was more sensitive to salt than all other *O. sativa* and rice varieties; this sensitivity was also confirmed by the low photosynthetic pigment content. However, population wr1, which showed an intermediate sensitivity to salt in the growth parameters, recorded the highest chlorophyll and carotenoid content reduction, thus highlighting a greater sensitivity, which would probably also have been observed in seedling growth if the salt conditions had lasted longer. Baldo and wr3 were once again confirmed to be the most tolerant to salt when photosynthetic pigments were considered.

The chlorophyll *a/b* ratio rose for increasing salt concentrations in all *O. sativa* populations and Baldo, while it did not vary at 50

Table 6. Chlorophyll *a*, chlorophyll *b*, chlorophyll *a/b* ratio, and carotenoid content of the different *Oryza sativa* (weedy rice) populations and rice varieties and the percentage difference of these values for each salt concentration compared with the control.^a

Population	NaCl concentration	Chl <i>a</i>	Chl <i>a</i> variation compared with the control ^b	Chl <i>b</i>	Chl <i>b</i> variation compared with the control ^b	Chl <i>a/b</i> ratio	Carotenoid	Carotenoid variation compared with the control ^b
	mM	mg g ⁻¹ SW	%	mg g ⁻¹ SW	%		mg g ⁻¹ SW	%
wr1	0	1.10	—	0.20	—	5.48	1.13	—
	50	0.21*	-81.20 a	0.03*	-85.45 a	7.08*	0.31*	-72.90 a
wr2	0	0.96	—	0.17	—	5.68	1.02	—
	50	1.22*	26.67 d	0.27*	56.92 d	4.58*	1.28*	25.26 d
wr3	0	0.53	—	0.08	—	6.23	0.64	—
	50	0.54*	2.39 c	0.07*	-16.66 c	7.67*	0.78*	21.13 c
Baldo	0	0.54	—	0.14	—	3.85	0.68	—
	50	0.90*	68.13 e	0.23*	67.90 d	3.96*	1.21*	78.50 e
CL80	0	0.97	—	0.12	—	8.28	0.78	—
	50	0.48*	-50.51 b	0.07*	-43.70 b	7.27	0.25*	-67.58 b

^aAbbreviations: Chl, chlorophyll; SW, shoot fresh weight.^bLowercase letters refer to the chlorophyll and carotenoid content variations among populations for 50 mM. Different letters indicate significantly different values according to Tukey's honest significant difference (HSD) test. Where no letters are present, the comparison was not significant.^cSignificant variations of the chlorophyll, chlorophyll *a/b* ratio, and carotenoid content, according to Tukey's HSD test, between the two NaCl concentrations (0 mM and 50 mM) for each population/rice variety.

mM compared with 0 mM in CL80 (Table 6); as a high ratio is an indicator of a healthy plant, this confirmed the higher salt sensitivity of CL80 (Duarte et al. 2013). Even though population wr1 displayed the greatest pigment reduction, it also recorded a high chlorophyll *a/b* ratio; this behavior could probably be explained by the fact that during the degradation of chlorophyll *b* due to salt, to prevent further damage, chlorophyll *b* may be converted to chlorophyll *a*, which is the primary photosynthetic pigment (Ashraf and Harris, 2013). The increase or moderate decrease in the chlorophyll content in salt-tolerant rice varieties observed in this study, as opposed to a marked decrease for salt-sensitive rice varieties, was also found in previous studies, thereby highlighting a high variability in salt response within *Oryza sativa* species (Kibria et al. 2017).

This study has highlighted a different tolerance to salinity of the tested species; *E. crus-galli* was found to be more tolerant than both *O. sativa* and rice. The level of tolerance varied according to the considered growth stage, and it was higher at emergence and lower at the seedling stage. The two weed species and rice were able to emerge up to 250 mM NaCl, while *E. crus-galli* was only able to grow for concentrations of up to 150 mM NaCl at the seedling stage, while rice and *O. sativa* only developed seedlings up to 50 mM. Previous studies found that salinity tolerance at the seedling stage was not correlated with salinity tolerance at other growth stages in rice (Ferdose et al. 2009; Kakar et al. 2019). A further demonstration of this finding is the behavior of population wr1, which was the most salt tolerant at emergence, but not at the seedling stage.

In some cases, relatively low concentrations of salt can stimulate plant growth, acting as a hormetic effect (Calabrese 2013). In our study, hormesis was observed in *E. crus-galli* emergence and in the plant height of almost all populations for 50 mM NaCl. This phenomenon was less evident in *O. sativa* and rice, even though it was observed in some populations for 50 mM in terms of emergence (wr1 and wr2) and plant height (wr1, wr2, and Baldo). This phenomenon was reported in previous studies; in China, for example, some weed species were stimulated to germinate in soil taken from coastal areas with a moderate level of salinity (Bai et al. 2014).

Wide variability in response to salt was not only found for the different species but also among populations within the same species. The tested populations of both *E. crus-galli* and *O. sativa* responded variably to salt, in terms of growth parameters. Baldo and CL80 demonstrated different salt tolerance, as CL80 was observed to be more sensitive than Baldo. Wide growth variability has been demonstrated for both *E. crus-galli* and *O. sativa*, even under nonsaline conditions, as weeds are generally characterized by wide phenotypic plasticity, as well as genetic variability, in response to different environmental conditions (Chauhan and Johnson 2009; Fogliatto et al. 2011, 2012).

This study also aimed to establish whether ALS inhibitor-resistant and ALS inhibitor-sensitive *E. crus-galli* populations and rice varieties responded differently to salinity. The ALS inhibitor-resistant *E. crus-galli* population r2 was affected more by salt, in terms of emergence, seedling growth, and carotenoid content, but this was less evident in the other resistant population r1; a behavior similar to that of r2 was observed for CL80, which is tolerant to an ALS-inhibiting herbicide. Even though only a few resistant populations were tested, the study highlighted a common behavior in response to salt in two out of three herbicide-resistant populations (r2 population for *E. crus-galli* and CL80); this behavior is probably caused by a fitness penalty that may be associated

with herbicide resistance, even though specific studies are needed to confirm this hypothesis (Keshtkar et al. 2019). Greater growth performance and higher competitive ability of herbicide-sensitive populations were also reported in a previous germination study on *O. sativa* and rice, while similar competitive ability was observed when comparing *E. crus-galli* populations resistant and sensitive to propanil and clomazone (Bagavathiannan et al. 2011; Fogliatto et al. 2019). However, previous studies carried out with other species that are resistant to different herbicides found a higher growth performance and tolerance to salt in resistant populations (Shrestha et al. 2018). A study conducted on junglerice [*Echinochloa colona* (L.) Link] found that glyphosate-resistant biotypes were more competitive, germinate better, and produce more biomass under moisture and saline conditions than sensitive populations (Shrestha et al. 2018). Therefore, a lower growth performance of resistant populations in response to diverse environmental stresses cannot be generalized, and it may vary for different species, different populations within a species, and different herbicides and their related modes of action (Shrestha et al. 2018).

This study highlighted that Baldo performed better under high salt concentrations than CL80; thus the availability of salt-tolerant rice varieties is particularly important, as the majority of the rice areas throughout the world are located near coastal areas and river deltas where salinity problems are more common (Formentin et al. 2018). Moreover, some weeds can be more competitive than others in saline environments (Fogliatto et al. 2020); according to our study, *E. crus-galli* is favored more than *O. sativa* under saline conditions, and thus more efforts should be made to control this weed. Populations resistant to ALS inhibitors showed variable behavior, with some populations that tolerate salt (such as the *E. crus-galli* r1 population) and others that can be more sensitive to salt (such as the *E. crus-galli* population r2 and the CL80 rice variety).

Acknowledgments. This research received no specific grant from any funding agency or the commercial or not-for-profit sectors. No conflicts of interest have been declared.

References

- Abdel Latef AAH, Abu Alhmad MF, Kordrostami M, Abo-Baker A-BA-E, Zakir A (2020) Inoculation with *Azospirillum lipoferum* or *Azotobacter chroococcum* reinforces maize growth by improving physiological activities under saline conditions. *J Plant Growth Regul* 39:1293–1306
- Acosta-Motos J-R, Diaz-Vivancos P, Álvarez S, Fernández-García N, Sanchez-Blanco MJ, Hernández JA (2015) Physiological and biochemical mechanisms of the ornamental *Eugenia myrtifolia* L. plants for coping with NaCl stress and recovery. *Planta* 242:829–846
- Acosta-Motos JR, Ortuño MF, Bernal-Vicente A, Diaz-Vivancos P, Sanchez-Blanco MJ, Hernandez JA (2017) Plant responses to salt stress: adaptive mechanisms. *Agronomy* 7:18
- Ahuja I, de Vos RCH, Bones AM, Hall RD (2010) Plant molecular stress responses face climate change. *Trends Plant Sci* 15:664–674
- Akbari M, Alamdarlo HN, Mosavi SH (2020) The effects of climate change and groundwater salinity on farmers' income risk. *Ecol Indic* 110:105893
- Andres A, Fogliatto S, Ferrero A, Vidotto F (2015) Growth variability of Italian weedy rice populations grown with or without cultivated rice. *Crop Sci* 55:394–402
- Ashraf M, Harris PJC (2013) Photosynthesis under stressful environments: an overview. *Photosynthetica* 51:163–190
- Aslam M, Qureshi RH, Ahmed N (1993) A rapid screening technique for salt tolerance in rice (*Oryza sativa* L.). *Plant Soil* 150:99–107
- Badia D (2000) Potential nitrification rates of semiarid cropland soils from the Central Ebro valley, Spain. *Arid Soil Res Rehabil* 14:281–292
- Bagavathiannan MV, Norsworthy JK, Jha P, Smith K (2011) Does resistance to propanil or clomazone alter the growth and competitive abilities of barnyardgrass (*Echinochloa crus-galli*)? *Weed Sci* 59:353–358
- Bai J, Huang L, Gao Z, Lu Q, Wang J, Zhao Q (2014) Soil seed banks and their germination responses to cadmium and salinity stresses in coastal wetlands affected by reclamation and urbanization based on indoor and outdoor experiments. *J Hazard Mater* 280:295–303
- Bertazzini M, Sacchi GA, Forlani G (2018) A differential tolerance to mild salt stress conditions among six Italian rice genotypes does not rely on Na⁺ exclusion from shoots. *J Plant Physiol* 226:145–153
- Calabrese EJ (2013) Hormetic mechanisms. *Crit Rev Toxicol* 43:580–606
- Chauhan BS, Abugho SB, Amas JC, Gregorio GB (2013) Effect of salinity on growth of barnyardgrass (*Echinochloa crus-galli*), horse purslane (*Trianthema portulacastrum*), junglerice (*Echinochloa colona*), and rice. *Weed Sci* 61:244–248
- Chauhan BS, Johnson DE (2009) Seed germination ecology of junglerice (*Echinochloa colona*): a major weed of rice. *Weed Sci* 57:235–240
- Cirillo V, Masin R, Maggio A, Zanin G (2018) Crop-weed interactions in saline environments. *Eur J Agron* 99:51–61
- Duarte B, Santos D, Marques JC, Caçador I (2013) Ecophysiological adaptations of two halophytes to salt stress: photosynthesis, PS II photochemistry and anti-oxidant feedback—implications for resilience in climate change. *Plant Physiol Biochem* 67:178–188
- Ferdose J, Kawasaki M, Taniguchi M, Miyake H (2009) Differential sensitivity of rice cultivars to salinity and its relation to accumulation and root tip structure. *Plant Prod Sci* 12:453–461
- Fogliatto S, Serra F, Patrucco L, Milan M, Vidotto F (2019) Effect of different water salinity levels on the germination of imazamox-resistant and sensitive weedy rice and cultivated rice. *Agronomy* 9:658
- Fogliatto S, Vidotto F, Ferrero A (2011) Germination of weedy rice in response to field conditions during winter. *Weed Technol* 25:252–261
- Fogliatto S, Vidotto F, Ferrero A (2012) Morphological characterisation of Italian weedy rice (*Oryza sativa*) populations. *Weed Res* 52:60–69
- Fogliatto S, Vidotto F, Ferrero A (2020) How can weedy rice stand against abiotic stresses? A review. *Agronomy* 10:1284
- Formentin E, Sudiro C, Perin G, Riccadonna S, Barizza E, Baldoni E, Lavezzo E, Stevanato P, Sacchi GA, Fontana P, Toppo S, Morosinotto T, Zottini M, Lo Schiavo F (2018) Transcriptome and cell physiological analyses in different rice cultivars provide new insights into adaptive and salinity stress responses. *Front Plant Sci* 9:204
- Gay F, Maraval I, Roques S, Gunata Z, Boulanger R, Audebert A, Mestres C (2010) Effect of salinity on yield and 2-acetyl-1-pyrroline content in the grains of three fragrant rice cultivars (*Oryza sativa* L.) in Camargue (France). *Field Crops Res* 117:154–160
- Goulart ICGR, Borba TCO, Menezes VG, Merotto A (2014) Distribution of weedy red rice (*Oryza sativa*) resistant to imidazolinone herbicides and its relationship to rice cultivars and wild *Oryza* species. *Weed Sci* 62:280–293
- Goulart ICGR, Matzenbacher FO, Merotto A (2012) Differential germination pattern of rice cultivars resistant to imidazolinone herbicides carrying different acetolactate synthase gene mutations: differential germination pattern of rice cultivars. *Weed Res* 52: 224–232
- Government of South Australia, (2015). Measuring Salinity. <https://landscape.sa.gov.au/mr/publications/measuring-salinity>. Accessed: June 12, 2020
- Guo Z, Yang N, Zhu C, Gan L (2017) Exogenously applied poly-γ-glutamic acid alleviates salt stress in wheat seedlings by modulating ion balance and the antioxidant system. *Env Sci Pollut Res* 24:6592–6598
- Hakim MA, Juraimi AS, Hanafi MM, Ali E, Ismail MR, Selamat A, Karim SMR (2014a) Effect of salt stress on morpho-physiology, vegetative growth and yield of rice. *J Environ Biol* 35:317–326
- Hakim MA, Juraimi AS, Hanafi MM, Ismail MR, Rafii MY, Aslani F, Selamat A (2014b) The effect of salinity on chlorophyll, proline and mineral nutrients in common weeds of coastal rice fields in Malaysia. *J Environ Biol* 35:855–864
- Hakim MA, Juraimi AS, Hanafi MM, Ismail MR, Rafii MY, Islam MM, Selamat A (2014c) The effect of salinity on growth, ion accumulation and yield of rice varieties. *J Anim Plant Sci* 24:874–885
- Hakim MA, Juraimi AS, Hanafi MM, Selamat A, Ismail MR, Karim SMR (2011) Studies on seed germination and growth in weed species of rice field under salinity stress. *J Environ Biol* 32:529–536

- Heap I (2020) The International Herbicide-Resistant Weed Database. <http://weedsdatabase.org/Home.aspx>. Accessed: June 1, 2020
- Hoang TML, Tran TN, Nguyen TKT, Williams B, Wurm P, Bellairs S, Mundree S (2016) Improvement of salinity stress tolerance in rice: challenges and opportunities. *Agronomy* 6:54
- Isla R, Aragüés R, Royo A (2003) Spatial variability of salt-affected soils in the middle Ebro Valley (Spain) and implications in plant breeding for increased productivity. *Euphytica* 134:325–334
- Kakar N, Jumaa SH, Redoña ED, Warburton ML, Reddy KR (2019) Evaluating rice for salinity using pot-culture provides a systematic tolerance assessment at the seedling stage. *Rice* 12:57
- Keshtkar E, Abdolshahi R, Sasanfar H, Zand E, Beffa R, Dayan FE, Kudsk P (2019) Assessing fitness costs from a herbicide-resistance management perspective: a review and insight. *Weed Sci* 67:137–148
- Kibria MG, Hossain M, Murata Y, Hoque MdA (2017) Antioxidant defense mechanisms of salinity tolerance in rice genotypes. *Rice Sci* 24:155–162
- Korres NE, Burgos NR, Travlos I, Vurro M, Gitsopoulos TK, Varanasi VK, Duke SO, Kudsk P, Brabham C, Rouse CE, Salas-Perez R (2019) New directions for integrated weed management: modern technologies, tools and knowledge discovery. *Adv Agron* 155:243–319
- Korres NE, Norsworthy JK, Tehranchian P, Gitsopoulos TK, Loka DA, Oosterhuis DM, Gealy DR, Moss SR, Burgos NR, Miller MR, Palhano M (2016) Cultivars to face climate change effects on crops and weeds: a review. *Agron Sustain Dev* 36:12
- Läuchli A, Grattan SR (2007) Plant growth and development under salinity stress. Pages 1–32 in Jenks MA, Hasegawa PM, Mohan Jain S, eds. *Advances in Molecular Breeding toward Drought and Salt Tolerant Crops*. Dordrecht, Netherlands: Springer Science & Business Media
- Machado RMA, Serralheiro RP (2017) Soil salinity: effect on vegetable crop growth. Management practices to prevent and mitigate soil salinization. *Horticulturae* 3:30
- Motamed MK, Asadi R, Rezaei M, Amiri E (2008) Response of high yielding rice varieties to NaCl salinity in greenhouse circumstances. *Afr J Biotechnol* 7:3866–3873
- Negrão S, Schmöckel SM, Tester M (2017) Evaluating physiological responses of plants to salinity stress. *Ann Bot* 119:1–11
- Opeña JL, Chauhan BS, Baltazar AM (2014) Seed germination ecology of *Echinochloa glabrescens* and its implication for management in rice (*Oryza sativa* L.). *PLoS ONE* 9:e92261
- Parida AK, Das AB (2005) Salt tolerance and salinity effects on plants: a review. *Ecotoxicol Environ Saf* 60:324–349
- Parihar P, Singh S, Singh R, Singh VP, Prasad SM (2015) Effect of salinity stress on plants and its tolerance strategies: a review. *Environ Sci Pollut Res* 22:4056–4075
- Pearcy R, Tumosa N, Williams K (1981) Relationships between growth, photosynthesis and competitive interactions for a C₃ and a C₄ Plant. *Oecologia* 48:371–376
- Rahman M, Ungar IA (1990) The effect of salinity on seed germination and seedling growth of *Echinochloa crus-galli*. *Ohio J Sci* 90:13–15
- R Core Team (2019) R: a language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria
- Ritz C, Baty F, Streibig JC, Gerhard D (2015) Dose-response analysis Using R. *PLoS ONE* 10:e0146021
- Ritz C, Cedergreen N, Jensen JE, Streibig JC (2006) Relative potency in non-similar dose-response curves. *Weed Sci* 54: 407–412
- Ritz C, Streibig JC (2016) Analysis of dose-response curve. Package “drc” guide. <https://cran.r-project.org/web/packages/drc/drc.pdf>. Accessed: September 10, 2020
- Serra F, Fogliatto S, Vidotto F (2018) Effect of salinity on *Echinochloa crus-galli* germination as affected by herbicide resistance. *Ital J Agron* 13:221–228
- Shrestha A, deSouza LL, Yang P, Sosnoskie L, Hanson BD (2018) Differential tolerance of glyphosate-susceptible and glyphosate-resistant biotypes of junglerice (*Echinochloa colona*) to environments during germination, growth, and intraspecific competition. *Weed Sci* 66:340–346
- Singh A (2018) Assessment of different strategies for managing the water resources problems of irrigated agriculture. *Agric Water Manag* 208:187–192
- Stepien P, Johnson GN (2009) Contrasting responses of photosynthesis to salt stress in the glycophyte *Arabidopsis* and the halophyte *Thellungiella*: role of the plastid terminal oxidase as an alternative electron sink. *Plant Physiol* 149:1154–1165
- Sudianto E, Beng-Kah S, Ting-Xiang N, Saldain NE, Scott RC, Burgos NR (2013) Clearfield® rice: its development, success, and key challenges on a global perspective. *Crop Prot* 49:40–51
- Tanji KK, Kielen NC (2002) *Agricultural Drainage Water Management in Arid and Semi-arid Areas*. FAO Irrigation and Drainage Paper 6. Rome: Food and Agriculture Organization of the United Nations. 1188 p
- Toscano S, Trivellini A, Cocetta G, Bulgari R, Francini A, Romano D, Ferrante A (2019) Effect of preharvest abiotic stresses on the accumulation of bioactive compounds in horticultural produce. *Front Plant Sci* 10:1212
- Vidotto F, Dalla Valle N, Fogliatto S, Milan M, De Palo F, Tabacchi M, Ferrero A (2020) Rapid increase of herbicide resistance in *Echinochloa* spp. consequent to repeated applications of the same herbicides over time. *Arch Agron Soil Sci*. DOI:10.1080/03650340.2020.1741554
- Vila-Aiub MM, Gundel PE, Preston C (2015) Experimental methods for estimation of plant fitness costs associated with herbicide-resistance genes. *Weed Sci* 63:203–216
- Wilson C, Read JJ (2006) Effect of mixed-salt salinity on growth and ion relations of a barnyardgrass species. *J Plant Nutr* 29:1741–1753
- Wu X, Zhang T, Pan L, Wang L, Xu H, Dong L (2016) Germination requirements differ between fenoxaprop-P-ethyl resistant and susceptible Japanese foxtail (*Alopecurus japonicus*) biotypes. *Weed Sci* 64: 653–663
- Zeng L, Shannon MC (2000) Salinity effects on seedling growth and yield components of rice. *Crop Sci* 40:996
- Zhan LJ, Fontana E, Tibaldi G, Nicola S (2009) Qualitative and physiological response of minimally processed garden cress (*Lepidium sativum* L.) to harvest handling and storage conditions. *J Food Agric Environ* 7:43–50