



## **Resolution of the *Eremophila tietkensii* (Scrophulariaceae) species complex based on congruence between morphological and molecular pattern analyses**

Authors: Curtis, Amy L., Grierson, Pauline F., Batley, Jacqueline, Naaykens, Jeremy, Fowler, Rachael M., et al.

Source: Australian Systematic Botany, 35(1) : 1-18

Published By: CSIRO Publishing

URL: <https://doi.org/10.1071/SB21005>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).







Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.



# Resolution of the *Eremophila tietkensii* (Scrophulariaceae) species complex based on congruence between morphological and molecular pattern analyses

Amy L. Curtis<sup>A,B,\*</sup> , Pauline F. Grierson<sup>B</sup> , Jacqueline Batley<sup>B</sup> , Jeremy Naaykens<sup>C</sup>,  
Rachael M. Fowler<sup>D</sup> , Anita Severn-Ellis<sup>B</sup>  and Kevin R. Thiele<sup>A,B</sup> 

For full list of author affiliations and declarations see end of paper

## \*Correspondence to:

Amy L. Curtis  
Western Australian Herbarium,  
Biodiversity and Conservation Science,  
Department of Biodiversity, Conservation  
and Attractions, Locked Bag 104, Bentley  
Delivery Centre, WA 6983, Australia  
Email: amy.curtis@dbca.wa.gov.au

## Handling Editor:

David Williams

Received: 12 February 2021  
Accepted: 17 November 2021  
Published: 2 March 2022

## Cite this:

Curtis AL et al. (2022)  
*Australian Systematic Botany*  
35(1), 1–18. doi:10.1071/SB21005

© 2022 The Author(s) (or their  
employer(s)). Published by  
CSIRO Publishing.  
This is an open access article distributed  
under the Creative Commons Attribution-  
NonCommercial-NoDerivatives 4.0  
International License (CC BY-NC-ND)

OPEN ACCESS

## ABSTRACT

*Eremophila* R.Br. comprises at least 238 species endemic to Australia, with many more having not yet been formally described. Three putative new taxa, namely, *E. sp.* Hamersley Range (K. Walker KW 136), *E. sp.* Calvert Range (A. A. Burbidge 738) and *E. sp.* Rudall River (P. G. Wilson 10512), were segregated from a broadly defined *E. tietkensii* F.Muell. & Tate by J. Hurter at the Western Australian Herbarium in 2012. Both *E. sp.* Hamersley Range and *E. sp.* Rudall River are listed as being of conservation concern in Western Australia, the former occurring in the Pilbara region in areas of prospective interest for mining development. We sought to determine whether these phrase-named entities should be formally described as new species, using multivariate analyses of morphometric and molecular data derived from specimens in the Western Australia Herbarium. *Eremophila sp.* Rudall River could not be adequately separated from *E. tietkensii* by either morphological or molecular data, and is here included within that species. By contrast, *E. sp.* Hamersley Range and *E. sp.* Calvert Range are clearly morphologically and genetically distinct. We thus describe them here as the new species *E. naaykensii* A.L.Curtis & K.R.Thiele and *E. hurteri* A.L.Curtis & K.R.Thiele. The recognition of these taxa will help inform their conservation prioritisation and subsequent management.

**Keywords:** congruence, ddRADseq, *Eremophila*, herbarium sampling, morphology, Pilbara, taxonomy, Western Australia.

## Introduction

*Eremophila* R.Br. (Scrophulariaceae) comprises ~238 species and ~58 subspecies of perennial shrubs endemic to arid and semi-arid Australia (Chinnock 2007a). Species diversity is highest in Western Australia (WA), where more than 90% of *Eremophila* species occur (Chinnock 2007a). Many putative new taxa within *Eremophila* still require formal delimitation and description (see e.g. Brown and Buirchell 2011; Chinnock and Doley 2011; Edginton 2015; Brown and Davis 2016; Buirchell and Brown 2016). Approximately half of these putative taxa are found in WA, and many are listed under federal and state level legislation as rare or threatened (Western Australian Herbarium's FloraBase, see <https://florabase.dpaw.wa.gov.au/>, accessed 10 February 2021). Reducing uncertainty of these unresolved taxa of *Eremophila* is thus important for determining management actions to protect potentially rare and endangered species. More generally, accurate circumscription and naming of taxa is fundamental to effective conservation and sustainable management (Burgman et al. 2000; Wege et al. 2015) and to formulate policies to protect genetic and ecological diversity, understand evolutionary processes, and mitigate risks from land-use change and development (Coates et al. 2014; Oliveira et al. 2017).

*Eremophila tietkensii* was described by von Mueller and Tate (1890) from specimens collected on the 1889 expedition of William Tietkens in the Northern Territory

(NT; Chinnock 2007b). *E. tietkensis* was considered by Ewart and Jarrett (1928) to be a variety of *E. latrobei*. However, Chinnock (2007b), after examining the type and resolving some nomenclatural confusion, reinstated *E. tietkensis* at the species level. Chinnock regarded *E. tietkensis* as widespread and distributed from the Cape Range and Carnarvon biogeographic region of WA to the NT border region (Chinnock 2007b; Western Australian Herbarium's FloraBase, see <https://florabase.dpaw.wa.gov.au/>). Substantial variation in leaf morphology across its geographic range led him to suggest that recognition of subspecies may be warranted.

In 2012, three putative taxa occurring within the Pilbara bioregion and adjacent Great Sandy and Little Sandy Desert bioregions in north-west WA were segregated from *E. tietkensis* and given the phrase names *E. sp.* Hamersley Range (K. Walker KW 136), *E. sp.* Rudall River (P. G. Wilson 10512), and *E. sp.* Calvert Range (A. A. Burbidge 738; Western Australian Herbarium's FloraBase, see <https://florabase.dpaw.wa.gov.au/>; Fig. 1). Of these, *E. sp.*

Hamersley Range was listed as Priority 3 and *E. sp.* Rudall River as Priority 2 (Western Australian Herbarium's FloraBase, see <https://florabase.dpaw.wa.gov.au/>). The third phrase-named taxon, *E. sp.* Calvert Range, was not conservation-listed despite being represented by fewer specimens collected from a more limited geographic range than the other two.

*Eremophila sp.* Hamersley Range is currently known from the Hamersley subregion of the Pilbara bioregion of north-western Australia and occurs across a geographic range of ~200 km from south of Paraburdoo to north-west of Newman (Western Australian Herbarium's FloraBase, see <https://florabase.dpaw.wa.gov.au/>). This taxon was segregated on the basis of having one to four flowers per axil, with pedicels 2.5–3 times the length of the flowers, and the ovary being ribbed and with a dense covering of glandular and eglandular hairs (Western Australian Herbarium's FloraBase, see <https://florabase.dpaw.wa.gov.au/>). *Eremophila sp.* Hamersley Range has been recorded as favouring high parts of the landscape such as breakaways and upper



**Fig. 1.** (a) *Eremophila tietkensis*. Photograph: R. Fowler. (b) *E. sp.* Rudall River. Photograph: J. Hurter. (c) *E. sp.* Calvert Range. Photograph: A. Brown. (d) *E. sp.* Hamersley Range. Photograph: J. Naaykens.



hill slopes (Department of Mines and Petroleum 2016). *Eremophila* sp. Rudall River is known from across a ~400-km range from the east of the Pilbara bioregion to the Great Sandy Desert around the Rudall and Oakover Rivers (Western Australian Herbarium's FloraBase, see <https://florabase.dpaw.wa.gov.au/>). It occurs on quartzitic scree slopes (Office of the Environmental Protection Authority 2014), and is characterised by persistent leaf bases, coriaceous, ovate leaves, and up to three flowers per axil (Western Australian Herbarium's FloraBase, see <https://florabase.dpaw.wa.gov.au/>). The third putative taxon, *E. sp.* Calvert Range, is poorly known from only three specimens from an area ~100 km north-west to ~50 km south of Lake Disappointment in the Little Sandy Desert bioregion (Western Australian Herbarium's FloraBase, see <https://florabase.dpaw.wa.gov.au/>). It was segregated from the other taxa on the basis of having sericeous eglandular hairs on the ovary, the leaf surface being strumose, and having sepals fused at base. Despite its apparently limited range, *E. sp.* Calvert Range was not conservation-listed, partly owing to the remoteness of its range and the likelihood that it has been under-collected. *Eremophila tietkensis* sens. str. (i.e. not including the phrase-named entities) is morphologically variable and overlaps in geographic range and leaf morphology with each of the other taxa (Fig. 1). There has been ongoing confusion in identifications within the *E. tietkensis* complex, with some specimens (e.g. PERTH 08731535, PERTH 08957274, PERTH 06017142, PERTH 06023983, PERTH 06570496, PERTH 06653561) being reassigned from *E. tietkensis* to *E. sp.* Hamersley Range and vice versa (Western Australian Herbarium's FloraBase, see <https://florabase.dpaw.wa.gov.au/>).

Phylogenetic relationships among the taxa in the *E. tietkensis* species complex remain largely unknown. A recent molecular phylogenetic study of *Eremophila* based on chloroplast, mitochondrial and nuclear rDNA (Fowler 2018) included three samples of *E. tietkensis* and one sample of *E. sp.* Hamersley Range; this study concluded that *E. tietkensis* was non-monophyletic. However, the clade in which these samples were placed (which contained representatives from *Eremophila* sections *Eremophila*, *Eremaea* Chinnock, and *Pulchriseplae* Chinnock) is poorly resolved overall, with low support values on most nodes (Fowler 2018). Given these uncertainties, the objective of this study was to determine, using a combined morphological and molecular approach, whether *E. sp.* Hamersley Range, *E. sp.* Rudall River and *E. sp.* Calvert Range should be formalised as new taxa or synonymised under *E. tietkensis*.

## Materials and methods

### Characterisation and quantification of morphometric traits

Morphological measurements of all specimens within the *E. tietkensis* species complex held at the Western Australian

Herbarium were assessed. Preliminary examination of herbarium specimens indicated that some *E. tietkensis* specimens had been misidentified and were likely to belong, instead, to one of the putative taxa (confirmed with herbarium experts); these specimens were re-determined accordingly. Morphological characters of 67 specimens of *E. tietkensis*, 18 of *E. sp.* Hamersley Range, 14 of *E. sp.* Rudall River and 6 of *E. sp.* Calvert Range were measured or assessed quantitatively and qualitatively (Table 1).

Leaf surface trichomes were examined using compound and scanning electron microscopy (SEM) to determine their morphology and anatomy. No discernible differences were found between abaxial and adaxial trichomes; so, data from both leaf surfaces were combined. For light microscopy (LM), leaf samples of ~5 × 5 mm were taken from herbarium specimens and placed on a glass slide with sufficient water to rehydrate. The sample was left to soften for 1 min to aid in removal of trichomes, which were scraped onto the slide with forceps and examined under a compound light microscope. For SEM, leaf samples of ~2 × 2 mm were taken from herbarium specimens, coated with gold using a JEOL Smart Coater sputter coater, and imaged using a Philips 505 Scanning Electron Microscope at the Western Australian Herbarium.


Measurements of vegetative, floral and fruit characters were made by hand with a ruler to the nearest 0.5 mm. For all quantitative characters, three measurements per individual were averaged, with ratios calculated before averaging. For leaf characters, the three largest fully developed leaves on each specimen were chosen. Of the 23 characters measured or scored, only those that were assessable on most specimens were included to ensure the number of individuals retained in the final dataset was maximised. The final dataset contained seven morphological characters: L1, L4, L5, L9, L10, L11, and O3 (Table 1). Leaf length (L1) and ratio (shape) characters (L4 and L5) were included, while leaf width (L2) and distance to widest point (L3) were excluded so as to remove logical auto-correlations between the ratios and their base measurements. The same reasoning was used to include petiole width (L9) and ratio of petiole length:width (L10) while excluding petiole length (L8). The apex shapes (L6 and L7) were excluded because they were highly variable within individual on specimens and could not be adequately scored. While potentially informative, comparison of sepal size and indumentum across all specimens was problematic because sepals in members of the *E. tietkensis* complex are accrescent, enlarging significantly during and after flowering, and glabrescent. Consequently, sepal length, width and indumentum were excluded. Density of glandular and eglandular hairs on the ovary, another potentially useful character, was not able to be used because of an insufficient number of specimens with ovaries at a comparable phenological stage; indumentum density was found to vary greatly as the ovary matured. Using mostly vegetative characters meant that a higher

**Table 1.** Traits measured for morphometric ordination and classification analyses, and description of character states scored for each trait.

Character	State	Type	Code
Leaves			
Average length (mm)	Largest, mature leaves	Continuous	L1
Average width (mm)	Largest, mature leaves	Continuous	L2
Average distance from leaf axil to widest point of leaf blade (mm)	Largest, mature leaves	Continuous	L3
Average leaf length:width		Ratio	L4
Average distance from leaf axil to widest point of leaf blade:length		Ratio	L5
Apex shape	(0) acute, (1) acuminate, (2) attenuate, (3) mucronate	Multistate	L6
Apex	(0) not recurved, (1) recurved		L7
Average petiole length (mm)	Largest, mature leaves	Continuous	L8
Average petiole width (mm)	Largest, mature leaves	Continuous	L9
Average petiole length:width		Ratio	L10
Trichomes, abaxial and adaxial	(0) completely septate, (1) half septate with apical cell longer than cells in bottom half	Binary	L11
Pedicels			
Average length (mm)	Corolla present, mid-flowering	Continuous	P1
Sepals			
Average length (mm)	Corolla present, mid-flowering	Continuous	S1
Average width (mm)	Corolla present, mid-flowering	Continuous	S2
Average length (mm)	Fruiting	Continuous	S3
Average width (mm)	Fruiting		
Density of hairs on margins	(0) very sparse, (1) sparse, (2) moderate, (3) dense	Multistate	S4
Apex shape	(0) obtuse, (1) rounded, (2) acute (3) retuse	Multistate	S5
Habit			
Plant height (m)	Herbarium specimen sheet	Continuous	H1
Plant width (m)	Herbarium specimen sheet	Continuous	H2
Ovary			
Density of glandular hairs	(0) absent, (1) sparse, (2) moderate [30% surface area covered], (3) dense [>30% covered]	Multistate	O1
Density of eglandular hairs	(0) absent, (1) sparse, (2) moderate [30% surface area covered], (3) dense [>30% covered]	Multistate	O2
Sericeous yellow eglandular trichomes	(0) absent, (1) present	Binary	O3
Texture	(0) not ribbed, (1) ribbed	Binary	O4

proportion of herbarium specimens could be included in analyses; this approach also had the advantage that plant identification can be made in the absence of flowers.

The dataset was analysed using non-metric multi-dimensional scaling (nMDS), principal coordinate analysis (PCoA) ordinations, and unweighted pair-group method with



**Fig. 2.** Collection locations of all PERTH voucher specimens of *Eremophila tietkensisii*, *E. sp. Hamersley Range*, *E. sp. Rudall River* and *E. sp. Calvert Range* in WA. Specimens were assigned to *a priori* groups numbered 1–7.

**Table 2.** Total number of *Eremophila* specimens used in morphometric and molecular analyses.

Species	Morphometric analysis		Molecular analysis				
	Herb. specimens	Morphometrics	Herb. specimens	Additional material	DNA extraction	Sequenced	SNP analysis
<i>E. sp. Hamersley Range</i>	18	18	18	1	19	18	15
<i>E. tietkensisii</i>	67	63	67	6	73	54	36
<i>E. sp. Rudall River</i>	14	14	14	1	14	11	10
<i>E. sp. Calvert Range</i>	6	6	6	–	6	6	6
Total	105	101	105	8	109	86	67

Herb., herbaceous.

arithmetic mean (UPGMA) classification in PRIMER (ver. 6.1, see <https://www.primer-e.com/>; Clarke and Gorley 2006). Differences within and among recovered groups were analysed using ANOSIM, which generates an R statistic with a theoretical range from 0 (indicates a random distribution) to 1 (perfect separation of groups; Clarke and Gorley 2006).

Specimens were assigned to broad geographic groups (Fig. 2) to enable assessment of geographic structuring or partitioning within the widespread *E. tietkensisii* complex.

**Molecular analysis**

In total, 101 herbarium specimens were sampled for a ddRADSeq molecular analysis. Four specimens from the

Western Australian Herbarium collections of *E. tietkensisii* and the phrase-named taxa were excluded because of insufficient material or poor quality (Table 2). Additional non-vouchered silica-dried material of each species, except *E. sp. Calvert Range*, was provided by B. Buirchell for inclusion in the molecular analysis. As the purpose of this study was to determine species boundaries rather than produce a phylogeny or test evolutionary hypotheses or hierarchies, an out-group was not included.

Genomic DNA was extracted using the Qiagen DNEasy Plant Minikit (Qiagen, Hilden, Germany) following the manufacturer’s protocol modified in the following ways: at Step 2, 1.35 mL of Buffer AP1, in addition to 1 µL of dithiothreitol for every 1 mL of Buffer AP1, and 4 µL of RNase A was

added to each tube containing 80 mg of plant material, vortexed, then samples were incubated for 30 min at 65°C; and at Step 11, 40 µL of Buffer AE heated to 65°C was added to each spin column filter for elution and then incubated at room temperature (15–25°C) for 10 min.

DNA concentration was quantified with a Qubit dsDNA BR Assay Kit and a Qubit 3.0 Fluorometer (Invitrogen, Carlsbad, CA, USA). Fragment lengths were checked with a LabChip GX Touch 24 Nucleic Acid Analyser using the HT DNA gDNA reagents (Perkin and Elmer, Waltham, MA, USA). Of the 109 samples, 86 were of suitable quality to be used for library preparation.

Library preparation followed the procedure in [Severn-Ellis \*et al.\* \(2020\)](#). DNA extracts were diluted to 300-ng concentration per sample with nuclease-free water and digested with 5 units of HpyCH4IV restriction enzyme (New England Biotechnologies (NEB), Ipswich, MA, USA), and 5 U of HinfI restriction enzyme (NEB) per sample and 3 µL of the master mix was added to each sample. Barcoded and common adapters designed to complement the pairs of restriction enzyme overhangs were prepared as described by [Peterson \*et al.\* \(2012\)](#). Ligated fragments were cleaned of excess, unligated adapters, and fragments within the range of 250–800 base pairs were selected using Ampure XP beads (Beckman Coulter, Brea, CA, USA) and enriched by polymerase chain reaction (PCR) amplification by using Phusion Hot-Start High-Fidelity Polymerase Master Mix (Thermo Fisher Scientific, Waltham, MA, USA).

Pooled libraries were cleaned (Ampure XP beads), quantified with Qubit and visualised and quality-checked using the LabChip GX Touch 24 with HT DNA HiSens Dual Protocol Reagents. Molarity of the final library was calculated and 50–100 µL of the ddRAD bead cleaned library was diluted to a final concentration of 10–20 nM by using normalisation buffer. The pooled ddRAD libraries were sequenced (2 × 150-bp reads) across two Illumina HiSeq X lanes at the Kinghorn Centre for Clinical Genomics (KCCG) Sequencing Laboratory in Darlinghurst, New South Wales, Australia.

## Sequence data analysis

*De novo* assembly of RAD loci and single-nucleotide polymorphism (SNP) calling was performed following the bioinformatics workflow described in [Severn-Ellis \*et al.\* \(2020\)](#), see [https://github.com/ascheben/RAD\\_analysis\\_workflow/](https://github.com/ascheben/RAD_analysis_workflow/). Pooled data were de-multiplexed using Stacks (ver. 2.53, see <https://catchenlab.life.illinois.edu/stacks/>) process\_radtags ([Catchen \*et al.\* 2013](#)), with barcode rescue (-r), quality filtering (-q, -c) and RAD tag checks (-renz\_1 HpyCH4IV -renz\_2 HinfI). Low-quality reads were discarded and reads were trimmed (144 bp) with Trimmomatic (ver. 0.39, see <http://www.usadellab.org/cms/index.php?page=trimmomatic>; [Bolger \*et al.\* 2014](#)). Quality checks using FastQC (ver. 0.11.9, see <https://www.bioinformatics.babraham.ac.uk/projects/fastqc/>; [Andrews 2010](#)) were conducted, followed by a

preliminary diversity assessment based on pairwise distances using Mashtree (ver. 0.37, see <https://metacpan.org/dist/Mashtree>; [Katz \*et al.\* 2019](#)), whereafter individuals with extremely low genotyping rates were removed. *De novo* assembly of RAD loci and SNP calling was performed using Stacks (ver. 2.53; [Catchen \*et al.\* 2013](#)). For high levels of intraspecific divergence, it has been recommended that  $n = M$  in the Stacks software ([Paris \*et al.\* 2017](#)), and was considered appropriate for the present study. Stacks parameters of M-3, m-3, and n-3 were chosen following experimentation with different parameter settings ([Paris \*et al.\* 2017](#)) rendering the highest number of polymorphic loci in 80% of the populations studied.

Further filtering with VCFtools (ver. 0.1.16, see <https://vcftools.github.io/>; [Danecek \*et al.\* 2011](#)) was executed to remove indels and retain only biallelic, high-quality SNPs. The minor allele frequency (MAF) filters rare alleles, and the threshold was set at 0.05. The filters that remove the most SNPs are usually depth (-minDP) and the missingness (max-missing) filters. For heterozygous samples, read depths  $\geq 5$  have been suggested to avoid undercalling of heterozygous genotypes ([Maruki and Lynch 2017](#); [Bilton \*et al.\* 2018](#)). Depth and missingness filters were fine-tuned to find a balance between the quantity and quality of SNPs. Read depths of  $\geq 3$  (DP3) and  $\geq 5$  (DP5) were evaluated and a read depth of 5 (MAF = 0.05) was chosen. Missingness was maintained at 90% across samples (max-missing = 0.8). Finally, individuals with low genotyping rates (missingness > 50%) based on the filtered SNPs were removed.


A principal component analysis (PCA) was conducted using the gdsfmt, SNPRelate, gridExtra, and ggrepel packages in R (RStudio, Inc., Boston, MA, USA) to visualise genetic diversity using detected SNPs and to assess whether samples formed clusters congruent with the results of the morphometric analyses.

## Results

### Key morphometric differences among taxa

Examination of abaxial and adaxial leaf trichomes by using both SEM and LM showed that trichomes of *Eremophila* sp. Hamersley Range were substantially different from those of the other three taxa in the complex ([Fig. 3a](#)). Trichomes in *E. sp.* Hamersley Range were distinctly septate with six to eight cells of approximately equal length, with the terminal cell with a rounded tip ([Fig. 3a–b, 4f–i](#)). By contrast, *E. tietkensii* trichomes had four to five cells per trichome, with the terminal cell comprising half to one-third of the total length of the trichome and being attenuate at the apex ([Fig. 3c–d, 4a–e](#)). Trichomes in *E. sp.* Rudall River ([Fig. 3e–f, 4j–k](#)) and *E. sp.* Calvert Range ([Fig. 3g–h, 4l–m](#)) were similar to those of *E. tietkensii* but were slightly longer (up to 350 µm in *E. sp.* Rudall River and 300 µm in *E. sp.* Calvert Range, compared with 250 µm in *E. sp.* Hamersley Range






**Fig. 3.** Scanning electron microscopy (SEM) images of adaxial trichomes on leaves of *Eremophila* sp. Hamersley Range: (a) PERTH 03557464, (b) PERTH 09105972; *E. tietkensis*: (c) PERTH 03856275, (d) PERTH 03899918; *E. sp.* Rudall River: (e) PERTH 03878759, (f) PERTH 04201159; and *E. sp.* Calvert Range: (g) PERTH 03878570, (h) PERTH 07512821.


and *E. tietkensis*) and narrower (12–25 µm wide in *E. sp.* Rudall River and 10–22 µm wide in *E. sp.* Calvert Range, compared with 20–30 µm in *E. sp.* Hamersley Range and *E. tietkensis*), and tended to have slightly longer terminal cells.

Overall, *E. sp.* Hamersley Range was the only taxon that could be distinguished on the basis of trichomes; the remaining three taxa differed only slightly and were highly variable.





**Fig. 4.** Trichomes under compound microscope at 400× of *Eremophila tietkensis*: (a) PERTH 08316899, (b) PERTH 03881202, (c) PERTH 06752519, (d) PERTH 03851281, (e) PERTH 07324545; *E. sp.* Hamersley Range: (f) PERTH 06653537, (g) PERTH 0653561, (h) PERTH 06017142, (i) PERTH 8521; *E. sp.* Rudall River: (j) PERTH 08305447, (k) PERTH 03878740; and *E. sp.* Calvert Range: (l) PERTH 07765886, (m) PERTH 07512821.



**Fig. 5.** Ovary indumentum of (a) *Eremophila* sp. Calvert Range, showing sericeous eglandular trichomes; and (b) *E. Hamersley* Range, showing glandular and eglandular trichomes.

*Eremophila* sp. Calvert Range had a sericeous eglandular indumentum on the ovary that is distinct from that of the other three species (Fig. 5a). Ovaries in *Eremophila* sp.


Hamersley Range had a sparse to moderate covering of glandular and eglandular hairs (Fig. 5b); those in *E. sp.* Rudall River mostly lacked eglandular hairs and had a

moderate covering of glandular hairs, and, in *E. tietkensisii*, ovaries mostly lacked eglandular hairs and had a moderate to dense covering of glandular hairs.

### Morphometric groupings of taxa

Non-metric multidimensional scaling (nMDS) and UPGMA clustering showed that *Eremophila* sp. Hamersley Range and

*E. sp.* Calvert Range are morphologically distinct from *E. tietkensisii* and *E. sp.* Rudall River; however, each taxon is morphologically closer to *E. tietkensisii* than to each other (Fig. 6a). Whereas, overall, *E. tietkensisii* and *E. sp.* Rudall River formed somewhat separate groups, some *E. tietkensisii* individuals were morphologically more similar to *E. sp.* Rudall River than they were to other *E. tietkensisii* specimens. *Eremophila tietkensisii* showed high levels of morphological



**Fig. 6.** (a) Non-metric multi-dimensional scaling (nMDS) ordination of all individuals on the basis of selected traits (Table I) overlain with groupings from cluster analysis (b); and (b) UPGMA classification of all individuals on the basis of selected traits (Table I).

variability, with several individuals forming groups away from the rest of *E. tietkensisii* when an 80% similarity threshold based on the UPGMA classification was applied (Fig. 6b). *Eremophila* sp. Hamersley Range and *E. sp.* Calvert Range each clustered as one group both at the 70 and 80% similarity threshold (Fig. 6a). *Eremophila* sp. Hamersley Range and *E. sp.* Calvert Range formed distinct groups in the UPGMA classification (Fig. 6b). *Eremophila* sp. Rudall River and *E. tietkensisii* formed weakly separated groups, with several individuals of *E. tietkensisii* not being sufficiently distinguished from *E. sp.* Rudall River to join the larger *E. tietkensisii* group (Fig. 6b).

Pairwise ANOSIM tests between *Eremophila* sp. Hamersley Range and *E. sp.* Calvert Range and between *E. sp.* Calvert Range and *E. sp.* Rudall River returned an R statistic of 1, indicating a complete separation of groups on the basis of morphological characters (Table 3). Comparisons of *Eremophila tietkensisii* and *E. sp.* Hamersley Range ( $R = 0.96$ ,  $P < 0.001$ ), and *E. tietkensisii* and *E. sp.* Calvert Range ( $R = 0.88$ ,  $P < 0.001$ ), also showed high separation between groups (Table 3). The weakest separation of groups was between *E. tietkensisii* and *E. sp.* Rudall River ( $R = 0.5$ ,  $P < 0.001$ ), as observed in the ordination (Fig. 6a). When *E. tietkensisii* and *E. sp.* Rudall River were analysed separately from the rest of the complex, the strength of separation of groups increased only slightly ( $R = 0.52$ ,  $P < 0.001$ , Table 3).

When considering only *Eremophila tietkensisii* and *E. sp.* Rudall River, the *E. sp.* Rudall River group was more similar to the larger *E. tietkensisii* group at the 60% similarity threshold than five *E. tietkensisii* individuals (PERTH 08317178, PERTH 08332215, PERTH 06752519, PERTH 03975320, PERTH 08316937; Fig. 7). At the 80% similarity threshold,

two *E. sp.* Rudall River specimens were more similar to *E. tietkensisii* than to each other. Individuals from the same geographic location did not cluster together, indicating that morphological characters were not correlated with geographical location (Fig. 7). Geographic Groups 1 and 3 clustered together at the 70% similarity threshold despite Group 3 being ~1000 km east of Group 1.

The five *Eremophila tietkensisii* individuals that clustered with *E. sp.* Rudall River had unusually short and wide leaves. One specimen (PERTH 08332215) occurred on the border of the *E. sp.* Rudall River population, and in the classification and ordination based on morphology was grouped with *E. sp.* Rudall River; it may have been misidentified initially. One specimen (PERTH 03975320) was the southernmost occurring *E. tietkensisii* individual, ~625 km from the nearest *E. sp.* Rudall River populations. Two specimens (PERTH 08316937 and PERTH 08317178) had a more compact habit than did the other three. They were collected 7 years apart and located ~1 km from each other, but ~500 km away from the nearest *E. sp.* Rudall River populations. One specimen (PERTH 06752519) occurred even closer to the *E. tietkensisii* geographic Group 1, some ~585 km from the nearest *E. sp.* Rudall River populations. These four specimens are unlikely to be misidentifications of *E. sp.* Rudall River; even if some specimens were reassigned from *E. tietkensisii* to *E. sp.* Rudall River and *vice versa*, the morphological separation between these two entities was found to be weak.

No clear structure was found in the principal coordinate analysis (PCoA) performed on *Eremophila tietkensisii* only. Axis 1 (PCo1) explained 54.9% of the variation and was correlated mainly with mean leaf and petiole length. Axis 2 (PCo2) explained 19.6% of the variation and was correlated with the distance to widest point: length ratio (Fig. 8). Despite the wide geographic range of the samples, there was no clear separation into geographically distinct morphotypes.

## Molecular analysis


Dried and carefully preserved herbarium material was found to be a generally excellent source of DNA material for ddRADSeq analysis. There was no clear correlation between the age of herbarium specimen and the quality or quantity of DNA suitable for use in the molecular analysis following demultiplexing and trimming (Supplementary Fig. S1). The two oldest samples that passed filtering and quality checks and were used in the SNP analysis were collected in 1941.

A total of 1 851 822 978 sequence reads were generated, with 1 712 241 716 remaining following demultiplexing. After trimming, 1 511 865 102 reads remained. The total number of variant sites before filtering was 14 164 739. After removing samples with >90% SNPs missing, 67 of 78 individuals remained, and after filters were applied, 39 396 variant sites remained. Finally, when a single SNP from each ddRAD locus based on the distance between SNPs


**Table 3.** ANOSIM R statistic and P-values of global and pairwise tests comparing *Eremophila* taxa on the basis of the morphometric dataset using 999 permutations.

Test and taxon		R statistic	P-value
Pairwise test between taxa			
<i>E. sp.</i> Calvert Range	<i>E. tietkensisii</i>	0.88	0.001
<i>E. sp.</i> Calvert Range	<i>E. sp.</i> Hamersley Range	1	0.001
<i>E. sp.</i> Calvert Range	<i>E. sp.</i> Rudall River	1	0.001
<i>E. tietkensisii</i>	<i>E. sp.</i> Hamersley Range	0.96	0.001
<i>E. tietkensisii</i>	<i>E. sp.</i> Rudall River	0.50	0.001
<i>E. sp.</i> Hamersley Range	<i>E. sp.</i> Rudall River	1	0.001
Global test		0.78	0.001
Global test and pairwise test between taxa			
<i>E. tietkensisii</i>	<i>E. sp.</i> Rudall River	0.521	0.001





**Fig. 7.** Non-metric multi-dimensional scaling (nMDS) ordination of *Eremophila tietkensisii* and *E. sp. Rudall River* individuals with pedicels present overlain with groupings from cluster analysis. Numbers indicate geographic areas of individuals as shown in Fig. 2.




**Fig. 8.** Principal coordinate analysis (PCoA) for six traits of all *Eremophila tietkensisii* individuals. Numbers represent geographic groups as per Fig. 2.

was randomly kept and samples with over 50% missing SNPs were removed, all 67 individuals remained, with a total of 7032 SNPs.

The first two components of the PCA based on SNPs explained 46.5% of the variation (Fig. 9). The broad

geographic groupings within *Eremophila tietkensisii* were recovered on the PCA ordination, except that *E. sp. Rudall River* sat between and mixed with *E. tietkensisii* geographic Groups 2 and 3 and was not genetically divergent from them (Fig. 9). The *E. tietkensisii* specimen from Exmouth (Group 4)



**Fig. 9.** Principal component analysis of detected SNPs of *Eremophila tietkensis*, *E. sp. Hamersley Range*, *E. sp. Calvert Range*, and *E. sp. Rudall River*. Inset shows positions of *E. tietkensis* and *E. sp. Rudall River* specimens.

clustered with geographic Groups 1 and 2 and did not appear to be genetically differentiated. *Eremophila tietkensis* Group 2 clustered with Groups 1 and 3, despite Group 3 being very geographically distant. *Eremophila sp. Hamersley Range* and *E. sp. Calvert Range* were strongly genetically segregated from each other and from the rest of *E. tietkensis* and *E. sp. Rudall River*, despite their close geographical proximity to all groups (Fig. 9).

## Discussion

### Congruence between morphology and molecular patterns of variation

There is a high degree of congruence between patterns of morphological and molecular variation in the *Eremophila tietkensis* species complex. Morphological variation resolved three groups, comprising *E. sp. Hamersley Range*, *E. sp. Calvert Range* and *E. tietkensis* (including *E. sp. Rudall River*). The same three groups were strongly resolved in the SNP analysis. *Eremophila sp. Rudall River* could not be adequately segregated from *E. tietkensis* in either the morphological or molecular analysis and resolved between *E. tietkensis* geographic Groups 2 and 3 either genetically or geographically. While the morphometric analysis did not resolve geographical structuring in morphology in *E. tietkensis*, there was distinct geographical patterning between the *a priori* geographic groups on the basis of the molecular analysis (Fig. 9). The genetic patterning was not strong enough to indicate that *E. tietkensis* could be reasonably

taxonomically split on the basis of geography; the spread of molecular variation across its range was slightly but not substantially greater than for the other taxa, indicating that gene flow, although restricted, is sufficient to have prevented allopatric speciation. The lack of consistency in morphological types of *E. tietkensis* within geographic groups, in conjunction with the genetic structuring among geographic groups, suggests that differences in morphology may be the result of phenotypic plasticity. Detailed analysis of substrate, habitat and underlying geology may explain differences in phenotypic variation and drivers of species distributions, because these factors have been used to explain variation and distribution in other *Eremophila* species (Coates et al. 2014).

*Eremophila sp. Hamersley Range* was genetically highly divergent from *E. tietkensis* despite its close geographic proximity. Although this analysis was unable to estimate time since divergence our results suggest that these two taxa have been genetically isolated for a long period of time. A growing number of studies of evolutionary history and distribution patterns of Pilbara biota are showing similar patterns of divergence between species (e.g. Anderson et al. 2016). Since the formation of the Great Sandy and Gibson deserts <1 million years ago, the Pilbara region has been separated from similar landforms by extensive sand plains (Pepper et al. 2013b). Species ranges contracted and expanded throughout this time, including those of *Acacia* spp. (Ladiges et al. 2006), *Triodia* spp. (Anderson et al. 2016), reptiles (Doughty et al. 2011; Pepper et al. 2013a), beetles (Guthrie et al. 2010) and spiders (Durrant et al. 2010). Owing to the topographical heterogeneity of the

Pilbara compared with surrounding low-lying plains, localised areas within the region were likely to be more climatically stable, acting as refugia for species that could not otherwise persist in the arid conditions outside of these environments (Byrne *et al.* 2016, 2017). These processes may have contributed to the high level of genetic divergence between *E. sp.* Hamersley Range and *E. tietkensisii*.

The resolution of the *Eremophila tietkensisii* complex provided here will allow the taxa to be included in future phylogenetic studies where reconstruction of patterns and timing of divergence among lineages can be inferred, providing insights into the evolutionary history of species in north-western Australia. These results will also assist in biological surveys, because these species can now be accurately discriminated, and population studies can be undertaken. We also note here the largely untapped potential of herbarium material for ddRADSeq molecular analyses for such studies. DNA extraction from herbarium material is often considered to be a supplement to field sampling (Beck and Semple 2015). We found that the quality of DNA obtained for molecular analyses was excellent in carefully stored specimens and not strongly correlated with specimen age. This observation is also consistent with those of recent similar studies elsewhere (Bakker *et al.* 2016; Forrest *et al.* 2019; Joyce *et al.* 2021). Overall, we have demonstrated that even without additional sampling, species delimitation can be achieved in an accurate, efficient and cost-effective manner by using ddRADseq to sample the genomes of herbarium specimens.

## Taxonomy

### Taxonomic implications

The congruence between morphological and molecular patterns of variation, together with discriminatory characters, allows the recognition of *Eremophila sp.* Hamersley Range and *E. sp.* Calvert Range as morphologically and genetically distinct from *E. tietkensisii*. These taxa occur in close geographic proximity and are likely to have the opportunity to interbreed. Their clear genetic and morphological separation indicates that they do not do so. This, in turn, indicates that they are likely to comprise independently evolving metapopulation lineages without significant gene exchange. Therefore, we believe that they should be recognised as distinct species under the unified species concept (de Queiroz 2007). By contrast, the similar morphology and lack of clear genetic separation between *E. tietkensisii* and *E. sp.* Rudall River indicates that the latter cannot be recognised at species rank and should be absorbed back into *E. tietkensisii*.

Accordingly, below we formally name and describe as new species *Eremophila naaykensisii* A.L.Curtis & K.R.Thiele

and *E. hurteri* A.L.Curtis & K.R.Thiele, and circumscribe *E. tietkensisii* to include specimens currently phrase-named as *E. sp.* Rudall River.

### Key to species of *Eremophila* section *Eremophila* (amended from Chinnock 2007a)

1. Leaves linear to linear-oblongate or lanceolate.....2  
Leaves ovate to obovate or oblanceolate.....8
2. Sepals separated at base.....115. *E. oppositifolia*  
Sepals imbricate at base.....3
3. Outside surface of sepals glabrous; branches and leaves green, pseudoglabrous, with hairs usually completely obscured by resin.....117. *E. cryptothrix*  
Outside surface of sepals pubescent; branches and leaves grey to grey-green, hairs obvious.....4
4. Sepals <6.5 mm long; corolla 5.5–7.5 mm long (Qld).....  
.....114. *E. arbuscula*  
Sepals and corolla >7 mm long (WA).....5
5. Leaves linear-oblongate, channelled, <3 mm wide, very prominently tuberculate.....122. *E. mirabilis*  
Leaves lanceolate to oblanceolate, flattened, >7.5 mm wide or if narrower, not prominently tuberculate.....6
6. Flowers 1–4 per axil; corolla lilac, white, yellow, pale blue, pink or mauve, anthers included.....6a  
Flowers 1 per axil; corolla cream or pink; anthers usually extending beyond throat.....7
- 6a. Leaf indumentum comprising simple, uniseriate hairs, the terminal cell much longer than the others and usually attenuate.....6b  
Leaf indumentum comprising simple, uniseriate hairs that are evenly septate, the terminal cell no longer than the others and with a bluntly rounded tip.....*E. naaykensisii*
- 6b. Ovary densely glandular-puberulous with scattered or numerous longer eglandular hairs, style glabrous or with a few scattered, simple, spreading, short eglandular hairs.....*E. tietkensisii*  
Ovary densely sericeous with yellow, simple, eglandular hairs; style with sparse, long spreading, eglandular hairs for most of its length.....*E. hurteri*
7. Hairs on vegetative parts matted, posterior sepal oblong to oblanceolate, broadly acute.....124. *E. macmilliana*  
Hairs on vegetative parts not matted, posterior sepal similar to anterior pair, widely ovate to suborbicular or sepals subequal, oblanceolate, obtuse.....123. *E. platycalyx*
8. Corolla white or cream sometimes tinged bluish-green on lobes.....  
.....9  
Corolla pale blue, blue, pale lilac, violet, pale mauve, pink or white-tinged lilac.....11
9. Fruit glabrous; corolla white.....116 *E. reticulata*  
Fruit pubescent; corolla cream sometimes tinged bluish-green.....  
.....10
10. Leaves very widely depressed, ovate, obtuse, very rigid; ovary/fruit with eglandular hairs.....120. *E. rigida*  
Leaves obovate, acute, flexible; ovary/fruit with eglandular hairs.....123. *E. platycalyx*
11. Leaves subopposite to opposite, widely depressed ovate, flabellate or spatulate (SA; NT).....118. *E. rotundifolia*  
Leaves alternate or irregularly opposite, ovate, spatulate or oblanceolate (WA).....12
12. Branches sulcate; sepals unequal, outer 3 broader than inner pair, oblanceolate to obovate; flowers 1, rarely 2, per axil.....  
.....119. *E. spatulata*  
Branches non-sulcate; sepals subequal, elliptic to oblanceolate; flowers 2–4 per axil.....121. *E. tietkensisii*



***Eremophila tietkensis* F.Muell & Tate, *Trans. Proc. & Rep. Roy. Soc. S. Australia* 8: 109 (1890)**

*Type*: Laura Vale, Northern Territory, [June] 1889, W. H. Tietkens s.n. (holo: MEL 82820).

*Eremophila latrobei* var. *tietkensis* (F.Muell. & Tate) Ewart & P.H. Jarrett [see <https://id.biodiversity.org.au/name/apni/114129/api/apni-format>], *Proc. Roy. Soc. Victoria* 40: 87 (1928) [see <https://id.biodiversity.org.au/instance/apni/548054>].

*Eremophila pachomai* Paczkowska & A.R.Chapman, *W. Austral. Fl. Descr. Cat.* 339 (2000), *nom. inval.* [manuscript name; no Latin description or diagnosis provided or referenced]

*Eremophila* sp. Rudall River (P. G. Wilson 10512), Western Australian Herbarium: L. J. Biggs & C. M. Parker, *Nuytsia* 23:504 (2013).

Rounded to flat-topped shrub 0.6–2(–3) m tall, aromatic. Young stems covered in a persistent, fine, grey to yellowish, appressed tomentum of simple hairs, obscurely tuberculate beneath the indumentum; older stems with grey to very pale grey, slightly fissured bark, at first with prominently raised and knob-like persistent leaf bases. Leaves scattered, pale greyish-green or grey, petiolate; petioles (2.5–)5–10(–16) mm long; lamina ovate to lanceolate, (19–)32–57(–91) × (4–)10–21(–32) mm, smooth; indumentum dense, very short, appressed, white to grey, velutinous, often matted-resinous, comprising simple, uniseriate hairs, the terminal cell much longer than the others and usually attenuate; margins entire; apex acute, attenuate, or mucronate. Flowers 2–4 per axil, pedicellate; pedicels (5–)10–14(–25) mm long, with indumentum as for stems. Sepals 5, imbricate, subequal, elliptic to oblanceolate, broadly acute to obtuse with a mucro, (7–)10–13(–18) × (1.5–)3–5(–7) mm, pinkish-purple to mauve, maroon or red, pubescent with ± appressed, tangled hairs, the margins more densely so, enlarging after flowering and then glabrescent and with prominent veins. Corolla 22–28 mm long, pale blue, blue, pale lilac to pale mauve, white tinged lilac, mauve or pink; outer surface of lobes and tube with scattered eglandular hairs particularly near the margins, often almost glabrous; mid-inner tube with moderate density of eglandular hairs. Stamens 4, included; filaments with long eglandular hairs towards base, glabrous above; anthers glabrous. Ovary ovoid-oblong, densely glandular-puberulous with scattered or numerous longer eglandular hairs; style glabrous or with a few scattered, simple, spreading, short eglandular hairs. Fruit dry, woody, ovoid-conical, ± beaked, ribbed, 6–7 × 3–4.5 mm; exocarp adhering to endocarp, glandular-puberulous but usually with some longer eglandular hairs, occasionally resinous; endocarp vertically ribbed, splitting into four segments towards apex.

## Distribution and habitat

Occurs from Exmouth on the western coast of WA in the Carnarvon IBRA bioregion (Thackway and Cresswell 1995) to just over the NT border in the east, throughout the

Pilbara, Gascoyne, Little Sandy Desert, Great Sandy Desert, Gibson Desert and Central Ranges IBRA bioregions, and down to the Murchison bioregion to the south. Occurs on a range of substrates and landscape positions, including red–brown sand, silty loam, skeletal loam over ironstone, rocky quartz, gravel, laterite, dolerite, and limestone on flats, undulating plains, saline clay plains, plateaus, gully slopes, valley floors, creeklines, scree slopes, and outcrops (Western Australian Herbarium's FloraBase, see <https://florabase.dpaw.wa.gov.au/>).

## Phenology

Flowers in late winter to at least mid-spring, with fruits maturing from early spring onward.

## Conservation status

*Eremophila tietkensis* is widespread in WA, including in several national parks and nature reserves, and is not considered to be under threat. It is known in the NT only from regions close to the WA border between Lakes Mackay and Neale.

## Notes

*Eremophila tietkensis* is a widespread and morphologically variable species. It differs from *E. naaykensis* in having shorter pedicels (shorter than or similar in length to the flowers cf. usually longer than the flowers) among less dense leaf clusters at branch apices, and having leaf indumentum trichomes with an elongated terminal cell, and from *E. hurteri* in its glandular-puberulous cf. densely silky ovary indumentum.

Plants from the vicinity of the Rudall and Oakover Rivers have leaves that are generally shorter and more ovate than is typical, and these populations were previously segregated as *Eremophila* sp. Rudall River. However, the differences in leaf shape are continuous and highly variable. Some plants distant from the Rudall-Oakover area have equally small leaves, while some plants from within that area have longer, more lanceolate leaves. Two disjunct, far-western collections from the Cape Range are typical for the species.

## Selected specimens examined

**WESTERN AUSTRALIA.** 17 km on Exmouth Road, Exmouth, *Anonymous* D 11305 (PERTH 03856208); 25.4 km NW of Cobra on the Gifford Creek Road, Upper Gascoyne, R. J. Chinnock 6888 (PERTH 08316937); The Gap, 1.4 km N of the turnoff to Christmas Pool, Paterson Range, R. J. Chinnock 6965 (PERTH 08317186); Mu Hills, Ngaanyatjaraku, R. J. Chinnock 8002 (PERTH 08669945); 12.5 km N of Towrana, R. J. Chinnock 8002 (PERTH 08316945); 21.4 km N of Gascoyne Junction, R. J. Chinnock 3796 (PERTH 08316899); Yalhall Creek near Mount Rica, Hamersley Range, Ashburton, C. A. Gardner 6420 (PERTH 03856275); 20 km WSW Parngurr, Little Sandy Desert, P. K. Latz 17817 (PERTH 08305447); Rudall River district, ~ 500 km S of Broome, P. G. Wilson 10512 (PERTH 03878740).

## *Eremophila naaykensii* A.L. Curtis & K. R. Thiele, sp. nov.

*Eremophila* sp. Hamersley Range (K. Walker KW 136) Western Australian Herbarium: L. J. Biggs & C. M. Parker, *Nuytsia* 23: 504 (2013).

**Type:** Hamersley Range (specifically Hancock Range) within mining tenement E-47/1329-I neighbouring Mining Area C, within Juna Downs Street, ~103 km WNW of Newman townsite, 5 km E of Great Northern Highway, Western Australia, 21 Feb. 2018, *C. van den Bergh* CV Opp 18 (holo: PERTH 09105972!).

Rounded to obconical shrubs or small trees 1–2.5(–3.5) m tall, aromatic. Young *stems* clothed in a persistent, fine, grey to yellowish, appressed tomentum of simple hairs, obscurely tuberculate beneath the indumentum; older stems with grey to very pale grey, slightly fissured bark, at first with prominently raised and knob-like persistent leaf bases. *Leaves* scattered but tending to be clustered towards the stem apices, pale green or grey-blue, petiolate; petioles (6–)9–13(–18) mm long; lamina lanceolate, (37–)55–71.5(–89) × (5–)7.5–12(–15) mm, smooth; indumentum dense, very short, appressed, white to grey, velutinous, often matted-resinous, comprising simple, uniseriate hairs that are evenly septate, the terminal cell no longer than the others and with a bluntly rounded tip; margins entire; apex attenuate. *Flowers* (1)2–4 per axil, appearing clustered in the dense, terminal leaf clusters, pedicellate; pedicels (20–)28–33.5(–40) mm long and ± sigmoidal, with indumentum as for stems. *Sepals* 5, imbricate, subequal, elliptic to oblanceolate, broadly acute to obtuse, sometimes mucronate, (7–)8–10(–14) mm × (2.5–)3–5(–6), yellowish, greenish, red or purple-tinged in flower (likely to be colouring further after anthesis), pubescent with ± appressed, tangled hairs, the margins more densely so, enlarging after flowering and then glabrescent and with prominent veins. *Corolla* 20–28 mm long, cream, pale blue, lilac, yellow, pink or purple sometimes with spots on upper lobe, the throat and inside of tube pale yellow to cream; outer surface of lobes and tube with scattered eglandular hairs particularly near the margins, often almost glabrous; mid-inner tube with moderately dense eglandular hairs. *Stamens* 4, included; filaments with long eglandular hairs towards base, glabrous above; anthers glabrous. *Ovary* sparsely to moderate pubescent with glandular and eglandular hairs, ribbed; style with short, patent, eglandular hairs for most of its length. Mature *fruits* not seen.

### Distribution and habitat

Endemic in the Pilbara IBRA bioregion (Thackway and Cresswell 1995). Current records indicate a geographic range of ~200 km from west to east in the southern half of the central to eastern portions of the Hamersley Ranges, occurring from the vicinity of Paraburdoo east to north-west of Newman (Western Australian Herbarium's FloraBase, see

<https://florabase.dpaw.wa.gov.au/>). Generally found in rocky ranges of the Hamersley Plateau, often high in the landscape on the tops of ironstone ranges, breakaways and on upper slopes, often in and around rocky gullies and gorges, associated with low open *Eucalyptus leucophloia* and *Corymbia ferritcola* woodlands with mixed *Acacia aneura* sens. lat. and *Acacia* spp. open shrublands and tall shrublands.

### Phenology

Flowers in late winter to at least mid-spring, often seasonally dependent, with fruits maturing from early spring onward.

### Conservation status

*Eremophila naaykensii* is currently known from six populations and is listed as a Priority Three species under the Conservation Codes for Western Australian flora, under the name *E. sp.* Hamersley Range (K. Walker KW 136; Western Australian Herbarium's FloraBase, see <https://florabase.dpaw.wa.gov.au/>).

### Etymology

Named in honour of Jeremy Naaykens, Senior Advisor Riparian Ecology and Botany at Rio Tinto Australia. Jeremy has contributed much to our knowledge of the flora of the Pilbara region, and has collected specimens from most known populations of *Eremophila naaykensii*. His enthusiasm for the species often led to his disappearance up rocky ravines and gorges to collect specimens when other more pressing work was required.

### Notes

*Eremophila naaykensii* was previously included in *E. tietkensii*, from which it can be distinguished by the presence of evenly septate hairs with rounded tips on the adaxial and abaxial leaf blades, leaves that are densely clustered at the stem apices (not terminally clustered in *E. tietkensii*), and pedicels longer than the flowers (generally the same length as the flower in *E. tietkensii*). It almost certainly belongs in the clade of *Eremophila* that contains sections *Eremaeae*, *Pulchrisepalae*, *Eremophila* and *Eriocalyx* Benth. (Fowler 2018). However, phylogenetic relationships within this clade are poorly resolved with low support, and the precise phylogenetic relationships of *E. naaykensii* are currently unknown.

### Selected specimens examined

**WESTERN AUSTRALIA.** [precise localities withheld for conservation reasons] *J. Bull* & *J. Waters* ONS PH 62.04 (PERTH 09126120); *S. Reiffer* & *H. Ajduk* WPT 1-TS (PERTH 08772088); *S. van Leeuwen* 3723 (PERTH 06023983); *S. van Leeuwen* 3828 (PERTH 06110134); *S. van Leeuwen* 4074 (PERTH 06017339); *M. E. Trudgen* MET 17478 (PERTH 06653561).

## *Eremophila hurteri* A.L.Curtis & K.R.Thiele sp. nov.

**Type:** base of Calvert Range (campsite), Calvert Range, WA, 7 August 2000, A. A. Burbidge 738 (*holo*: PERTH 07512821!).

*Eremophila* sp. Calvert Range (A. A. Burbidge 738) Western Australian Herbarium: L. J. Biggs & C. M. Parker, *Nuytsia* 23: 504 (2013).

Intricate flat-topped shrubs 1–1.5 m tall, aromatic. Young *stems* with indumentum of short, woolly, usually yellowish, sometimes grey, hairs, sometimes appearing sericeous, obscurely tuberculate beneath the indumentum; older stems grey to dark brown, scarcely fissured, often distinctly tuberculate, at first with prominently raised and knob-like persistent leaf bases. *Leaves* scattered, silvery, petiolate; petioles (7–)8–10(–11) mm long, decurrent; lamina lanceolate, (45–)50–69.5(–84.5) × (9–)9.5–12.5(–14) mm, finely strumose; indumentum dense, very short, white to grey, woolly, often matted-resinous, comprising simple, uniseriate hairs, the terminal cell much longer than the others and attenuate; margins entire; apex attenuate. *Flowers* 1 or 2 per axil, pedicellate; pedicels (4.5–)9–13(–15) mm long, straight to curved, with indumentum as for stems. *Sepals* 5, imbricate, subequal, elliptic to oblanceolate, broadly acute to obtuse, sometimes mucronulate, 7–9 × 2–3.5 mm in flower, yellow in bud, turning white or pink or mauve at anthesis, densely short-tomentose with ± silky hairs, enlarging after flowering and then glabrescent and with prominent veins. *Corolla* 20–28 mm long, white to pale purple or mauve; outer surface of lobes and tube with scattered eglandular hairs particularly near the margins, often almost glabrous; mid-inner tube lanate with eglandular hairs. *Stamens* 4, included; filaments with woolly eglandular hairs towards base, glabrous above; anthers glabrous. *Ovary* densely sericeous with yellow, simple, eglandular hairs; style with sparse, long spreading, eglandular hairs for most of its length. *Mature fruits* not seen.

### Distribution and habitat

Endemic in the Little Sandy Desert IBRA bioregion (Thackway and Cresswell 1995). Current records indicate a geographic range of ~220 km from north to south either side of Lake Disappointment (Western Australian Herbarium's FloraBase, see <https://florabase.dpaw.wa.gov.au/>). Occurs on sandstone ranges, rocky scree slopes and stony plains at the bases of low ranges.

### Phenology

Flowers in late winter to at least mid-spring, with fruits maturing from early spring onward.

### Conservation status

*Eremophila hurteri* is currently known from six populations. It is not currently listed under the Conservation Codes for

Western Australian flora (Western Australian Herbarium's FloraBase, see <https://florabase.dpaw.wa.gov.au/>).

### Etymology

Named in honour of Johan Hurter, ecologist and botanist at EcoRex Environmental Consulting and previously the Rio Tinto Identification Botanist at the Western Australian Herbarium. Johan first suggested that there may be multiple species within *Eremophila tietkensii*, and segregated *E. hurteri* (as *E. sp.* Calvert Range), *E. naaykensii* (as *E. sp.* Hamersley Range) and *E. sp.* Rudall River.

### Notes

*Eremophila* sp. Calvert Range was previously included in *E. tietkensii*, from which it can be distinguished by an indumentum of yellow, sericeous, simple, eglandular hairs on the ovary (simple eglandular and glandular hairs in *E. tietkensii*) and by the strumose leaf surfaces (not strumose in *E. tietkensii*). It almost certainly belongs in the clade of *Eremophila* that contains sections *Eremaeae*, *Pulchrisepalae*, *Eremophila* and *Eriocalyx* (Fowler 2018). However, phylogenetic relationships within this clade are poorly resolved with low support, and the precise phylogenetic relationships of *E. hurteri* are currently unknown.

### Other specimens examined

**WESTERN AUSTRALIA.** At base of Durba Hills, Wiluna, A. A. Burbidge 733 (PERTH 07765886); Rudall River Region, East Pilbara, R. P. Hart 571 (PERTH 01226991); 4.5 km Sth Parngurr, Little Sandy Desert, P. K. Latz 17825 (PERTH 08305382); 28 Aug. 2004, W. P. Muir WPM 1046 (PERTH 08609942); 40 km S of Rudall River, ~500 km S of Broome, East Pilbara, P. G. Wilson 10540 (PERTH 03878570).

### Supplementary material

Supplementary material is available [online](#).

### References

- Anderson BM, Barrett MD, Krauss SL, Thiele K (2016) Untangling a species complex of arid zone grasses (*Triodia*) reveals patterns congruent with co-occurring animals. *Molecular Phylogenetics and Evolution* **101**, 142–162. doi:10.1016/j.ympev.2016.05.014
- Andrews S (2010) FastQC: a quality control tool for high throughput sequence data. Available at <https://www.bioinformatics.babraham.ac.uk/projects/fastqc/> [Verified 15 May 2020]
- Bakker FT, Lei D, Yu J, Mohammadin S, Wei Z, van de Kerke S, Gravendeel B, Nieuwenhuis M, Staats M, Alquezar-Planas DE, Holmer R (2016) Herbarium genomics: Plastome sequence assembly from a range of herbarium specimens using an Iterative Organelle Genome Assembly pipeline. *Biological Journal of the Linnean Society. Linnean Society of London* **117**, 33–43. doi:10.1111/bj.12642
- Beck JB, Semple JC (2015) Next-generation sampling: pairing genomics with herbarium specimens provides species-level signal in Solidago (Asteraceae). *Applications in Plant Sciences* **3**, 1500014. doi:10.3732/app.1500014
- Bilton TP, McEwan JC, Clarke SM, Brauning R, van Stijn TC, Rowe SJ, Dodds KG (2018) Linkage disequilibrium estimation in low coverage



- high-throughput sequencing data. *Genetics* **209**, 389–400. doi:10.1534/genetics.118.300831
- Bolger AM, Lohse M, Usadel B (2014) Trimmomatic: a flexible trimmer for Illumina sequence data. *Bioinformatics* **30**, 2114–2120. doi:10.1093/bioinformatics/btu170
- Brown A, Buirchell B (2011) 'A field guide to the Eremophilas of Western Australia.' (Simon Nevill Publications: York, WA, Australia)
- Brown AP, Davis RW (2016) *Eremophila buirchellii* and *E. calcicola* (Scrophulariaceae), two new species from Western Australia. *Nuytsia* **27**, 211–216.
- Buirchell BJ, Brown AP (2016) New species of *Eremophila* (Scrophulariaceae): thirteen geographically restricted species from Western Australia. *Nuytsia* **27**, 253–283.
- Burgman M, Maslin BR, Andrewartha D, Keatley MR, Boek C, McCarthy M (2000) Inferring threat from scientific collections: power tests and an application to Western Australian *Acacia* Species. In 'Quantitative Methods for Conservation Biology'. (Eds S Ferson, M Burgman) pp. 7–26. (Springer New York: New York, NY, USA)
- Byrne M, Coates DJ, Macdonald BM, Hankinson M, McArthur SM, Van Leeuwen S (2016) High nuclear genetic differentiation, but low chloroplast diversity in a rare species, *Aluta quadrata* (Myrtaceae), with a disjunct distribution in the Pilbara, Western Australia. *Australian Journal of Botany* **64**, 687–695. doi:10.1071/BT16128
- Byrne M, Millar MA, Coates DJ, Macdonald BM, McArthur SM, Zhou M, van Leeuwen S (2017) Refining expectations for environmental characteristics of refugia: two ranges of differing elevation and topographical complexity are mesic refugia in an arid landscape. *Journal of Biogeography* **44**, 2539–2550. doi:10.1111/jbi.13057
- Catchen J, Hohenlohe PA, Bassham S, Amores A, Cresko WA (2013) Stacks: an analysis tool set for population genomics. *Molecular Ecology* **22**, 3124–3140. doi:10.1111/mec.12354
- Chinnock RJ (2007a) 'Eremophila and allied genera: a monograph of the plant family Myoporaceae.' (Rosenberg Pub Pty Ltd: Sydney, NSW, Australia)
- Chinnock RJ (2007b) *Eremophila tietkensii* F.Muell. & Tate (Myoporaceae), a misinterpreted species. *Journal of the Adelaide Botanic Gardens* **21**, 1–3.
- Chinnock RJ, Doley AB (2011) *Eremophila koobabbiensis* (Scrophulariaceae), a new, rare species from the wheatbelt of Western Australia. *Nuytsia* **21**, 158–161.
- Clarke KR, Gorley RN (2006) 'PRIMER v6: user manual/tutorial.' (Plymouth Marine Laboratory: Plymouth, UK)
- Coates DJ, Llorens TM, Byrne M, McArthur S, Macdonald B (2014) Disjunct, highly divergent genetic lineages within two rare *Eremophila* (Scrophulariaceae: Myoporeae) species in a biodiversity hotspot: implications for taxonomy and conservation. *Botanical Journal of the Linnean Society* **177**, 96–111. doi:10.1111/boj.12228
- Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, Handsaker RE, Lunter G, Marth GT, Sherry ST (2011) The variant call format and VCFtools. *Bioinformatics* **27**, 2156–2158. doi:10.1093/bioinformatics/btr330
- de Queiroz K (2007) Species concepts and species delimitation. *Systematic Biology* **56**, 879–886. doi:10.1080/10635150701701083
- Department of Mines and Petroleum (2016) Clearing permit decision report. Government of Western Australia. Available at [ftp://ftp.dwer.wa.gov.au/permit/4615/4615-6%20Decision%20Report.pdf](http://ftp.dwer.wa.gov.au/permit/4615/4615-6%20Decision%20Report.pdf) [Verified 15 May 2020]
- Doughty P, Rolfe JK, Burbidge AH, Pearson DJ, Kendrick PG (2011) Herpetological assemblages of the Pilbara biogeographic region, Western Australia: ecological associations, biogeographic patterns and conservation. *Records of the Western Australian Museum* **78**(Suppl.), 315–341. doi:10.18195/issn.0313-122x.78(2).2011.315-341
- Durrant BJ, Harvey M, Framenau V, Ott R, Waldoek JM (2010) Patterns in the composition of ground-dwelling spider communities in the Pilbara bioregion, Western Australia. *Records of the Western Australian Museum* **78**(Suppl.), 185–204. doi:10.18195/issn.0313-122x.78(1).2010.185-204
- Edginton MA (2015) *Eremophila woodiae* Edginton (Scrophulariaceae), a new species from Queensland. *Austrobaileya* **9**, 408–415.
- Ewart AJ, Jarrett PH (1928) Contributions to the Flora of Australia, No. 34. Additions to the Flora of the Northern Territory and locality records. *Proceedings of the Royal Society of Victoria* **40**, 81–87.
- Forrest LL, Hart ML, Hughes M, Wilson HP, Chung KF, Tseng YH, Kidner CA (2019) The limits of Hyb-Seq for herbarium specimens: impact of preservation techniques. *Frontiers in Ecology and Evolution* **7**, 439. doi:10.3389/fevo.2019.00439
- Fowler RM (2018) Phylogeny of *Eremophila* and tribe Myoporeae (Scrophulariaceae). PhD thesis, The University of Melbourne, Melbourne, Vic., Australia.
- Guthrie NA, Weir T, Will K (2010) Localised and regional patterns in ground-dwelling beetle assemblages in a semi-tropical arid zone environment. *Records of the Western Australian Museum* **78**, 169–184. doi:10.18195/issn.0313-122x.78(1).2010.169-184
- Joyce EM, Pannell CM, Rossetto M, Yap JYS, Thiele KR, Wilson PD, Crayn DM (2021) Molecular phylogeography reveals two geographically and temporally separated floristic exchange tracks between Southeast Asia and northern Australia. *Journal of Biogeography* **48**(5), 1213–1227. doi:10.1111/jbi.14072
- Katz LS, Griswold T, Morrison S, Caravas J, Zhang S, den Bakker HC, Deng X, Carleton HA (2019) Mashtree: a rapid comparison of whole genome sequence files. *Journal of Open Source Software* **4**, 1762. doi:10.21105/joss.01762
- Ladiges PY, Ariati SR, Murphy DJ (2006) Biogeography of the *Acacia victoriae*, *pyrifolia* and *murrayana* species groups in arid Australia. *Journal of Arid Environments* **66**, 462–476. doi:10.1016/j.jaridenv.2006.01.012
- Maruki T, Lynch M (2017) Genotype calling from population-genomic sequencing data. *G3: Genes, Genomes, Genetics* **7**, 1393–1404. doi:10.1534/g3.117.039008
- Office of the Environmental Protection Authority (2014) Kintyre Uranium project, Government of Western Australia. Available at [https://www.epa.wa.gov.au/sites/default/files/EPA\\_Report/Rep%201522%20Kintyre%20PER%20280714.pdf](https://www.epa.wa.gov.au/sites/default/files/EPA_Report/Rep%201522%20Kintyre%20PER%20280714.pdf) [Verified 15 August 2021]
- Oliveira U, Soares-Filho BS, Paglia AP, Brescovit AD, De Carvalho CJB, Silva DP, Rezende DT, Leite FSF, Batista JAN, Barbosa JPPP, Stehmann JR, Ascher JS, De Vasconcelos MF, De Marco P, Löwenberg-Neto P, Ferro VG, Santos AJ (2017) Biodiversity conservation gaps in the Brazilian protected areas. *Scientific Reports* **7**, 9141. doi:10.1038/s41598-017-08707-2
- Paris JR, Stevens JR, Catchen JM (2017) Lost in parameter space: a road map for STACKS. *Methods in Ecology and Evolution* **8**, 1360–1373. doi:10.1111/2041-210X.12775
- Pepper M, Doughty P, Fujita MK, Moritz C, Keogh JS (2013a) Speciation on the rocks: integrated systematics of the *Heteronotia spelea* species complex (Gekkota; Reptilia) from western and central Australia. *PLoS One* **8**, e78110. doi:10.1371/journal.pone.0078110
- Pepper M, Doughty P, Keogh JS (2013b) Biodiversity and endemism in the iconic Australian Pilbara region: a review of landscape evolution and biotic response in an ancient refugium. *Journal of Biogeography* **40**, 1225–1239. doi:10.1111/jbi.12080
- Peterson BK, Weber JN, Kay EH, Fisher HS, Hoekstra HE (2012) Double digest RADseq: an inexpensive method for *de novo* SNP discovery and genotyping in model and non-model species. *PLoS One* **7**, e37135. doi:10.1371/journal.pone.0037135
- Severn-Ellis AA, Scheben A, Neik TX, Saad NSM, Pradhan A, Batley J (2020) Genotyping for species identification and diversity assessment using double-digest restriction site-associated DNA Sequencing (ddRAD-Seq). In 'Legume genomics'. (Eds M Jain, R Garg) pp. 159–187. (Springer New York: New York, NY, USA)
- Thackway R, Cresswell ID (1995) An interim biogeographic regionalisation for Australia: a framework for setting priorities in the National Reserves System Cooperative Program. (Australian Nature Conservation Agency, Reserve Systems Unit: Canberra, ACT, Australia). Available at <https://www.awe.gov.au/sites/default/files/documents/ibra-framework-setting-priorities-nrs-cooperative-program.pdf>
- von Mueller FJH, Tate R (1890) List of Plants collected during Mr Tietkens' expedition into Central Australia, 1889. *Transactions, Proceedings and Reports of the Royal Society of South Australia* **13**, 94–109.
- Wege J, Thiele K, Shepherd K, Butcher R, Macfarlane T, Coates D (2015) Strategic taxonomy in a biodiverse landscape: a novel approach to maximizing conservation outcomes for rare and poorly known flora. *Biodiversity and Conservation* **24**, 17–32. doi:10.1007/s10531-014-0785-4

**Data availability.** The data that support this study are available in GenBank at <https://www.ncbi.nlm.nih.gov/bioproject/765188>.

**Conflicts of interest.** Kevin R. Thiele is an Associate Editor for *Australian Systematic Botany*. Despite this relationship, he did not at any stage have Associate Editor-level access to this manuscript while in peer review, as is the standard practice when handling manuscripts submitted by an editor to this journal. *Australian Systematic Botany* encourages its editors to publish in the journal and they are kept totally separate from the decision-making process for their manuscripts. The authors declare that they have no further conflicts of interest.

**Declaration of funding.** This project was funded in part by Rio Tinto.

**Acknowledgements.** We thank the Curator and staff at the Western Australian Herbarium for access to equipment and specimens and for providing material for both morphometric and molecular work, particularly Steve Dillon for his careful initial observations on trichome morphology in this group, and for assistance with SEM. The Batley Laboratory at the University of Western Australia provided access to resources for the molecular work. Thanks go to Bevan Buirchell for providing additional field samples, and Johan Hurter for photographs and for the initial groundwork that led to this project. Finally, we thank the reviewers whose comments improved an earlier version of this paper.

**Author affiliations**

<sup>A</sup>Western Australian Herbarium, Biodiversity and Conservation Science, Department of Biodiversity, Conservation and Attractions, Locked Bag 104, Bentley Delivery Centre, WA 6983, Australia.

<sup>B</sup>School of Biological Sciences, Faculty of Science, The University of Western Australia, 35 Stirling Highway, Crawley, WA 6009, Australia.

<sup>C</sup>Rio Tinto Iron Ore, 152–158 St Georges Terrace, Perth, WA 6000, Australia.

<sup>D</sup>School of BioSciences, The University of Melbourne, Parkville, Vic. 3010, Australia.