



Improving the Phenotypic Properties of the *Ceratitis capitata* (Diptera: Tephritidae) Temperature-Sensitive Lethal Genetic Sexing Strain in Support of Sterile Insect Technique Applications

Authors: Porras, Mitzy F., Meza, Jose S., Rajotte, Edwin G., Bourtzis, Kostas, and Cáceres, Carlos

Source: Journal of Economic Entomology, 113(6) : 2688-2694

Published By: Entomological Society of America

URL: <https://doi.org/10.1093/jee/toaa220>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Improving the Phenotypic Properties of the *Ceratitis capitata* (Diptera: Tephritidae) Temperature-Sensitive Lethal Genetic Sexing Strain in Support of Sterile Insect Technique Applications

Mitzy F. Porras,^{1,3} Jose S. Meza,^{1,4} Edwin G. Rajotte,² Kostas Bourtzis,¹ and Carlos Cáceres^{1,5}

¹Insect Pest Control Laboratory, Joint FAO/IAEA Division of Nuclear Techniques in Food and Agriculture, International Atomic Energy Agency, A-2444 Seibersdorf, Austria, ²Department of Entomology, The Pennsylvania State University, 501 ASI Building, University Park, PA 16802, ³Present address: Department of Entomology, The Pennsylvania State University, 501 ASI Building, University Park, PA 16802, ⁴Present address: Programa Moscafrut, AGRICULTURA/SENESICA-IICA, Metapa de Domínguez, 30860 Chiapas, México, and ⁵Corresponding author, e-mail: c.e.caceres-barrios@iaea.org

Subject Editor: Kent Shelby

Received 11 June 2020; Editorial decision 19 August 2020

Abstract

The genetic sexing strain (GSS) of the Mediterranean fruit fly (*Ceratitis capitata* (Wiedemann)) Vienna 8^{D53} is based on a male-linked translocation system and uses two selectable markers for male-only production, the *white pupae* (*wp*) and the *temperature sensitivity lethal* (*tsl*) genes. In this GSS, males emerge from brown pupae and are resistant to high temperatures while females emerge from white pupae, are sensitive to high temperatures. However, double homozygous females (*wp tsl/wp tsl*) exhibit a slower development rate compared to heterozygous males (*wp⁺ tsl/wp tsl*) during the larval stage, which was attributed to the pleiotropic effects of the *tsl* gene. We present the first evidence that this slower development is due to a different gene, here namely *slow development* (*sd*), which is closely linked to the *tsl* gene. Taking advantage of recombination phenomena between the two loci, we report the isolation of a novel temperature sensitivity lethal strain using the *wp* mutation as a morphological marker, which showed faster development (*wp tsl* FD) during the larval stage and increased in its temperature sensitivity compared with the normal *tsl* strain. Moreover, the introgression of this novel *wp tsl* FD combined trait into the Vienna 8^{D53} GSS, resulted in a novel Vienna 8^{D53} FD GSS, where females showed differences in the thermal sensibility, larval development speed, and productivity profiles. The modification of these traits and their impact on the mass rearing of the GSS for sterile insect technique applications are discussed.

Key words: Medfly, mass rearing, insect pest control, mutation

The Mediterranean fruit fly (Medfly), *Ceratitis capitata* (Wiedemann), is a major agricultural pest that causes direct damage to more than 400 fruit crops worldwide (De Meyer et al. 2002). The sterile insect technique (SIT) is successful tool for the areawide integrated pest management (IPM) programs of Medfly (Dyck et al. 2006). In SIT, mass-reared males are sterilized by irradiation and released into the field to mate with wild females, inseminating the wild females with sterile sperm resulting in fertility reduction and driving fly population densities below economic thresholds, or even local eradication (Dyck et al. 2006). SIT success relies on the mating efficiency of sterile males, which increases by threefold when only males are released (McInnis et al. 1994, Rendón et al. 2004).

Several genetic sexing strains (GSS) were developed that allow male-only releases, to improve *C. capitata* SIT efficiency (Franz 2005). These GSS are based on a reciprocal translocation between the Y-chromosome and the region of chromosome (autosome) 5 which includes the genes used as selectable markers, the *white pupae* (*wp*) and the *temperature sensitivity lethal* (*tsl*), both located on chromosome 5 (Robinson 1999, Franz 2005). In the GSS, only the *white pupae* (*wp*) marker (GSS-*wp*) was used, allowing the separation of white pupae females from brown (wild-type color) pupae males using a sorting machine (Robinson and Van Heemert 1982). Subsequently, the *temperature sensitivity lethal* (*tsl*) marker, closely linked to the *wp* gene, was isolated and used to construct a second

GSS in which the males were heterozygous for the wild-type alleles of both the *w^p* and *tsl* gene markers (Vienna 8^{D53-}) emerged from brown pupae and were resistant to elevated temperatures while the females were homozygous for the recessive alleles of the *w^p* and *tsl* gene markers, emerged from white pupae and were sensitive to high temperatures (Kerremans and Franz 1994, Franz 2005). Based on these traits, a protocol was developed for male-only production, which allowed the elimination of the double homozygous mutant females (*w^p tsl/w^p tsl*) by incubating 24-h-old embryos (eggs) at 34°C for 24 h while allowing the heterozygous males to survive.

The slow larval development phenotype was previously reported in *C. capitata* (*sw*) as well as in *Anastrepha ludens* (*sl*) (Diptera: Tephritidae) (Delprat et al. 2002, Meza et al. 2019). In both cases, the genes responsible for the slow larval development phenotype were mapped on chromosome 2. The slow larval development observed in Vienna 8^{D53-} females has been so far considered as a pleiotropic effect of the *tsl* gene, which resides on the right arm of chromosome 5 (Caceres 2002, Franz 2005). However, this genetic correlation of thermal sensitivity and slow larval development could also arise from a tight, independent linkage between two different genes controlling the respective traits. In the case of pleiotropy, the correlation between traits is inherent, while a linkage can be broken by genetic recombination (Paaby and Rockman 2013).

Genetic recombination phenomena have been observed in the Vienna 8^{D53-} GSS. For example, type-1 male recombination in Vienna 8^{D53-} can occur either between the translocation breakpoint and the *w^p* gene producing *w^p tsl* male and *w^p+* *tsl*⁻ female recombinants (type-1a) or between the *w^p* gene and the *tsl* gene producing *w^p+* *tsl* male and *w^p tsl*⁻ female recombinants (type-1b) (Franz 2002, 2005). To contain the threat of recombination to the genetic integrity of the GSS, a filter rearing system (FRS) consisting of the establishment of a small mother colony, the only colony contributing biological material to the mass-rearing system, systematically eliminating the recombinant individuals to avoid its massive accumulation (Fisher and Caceres 2000). An additional approach was the development of inversions, which are known recombination suppressors, such as the chromosomal inversion D53 used for the construction of the Vienna 8^{D53+} GSS (Franz 2002, Augustinos et al. 2017, Zacharopoulou et al. 2017).

In the mass-rearing facilities, six consecutive daily larval collections are usually obtained during the rearing of the Vienna 8^{D53-} strain. Brown pupae (males) are collected mainly during the first 2 d, while white pupae (females) are collected in the last 3 d (Caceres 2002). The slow larval development in females potentially can be used as a sexing mechanism alone or in combination with other traits like the pupal color morphological markers. However, in Medfly, the main sexing system is based on the temperature-sensitive lethal gene with the white pupae mutation acting as a selectable morphological marker. The slow development of female larvae is considered a negative trait because: 1) females may be forced to complete their development on a poor-quality larval diet since important nutrients may have been consumed by the fast developing males and 2) larval trays must be kept in the respective rearing area for two to three additional days with significant accompanying costs in terms of space, labor, and energy needed to maintain the necessary environmental rearing conditions (Caceres 2002, Meza et al. 2018).

In the present study, we report the construction of a novel *tsl*-based GSS, the Vienna 8^{D53-} FD GSS, in which the females have faster development (FD) than in the conventional *tsl* strain. The new strain was developed using classical genetic approaches and exploiting genetic recombination phenomena between the two closely linked loci, *tsl* and *slow development* (*sd*), and has novel biological

characteristics that can be used for the improvement of the mass-rearing efficiency as well as the overall enhancement and cost-effectiveness of SIT applications.

Materials and Methods

Study Site and Strains

The study was carried out at the Insect Pest Control Laboratory (IPCL) of the Joint FAO/IAEA Division of Nuclear Techniques in Food and Agriculture, Seibersdorf, Austria, using the pure *w^p*, *w^p tsl* double mutant, and the *FiM1* from the IPCL stock. The heterozygous lethal *FiM1* balancer strain, which carries the *w^p* recessive allele in repulsion and has the Sergeant dominant mutation (*Sr*²) as morphological marker (Gourzi et al. 2000), was used to avoid recombination during the isolation of the *sd*⁺ wild-type allele from an exceptional *w^p tsl sd*⁺ female, resulting in a temperature-sensitive lethal strain marked with white pupae and fast development during the larval stage (*w^p tsl* FD). We also used the Vienna 8^{D53-} strain, which is a GSS generally used for mass rearing and SIT applications almost worldwide, from two different sources (although both strains had a common origin): 1) Valencia, Spain, refreshed in 2013 using flies from Valencia (J. Garcia, personal communication) and 2) El Pino, Guatemala, refreshed in 2012 with wild flies from Guatemala (E. Ramirez, personal communication). In a preliminary test, the Vienna 8^{D53-} strain from Spain showed a higher number of white pupae in the first collection compared to the one from Guatemala and was therefore used to screen the first larval collection for the presence of *w^p*, while the genetic background of Vienna 8^{D53-} males from Guatemala were used for the construction of the novel GSS. During the comparative analysis of the novel GSS, the Vienna 8^{D53-} from Guatemala was used as a positive *tsl* reference. In addition, a GSS-*w^p* strain was constructed by backcrossing Guatemala Vienna 8^{D53-} males to pure *w^p* females and was used as a negative *tsl* reference (absence of *tsl* allele). All colonies were kept at 25 ± 1°C, 65% RH, and a photoperiod of 14:10 (L:D) h, with a photophase starting at 0800 hours.

Isolation of a Fast Developing Strain and the Construction of a Novel GSS (Vienna 8^{D53-} FD)

In an extensive screening of approximately 250,000 pupae Vienna 8^{D53-} from Spain, 50 exceptional *w^p* females were collected from the first larval collection and were crossed in single pairs with males from the *FiM1* balancer strain. From the F₁ offspring of every single cross (family), the *w^p* mutants were discarded, and the *Sr*² mutants that emerged from the brown pupae were interbred (*FiM1*, *Sr*²/*w^p tsl*) *en masse* (Fig. 1). The larval development rate (larvae were collected daily; Supp. Table S1 [online only]) was recorded in the F₂ offspring of each family. From families in which the white pupae exhibited similar larval development time as the brown pupae, all white pupae individuals with the fast development phenotype were selected and interbred, leading to the isolation of *w^p tsl* FD lines (fast developing strains).

The thermal sensitivity pattern of the fast developing strains was determined at six different heat treatments (31, 32, 33, 34, 35, and 36°C), using the *w^p* and *w^p tsl* mutant strains as a negative and positive reference, respectively. Then, the fast developing strains were selected and were used to construct the novel Vienna 8^{D53-} FD GSS by crossing and backcrossing the Vienna 8^{D53-} males from the Guatemala strain with the previously selected females from the fast developing strains.




Fig. 1. Crossing scheme used to recover the fast developing strain carrying the white pupae marker, from a single *wp tsl sd+* homozygous female (circle = centromere).

Egg Heat Treatment

Immediately after collection, the eggs were aerated in bubbling water for 24 h at 24°C and then subjected to the heat treatment (31–36°C) for another 24 h. Egg hatch was used as a response variable to measure thermal sensitivity.

Thermal Sensitivity

To measure the thermal sensitivity of the novel Vienna 8^{DS3}-FD, 1,000 eggs from each strain were exposed to four different heat treatments (24, 34, 35, and 36°C), with each treatment being replicated 10 times. After 24 h of incubation at 24°C, eggs were subject to heat treatment and then were placed onto carrot diet for eclosion and larval development (Tanaka et al. 1970). Eggs kept at 24°C during the entire incubation period were used as control. The number of white and wild-type (brown) pupae was recorded. The GSS-*wp* and Vienna 8^{DS3}- from Guatemala were used as negative and positive *tsl* references.

Larval Developmental Duration and Pupal Production Pattern

One thousand eggs from each one of the following strains GSS-*wp*, Vienna 8^{DS3}-, Vienna 8^{DS3}-FD_28, and Vienna 8^{DS3}-FD_37 were incubated for 2 d (25 ± 1°C, 65% RH) on moist filter paper placed in Petri dishes. After incubation, mature embryos were transferred into carrot diet (26°C, 50% RH) in a Petri dish (20 cm diameter × 2 cm) located inside a plastic rearing try (30 cm length × 30 cm width × 8 cm height) with the bottom covered by sawdust (Augustinos et al. 2017). This provided ventilation and space for mature larvae that leave the diet for pupation. Mature larvae and prepupae were collected daily, and upon the completion of ecdysis, the number of white and wild-type (brown) pupae was daily recorded. The pupae of each group (brown and white) were transferred to Petri dishes where adult emergence was also recorded daily. This procedure was replicated five times per strain.

Fecundity Following Mass-Rearing Procedures

Thirty single pair crosses were set up for each of the following GSS, GSS-*wp*, Vienna 8^{DS3}-, Vienna 8^{DS3}-FD_28, and Vienna

8^{DS3}-FD_37 strains in a transparent plastic cages (0.4-liter capacity) using freshly emerged adults. Each cage was provided with a standard diet of enzymatic yeast hydrolysate and sugar (1:3) with water supplied ad libitum. An oviposition device filled with water and covered with mesh was located on the cage side. Stress-related mortality (e.g., transfer) was extremely low (<1% mortality during the first 48 h postcapture). Survival of both sexes was monitored daily, and the number of eggs laid by individual females was recorded for 20 d postemergence.

Statistical Analyses

All data were tested for normality (Shapiro–Wilks test) and homogeneity of variance (Levene's test). For the isolation of fast developing strain and the construction of a novel Vienna 8^{DS3}-FD GSS experiment, the thermal sensitivity and pupal production patterns were analyzed for differences among the strains using a two-way analysis of variance (ANOVA) followed by post hoc multiple comparisons and Tukey–Kramer tests. The response variable was egg hatch and the factors were the strains and temperature (thermal treatment). For the experiment of thermal sensitivity, the response variable was the number of white pupae and brown pupae, with thermal treatment as an independent variable. A two-way ANOVA was used to analyze the differences in pupae production among the strains, while one-way ANOVAs were used to independently analyze the production of white and brown pupae. For the experiment larval developmental duration and pupal production pattern, we used ANOVA to identify differences in the time required for larval development until pupation among the GSS. For the experiment fecundity following mass-rearing procedures, we used a generalized linear model (GLM) with Poisson distribution to test for differences in fecundity (number of eggs per female) among the strains, with strains and time (days) as factors. All analyses were performed using JMP-Pro version 13.1.0.




Fig. 2. Thermal sensitivity pattern of the six selected fast developing strains compared to the *wp* and *wp tsl* mutant strains. Note that the thermal sensitivity pattern of the *wp tsl* FD₁₅ and *wp tsl* FD₇ strains suggested that the complete lethality for these two lines is achieved at 2°C higher than the other strains (mean \pm SE; $N = 10$).




Fig. 3. Number of white pupae and brown pupae (wild-type) recovered from 1,000 eggs exposed to four different thermal treatments (24, 34, 35, 35°C). Note that the number of brown pupae of Vienna 8^{D53}-FD₃₇ is higher than that of the other strains (mean \pm SE; $N = 10$).

Results

Establishment of Fast Developing Strains and Construction of a Novel Vienna 8^{D53}- FD GSS

Establishment of fast developing strains

After applying the appropriate crossing scheme illustrated in Fig. 1, fast development was observed in 6 out of 50 *wp* families where the first larval collection resulted in the production of white pupae (Supp Table S1 [online only]). The egg hatch of these six *wp* strains was greatly affected by temperature treatments (*wp tsl* FD) (two-way




Fig. 4. Daily records of the larval developmental duration for females (white pupae) and males (brown pupae), as well as daily records of the number of pupae produced. Note that the GSS-*wp* and Vienna 8^{D53}-FD₃₇ strains have a similar pattern of brown pupae production, and the females of the Vienna 8^{D53}-FD₃₇ and Vienna 8^{D53}-FD₂₈ are faster than Vienna 8^{D53}- (mean \pm SE; $N = 10$).




Fig. 5. Oviposition pattern of GSS-*wp*, Vienna 8^{D53}-, Vienna 8^{D53}-FD₂₈, and Vienna 8^{D53}-FD₃₇ strains (mean \pm SE; $N = 30$).

ANOVA: $F_{15,263} = 54.42$, $P < 0.0001$; Supp Tables S2 and S3 [online only]; Fig. 2). The strain, temperature, and their interaction significantly affected egg hatching. The *wp tsl* FD₇, ₉, ₁₅, ₂₈, ₃₆, and ₃₇ strains had different thermal sensitivity patterns compared to the normal *tsl* strain. Specifically, the *wp tsl* FD₉, ₂₈, ₃₆, and ₃₇ lines presented 100% lethality at 35°C, while the *wp tsl* FD₇ and ₁₅ strains demonstrated, complete lethality at 36°C.

Vienna 8D53- FD construction

Due to their higher productivity (recovery of pupae and adults per given number of embryos; data not shown) and complete lethality at 35°C, the *wp tsl* FD₂₈ and ₃₇ strains were selected for the construction of two




Fig. 6. Schematic representation of type-1b and the possible type-1c recombination outcomes. The sd^+ allele results in different phenotypic traits that can be obtained by type-1c recombination and can only be evident in an eventual homozygosis. Type-1b recombination results in the production of heterozygous $tsl sd$ females, which are resistant to elevated temperatures, thus disrupting the sexing mechanism. Type-1c recombination results in the production of homozygous tsl females but heterozygous for sd , which are sensitive to elevated temperatures and develop faster than the original Vienna 8^{D53-} .

novel GSS. This was achieved by crossing the first-emerging female from the $wp tsl$ FD strain with Vienna 8^{D53-} Guatemala males and backcrossing the F1 males with the respective females of each $wp tsl$ FD strain, thus resulting in the Vienna 8^{D53-} FD_28 and Vienna 8^{D53-} FD_37 GSS.

Thermal Sensitivity

The pupal productivity of the novel Vienna 8^{D53-} FD_28, Vienna 8^{D53-} FD_37 strains as well as the existing Vienna 8^{D53-} and GSS- wp strains was tested under normal rearing conditions (24°C), and their thermal sensitivity was determined. At 24°C, the Vienna 8^{D53-} FD_37 strain was the most productive, followed by the GSS- wp . After heat treatment, white pupae were only produced by the GSS- wp , the strain that did not carry the tsl gene. The number of brown pupae significantly varied among the strains (two-way ANOVA: $F_{7,152} = 41.02$, $P < 0.0001$; [Supp Tables S4 and S5](#) [online only]; [Fig. 3](#)). At 34, 35, and 36°C, the number of brown pupae produced depended on the temperature ($F_{1,152} = 25.34$, $P < 0.0001$), the strain ($F_{3,152} = 79.34$, $P < 0.0001$), and their interaction ($F_{1,152} = 7.92$, $P < 0.0001$), but the Vienna 8^{D53-} FD_37 strain kept their productivity level in all tested temperatures. Similarly, the number of white pupae produced depended on the temperature, strain, and their interaction of both factors ($F_{1,152} = 10.08$, $P = 0.0018$; $F_{3,152} = 119.75$, $P < 0.0001$; $F_{1,152} = 6.60$, $P = 0.01$; [Fig. 3](#)).

Larval Developmental Duration and Pupal Production Pattern

As shown in [Fig. 4](#), larval development required significantly less time for the GSS- wp compared to all other strains, which were all carrying the tsl gene (Vienna 8^{D53-} FD_28, Vienna 8^{D53-} FD_37, and Vienna 8^{D53-}), with no evidence of sex separation ($F_{3,16} = 15.16$, $P < 0.0001$). The production pattern of brown pupae was significantly different among the four strains studied, but only males from GSS- wp were statistically faster than Vienna 8^{D53-} , Vienna 8^{D53-} FD_28 and _37 ($F_{3,16} = 10.90$, $P = 0.0004$). The wp production pattern of both novel GSS (Vienna 8^{D53-} FD_28 and _37) proved to be faster than Vienna 8^{D53-} but slower than GSS- wp , where the tsl is absent.

Fecundity Following Mass-Rearing Procedures

The fecundity of Vienna 8^{D53-} FD_28, Vienna 8^{D53-} FD_37, Vienna 8^{D53-} , and GSS- wp strains was significantly greater in the Vienna 8^{D53-} FD_37 compared to the other three strains ($F_{79,1318} = 32.28$, $P < 0.0001$; [Supp Tables S6 and S7](#) [online only]; [Fig. 5](#)); the Vienna 8^{D53-} FD_37 females laid 85% more eggs than the females of the other strains. The first oviposition was asynchronous, with the Vienna 8^{D53-} FD_37 females laying eggs a day earlier than the other three strains. As clearly shown

in Fig. 5, the most productive period for all four strains was between the 4th and the 10th day after emergence, while a significant reduction in egg laying was observed after the 14th day (Fig. 5).

Discussion

The present study presents clear evidence that the *tsl* genetic region of *C. capitata* does directly control the slow larva development. Through the use of classical genetic approaches and the exploitation of genetic recombination phenomena between the two closely linked loci, *tsl* and *sd*, we were able to isolate a new version of the *tsl* strain in which the larvae develop faster, almost like wild-type strains (Fig. 4). This new *w^p tsl* FD phenotype was then used for the construction of novel GSS (Vienna 8^{DS3-} FD), which shortens the larval developmental time of females, exhibits differences in their pupal production pattern, and also affects the thermosensitivity profile, properties which are expected to significantly improve mass rearing, reduce costs, and enhance SIT applications.

Traditionally, during the mass rearing of Vienna 8^{DS3-}, white pupae females have been detected during the first day of larvae collection. In most cases, these white pupae females were resistant to elevated temperatures, and this trait was attributed to the type-1b recombination phenomena, where the *tsl* allele is lost (Fig. 6; type-1b recombination). However, we showed that some of these white pupae females are still sensitive to high temperatures, which is probably due to a different type of recombination (Fig. 6; type-1c recombination), resulting in the modification of the phenotypic properties like developmental time and fitness of the normal *tsl* strain.

The isolation of recombinants between the *tsl* and *sd* genetic loci allowed the selection of the proper alleles of these two selectable markers for the construction of a novel GSS using the genetic background of Vienna 8^{DS3-}, namely Vienna 8^{DS3-} FD, which displayed differences in thermal sensitivity, pupal production pattern, and the fecundity profile compared to the parental strain. Although the Vienna 8^{DS3-} FD showed a faster female developmental time than that of the Vienna 8^{DS3-} females, it was not as fast as that of the wild-type male (*w^p tsl⁺*). Furthermore, it was necessary to increase the temperature during the heat treatment from 34 to 35 or even 36°C to induce the 100% female lethality required to allow the production of male-only pupae. Indeed, in the case of the Vienna 8^{DS3-} FD₂₈ and Vienna 8^{DS3-} FD₃₇ strains, complete lethality was achieved at 35°C without affecting fecundity and male survival. It is worth noting that the Vienna 8^{DS3-} FD₃₇ GSS was shown to be the most productive in terms of fecundity and egg to pupa survival. Along with variations in the rearing practices, the phenotypic traits of the new *w^p tsl* FD strain may explain the differences observed in the Vienna GSS, all of common origin, used in the Medfly mass-rearing facilities worldwide (Augustinos et al. 2017).

Optimization and cost reduction of the mass-rearing process has the potential to improve the cost-benefit ratio of current operational SIT programs and facilitate the implementation of new ones. Therefore, the development of new Medfly GSS with better biological performance, such as faster development of females during the larval stage and females with higher fecundity (Vienna 8^{DS3-} FD₃₇) and viability, suggesting that the Vienna 8^{DS3-} FD has real potential to improve the mass-rearing process, efficiency, and cost-effectiveness. The thermotolerance observed in the Vienna 8^{DS3-} FD will benefit the effectiveness of the strain because the broad thermal range tolerated by females may permit more flexibility in the rearing temperatures (Caceres

2002). The potential use of this novel GSS for SIT applications will first require the development of novel protocols for mass rearing as well as its thorough evaluation as concerns the rearing efficiency, genetic stability, and biological quality, including male mating competitiveness.

Supplementary Data

Supplementary data are available at *Journal of Economic Entomology* online.

Acknowledgments

We thank Jaime Garcia and Edwin Ramirez for kindly providing strains used in the present study. We also thank the technicians of the Plant Pests and Genetics and Molecular Biology groups of the Insect Pest Control Laboratory for their excellent technical support in the frame of this study.

References Cited

- Augustinos, A., A. Targovska, E. Cancio-Martinez, E. Schorn, G. Franz, C. Cáceres, A. Zacharopoulou, and K. Bourtzis. 2017. *Ceratitis capitata* genetic sexing strains: laboratory evaluation of strains from mass-rearing facilities worldwide. *Entomol. Exp. Appl.* 164: 305–317.
- Caceres, C. 2002. Mass rearing of temperature sensitive genetic sexing strains in the Mediterranean fruit fly (*Ceratitis capitata*). *Genetica*. 116: 107–116.
- Delprat, M. A., C. E. Stolar, F. C. Manso, and J. L. Cladera. 2002. Genetic stability of sexing strains based on the locus *sw* of *Ceratitis capitata*. *Genetica*. 116: 85–95.
- De Meyer, M., R. S. Copeland, R. A. Wharton, B. A. McPheron, and B. N. Barnes. 2002. On the geographic origin of the Medfly *Ceratitis capitata* (Wiedemann) (Diptera: Tephritidae). pp. 45–53. *In* Proceedings of the 6th International Symposium on Fruit Flies of Economic Importance, Stellenbosch, 6–10 May, 2002, Iste Scientific Publications, Irene, South Africa.
- Dyck, V. A., J. Hendrichs, and A. S. Robinson. 2006. Sterile insect technique: principles and practice in area-wide integrated pest management. Springer, Dordrecht, The Netherlands.
- Fisher, K., and C. Caceres. 2000. A filter rearing system for mass reared genetic sexing strains of Mediterranean fruit fly (Diptera: Tephritidae), pp. 543–550. *In* Area-wide control of fruit flies and other insect pests. Joint proceedings of the international conference on area-wide control of insect pests, 28 May–2 June, 1998 and the Fifth International Symposium on Fruit Flies of Economic Importance, Penang, Malaysia, 1–5 June 1998. Penerbit Universiti Sains Malaysia, Penang, Malaysia.
- Franz, G. 2002. Recombination between homologous autosomes in Medfly (*Ceratitis capitata*) males: type-1 recombination and the implications for the stability of genetic sexing strains. *Genetica*. 116: 73–84.
- Franz, G. 2005. Genetic sexing strains in Mediterranean fruit fly, an example for other species amenable to large-scale rearing for the sterile insect technique, pp. 427–451. *In* V. A. Dyck, J. Hendrichs, and A. S. Robinson (eds.), Sterile insect technique principles and practice in area-wide integrated pest management. Springer, Dordrecht, The Netherlands.
- Gourzi, P., D. Gubb, Y. Livadaras, C. Caceres, G. Franz, C. Savakis, and A. Zacharopoulou. 2000. The construction of the first balancer chromosome for the Mediterranean fruit fly, *Ceratitis capitata*. *Mol. Gen. Genet.* 264: 127–136.
- Kerremans, P., and G. Franz. 1994. Cytogenetic analysis of chromosome 5 from the Mediterranean fruit fly, *Ceratitis capitata*. *Chromosoma*. 103: 142–146.
- McInnis, D., S. Tam, C. Grace, and D. Miyashita. 1994. Population suppression and sterility rates induced by variable sex ratio, sterile insect releases of *Ceratitis capitata* (Diptera: Tephritidae) in Hawaii. *Ann. Entomol. Soc. Am.* 87: 231–240.
- Meza, J. S., I. Ul Haq, M. J. B. Vreysen, K. Bourtzis, G. A. Kyritsis, and C. Cáceres. 2018. Comparison of classical and transgenic genetic sexing

- strains of Mediterranean fruit fly (Diptera: Tephritidae) for application of the sterile insect technique. *PLoS One*. 13: e0208880.
- Meza, J. S., C. Cáceres, and K. Bourtzis. 2019. Slow larvae mutant and its potential to improve the pupal color-based genetic sexing system in Mexican fruit fly, (Diptera: Tephritidae). *J. Econ. Entomol.* 112: 1604–1610.
- Paaby, A. B., and M. V. Rockman. 2013. The many faces of pleiotropy. *Trends Genet.* 29: 66–73.
- Rendón, P., D. McInnis, D. Lance, and J. Stewart. 2004. Medfly (Diptera: Tephritidae) genetic sexing: large-scale field comparison of males-only and bisexual sterile fly releases in Guatemala. *J. Econ. Entomol.* 97: 1547–1553.
- Robinson, A. 1999. Genetic sexing strains in the Medfly, *Ceratitis capitata*: development, mass rearing and field application. *Trends Entomol.* 2: 81–104.
- Robinson, A., and C. Van Heemert. 1982. *Ceratitis capitata*—a suitable case for genetic sexing. *Genetica*. 58: 229–237.
- Tanaka, N., R. Okamoto, and D. Chambers. 1970. Methods of mass rearing the Mediterranean fruit fly currently used by the US Department of Agriculture, pp. 19–23. *In* The Proceedings on the Sterile Male Techniques for Control of Fruit Flies. International Atomic Energy Agency, 1–5 September 1969, Vienna, Austria.
- Zacharopoulou, A., A. Augustinos, E. Drosopoulou, K. Tsoumani, A. Gariou-Papalexiou, G. Franz, K. Mathiopoulos, K. Bourtzis, and P. Mavragani-Tsipidou. 2017. A review of more than 30 years of cytogenetic studies of Tephritidae in support of sterile insect technique and global trade. *Entomol. Exp. Appl.* 164: 204–225.