

## **Landscape genetics of wolverines (*Gulo gulo*): scale-dependent effects of bioclimatic, topographic, and anthropogenic variables**

Authors: Balkenhol, Niko, Schwartz, Michael K., Inman, Robert M., Copeland, Jeffrey P., Squires, John S., et al.

Source: Journal of Mammalogy, 101(3) : 790-803

Published By: American Society of Mammalogists

URL: <https://doi.org/10.1093/jmammal/gyaa037>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.



## Landscape genetics of wolverines (*Gulo gulo*): scale-dependent effects of bioclimatic, topographic, and anthropogenic variables

NIKO BALKENHOL,\* MICHAEL K. SCHWARTZ, ROBERT M. INMAN, JEFFREY P. COPELAND, JOHN S. SQUIRES, NEIL J. ANDERSON, AND LISETTE P. WAITS

Wildlife Sciences, University of Goettingen, Buesgenweg 3, 37077 Goettingen, Germany (NB)

USDA Forest Service Rocky Mountain Research Station, 800 E. Beckwith, Missoula, MT 59801, USA (MKS, JPC, JSS)

Wildlife Conservation Society, 222 East Main Street, Ennis, MT 5972, USA (RMI)

Montana Fish, Wildlife and Parks, 490 North Meridian Road, Kalispell, MT 59901, USA (NJA)

Department of Fish & Wildlife Sciences, University of Idaho, 875 Perimeter Dr., Moscow, ID 83844-1136, USA (NB, LPW)

Present address of RMI: Montana Fish, Wildlife and Parks, 1420 East 6th Ave., Helena, MT 59620, USA

\*Correspondent: [nbalken@gwdg.de](mailto:nbalken@gwdg.de)

Climate change can have particularly severe consequences for high-elevation species that are well-adapted to long-lasting snow conditions within their habitats. One such species is the wolverine, *Gulo gulo*, with several studies showing a strong, year-round association of the species with the area defined by persistent spring snow cover. This bioclimatic niche also predicts successful dispersal paths for wolverines in the contiguous United States, where the species shows low levels of genetic exchange and low effective population size. Here, we assess the influence of additional climatic, vegetative, topographic, and anthropogenic, variables on wolverine genetic structure in this region using a multivariate, multiscale, landscape genetic approach. This approach allows us to detect landscape-genetic relationships both due to typical, small-scale genetic exchange within habitat, as well as exceptional, long-distance dispersal among habitats. Results suggest that a combination of snow depth, terrain ruggedness, and housing density, best predict gene flow in wolverines, and that the relative importance of variables is scale-dependent. Environmental variables (i.e., isolation-by-resistance, IBR) were responsible for 79% of the explained variation at small scales (i.e., up to ~230 km), and 65% at broad scales (i.e., beyond ~420 km). In contrast, a null model based on only space (i.e., isolation-by-distance, IBD) accounted only for 17% and 11% of the variation at small and broad scales, respectively. Snow depth was the most important variable for predicting genetic structures overall, and at small scales, where it contributed 43% to the variance explained. At broad spatial scales, housing density and terrain ruggedness were most important with contributions to explained variation of 55% and 25%, respectively. While the small-scale analysis most likely captures gene flow within typical wolverine habitat complexes, the broad-scale analysis reflects long-distance dispersal across areas not typically inhabited by wolverines. These findings help to refine our understanding of the processes shaping wolverine genetic structure, which is important for maintaining and improving functional connectivity among remaining wolverine populations.

**Key words:** corridor, genetic distance, hierarchical partitioning, landscape connectivity, mating movements, microsatellites, multiple regression on distance matrices, mustelid, spatial variance partitioning, weighted distances

The wolverine (*Gulo gulo*) is the largest terrestrial member of the mustelid family and is globally distributed throughout the boreal zone of the northern hemisphere. Wolverine occurrences worldwide are strongly associated with cold areas offering persistent spring snow cover, and year-round habitat use of the species generally is restricted to this narrow bioclimatic niche (Aubry et al. 2007; Copeland et al. 2010; Inman et al. 2013). In

the contiguous United States, historical wolverine populations were extirpated by the early 20th century, and contemporary populations in the lower 48 States were reestablished by immigrants from the north (McKelvey et al. 2014). The species still is absent from Utah and only single individuals have been confirmed in California (Moriarty et al. 2009), Colorado (Packila et al. 2017), and Oregon (Oregon Wild 2019), all of which were

© The Author(s) 2020. Published by Oxford University Press on behalf of the American Society of Mammalogists.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits non-commercial re-use, distribution, and reproduction in any medium, provided the original work is properly cited. For commercial re-use, please contact [journals.permissions@oup.com](mailto:journals.permissions@oup.com)

part of the historical distribution of the species (McKelvey et al. 2014). Currently, the species occupies insular, high-elevation habitat patches in Idaho, Montana, Washington, and Wyoming (Banci 1994; Aubry et al. 2007; McKelvey et al. 2014). In this southernmost part of their distribution, wolverines live at low densities (e.g., Inman et al. 2012a) and have an estimated effective population size of only 28–52 individuals (Schwartz et al. 2009). Compared to wolverines in Canada and Alaska, wolverines in the contiguous United States also show lower levels of genetic diversity, and higher levels of genetic differentiation (Kyle and Strobeck 2001, 2002; Cegelski et al. 2003, 2006; Schwartz et al. 2007; McKelvey et al. 2011). To ensure the long-term viability of wolverines in the lower 48 States, it is particularly important to understand and maintain population connectivity and gene flow across broad spatial scales (Carroll et al. 2001, 2015; Schwartz et al. 2009; Inman et al. 2013).

Landscape genetic approaches make it possible to study the effects of landscape heterogeneity on dispersal and population connectivity in highly mobile, low-density species such as the wolverine (Storfer et al. 2007; Waits et al. 2016). A typical way for evaluating such landscape-genetic relationships is through landscape resistance modeling (Spear et al. 2010, 2016). In this approach, the study landscape is modeled as a raster layer where each cell receives a value reflecting the hypothesized resistance of the landscape to animal movement and gene flow, for example due to physical or behavioral constraints (Zeller et al. 2012). This model then is used to estimate spatial proximity among sampling locations via “effective distances” (e.g., through least-cost paths), which account for the hypothesized landscape resistances. Finally, various statistical methods can be used to test for relationships between measures of genetic connectivity (e.g., genetic distances) and the estimates of landscape connectivity (i.e., effective distances—Wagner and Fortin 2016; Fortune et al. 2018; Shirk et al. 2018). A significant relationship between genetic and effective distances indicates that the modeled landscape resistance predicts spatial-genetic structures, a pattern often referred to as “isolation-by-resistance” (IBR—Cushman et al. 2006; McRae 2006). IBR often predicts genetic structures better than classic “isolation-by-distance” (IBD—Wright 1943), where genetic distances increase with increasing geographic (i.e., straight-line) distances among sampling locations. While IBD assumes that the landscape is homogeneous from the perspective of the study species, IBR accounts for species-specific landscape heterogeneity.

Such a landscape genetic approach was used to show that the distribution of persistent spring snow cover predicts wolverine gene flow in the contiguous United States (Schwartz et al. 2009). Wolverines have a strong association with the area defined by persistent spring snow cover, and the species depends on deep, persistent snow for denning and food caching (Copeland et al. 2010; Inman et al. 2013). Areas with persistent spring snow cover also predict wolverine occurrences during summer months, highlighting the year-round association of the species with a narrow bioclimatic niche (Copeland et al. 2010).

However, in addition to its narrow bioclimatic niche, gene flow in wolverines likely can be predicted by other landscape


variables; for example, those related to terrain or anthropogenic infrastructures (e.g., May et al. 2006; Webb et al. 2016; Heim et al. 2017; Kortello et al. 2019). To understand landscape effects on wolverine dispersal and gene flow, and to effectively maintain meta-population connectivity (e.g., through conservation corridors), it is therefore important to evaluate the influence of additional landscape variables on wolverine dispersal and gene flow.

Dispersal and gene flow in mammals can arise from multiple underlying processes that act across different spatial scales (e.g., Travis et al. 2013). Specifically, mammals usually disperse over small spatial scales, while long-distance dispersal in contrast is rare (Sutherland et al. 2000; Whitmee and Orme 2013). Occasional long-distance dispersal events often require individuals to venture into unfamiliar areas, where they encounter landscape elements and influencing factors not present in their typical habitat (e.g., Walton et al. 2018; Bartoń et al. 2019). On account of these factors, several recent studies have shown that it is challenging to predict gene flow from habitat suitability models (e.g., Mateo-Sánchez et al. 2015; Roffler et al. 2016), because these models capture the suitability of the landscape for movement and occurrence of the species, but not necessarily the resistance of the landscape to long-distance dispersal movements. Often, individuals still are able and willing to disperse through certain areas, even if these areas are unsuitable for settlement and breeding (Mateo-Sánchez et al. 2015; Keeley et al. 2016; Abrahms et al. 2017). Current landscape genetic approaches do not distinguish among these different processes leading to gene flow, and rarely test for scale effects on landscape-genetic relationships.

Here, we extend the analysis of Schwartz et al. (2009) by including additional landscape variables in a multivariate, multiscale landscape genetics framework that can detect environmental predictors of genetic structure resulting both from typical, small-scale genetic exchange within a habitat, as well as exceptional, long-distance dispersal among habitats. Specifically, our study addresses three interrelated questions: (1) What is the relative importance of pure space (IBD) versus landscape heterogeneity (IBR) for predicting wolverine genetic structure? (2) What is the relative importance of different landscape variables for predicting wolverine genetic structure? (3) Are scale effects detectable in the data set and, if so, how do they affect the answers to questions (1) and (2)?

## METHODS

*Genetic data set.*—The genetic data set used in this study previously was analyzed by Schwartz et al. (2009) and includes 210 wolverine tissue samples gathered across northern and central Idaho, western Montana, and northwestern Wyoming. The study area covers approximately 330,000 km<sup>2</sup>, including the entire Rocky Mountain range of wolverines in the contiguous United States (Fig. 1). Most samples are from harvested animals ( $N = 142$ ), with the remainder stemming from different research projects. Samples were genotyped at 16 microsatellite loci with an expected and observed heterozygosity



**Fig. 1.**—The study area includes states of Idaho (ID), Montana (MT), and Wyoming (WY) and borders Canada (CAN) in the north. Darker colors depict higher wolverine-specific landscape resistances inferred from the multivariate, broad-scale analysis (see main text for details). Locations of genetic wolverine samples are shown as white circles. The inset shows the location of the study area in the western United States.

of 0.504 and 0.473, respectively. Mean number of alleles per locus was 4.06; a total of 65 alleles were detected. Additional details about sampling and laboratory analyses are detailed by Schwartz et al. (2009). This data set was used to calculate Miller's genetic distance  $D_{ij}$  for all pairs of individuals  $i$  and  $j$  with the software *Alleles in Space* (Miller 2005). This distance is the individual-based equivalent to the commonly used measure  $D$  for estimating genetic differentiation among populations (Nei et al. 1983) and takes on values between 0 (identical genotypes) and 1 (no allele in common). Since landscape genetic conclusions can depend somewhat on the choice of the genetic distance metric employed (Shirk et al. 2017), we also estimated alternative measures and compared results obtained with each genetic distance. Specifically, we estimated Rousset's  $a$  (Rousset 2000) in software Spagedi (Hardy and Vekemans 2002), and Smouse and Peakall's  $r$  (Smouse and Peakall 1999) in software Genalex (Peakall and Smouse 2005).

#### *Landscape data and suitability for providing connectivity.*

As a first step for landscape resistance modeling, we chose eight digital landscape layers of variables that we hypothesized to be predictors of wolverine functional connectivity (Table 1). These variables were among those used by Inman et al. (2013) to predict habitat suitability for wolverines across the western United States. We chose these variables because wolverines in our study area occupy rugged, high-elevation habitats that generally follow the timberline (Hornocker and Hash 1981; Copeland 1996; Edelmann and Copeland 1999; Squires et al. 2006; Aubry et al. 2007; Copeland et al. 2007; Inman et al. 2012a, 2013). These habitats include conifer forests as well as forest edge, e.g., areas where forested and nonforested areas are adjacent (Brock et al. 2007). Wolverines also appear sensitive to anthropogenic influences (Copeland 1996; Rowland et al. 2003; Heim et al. 2017), and require deep spring snowpack for denning (Magoun and Copeland 1998; Aubry et al. 2007). Variables chosen for this study thus included a climatic



**Table 1.**—GIS data layers and sources used in the dispersal suitability models.

Type of variable	Variable	Abbreviation	Description (assumed effect on connectivity)	Source	Original resolution	Reference
<i>Climatic</i>	Snow depth	snow	Average 1-year snow depth (positive)	Snow Data Assimilation System (SNODAS)	1 km	Barret (2003)
<i>Topographic</i>	Absolute elevational difference from timberline	absdiff	Difference in elevation from 300-m buffer around estimated occurrences of ecological timberline (negative)	National Elevation Data	30 m	Brock and Inman (2006)
	Elevational difference from buffered timberline	timberline	Difference in elevation from estimated occurrence of ecological timberline (negative)	National Elevation Data	30 m	Brock and Inman (2006)
	Terrain ruggedness index	tri	Topographic heterogeneity (positive)	National Elevation Data	30 m	Riley et al. (1999)
<i>Anthropogenic</i>	Housing density	hden	Census block housing density per 1 km <sup>2</sup> (negative)	Wildland-Urban Interface Data (WUI)	1:100,000	Radeloff et al. (2005)
	Human population density	popden	Interpolated human population density (negative)	US Census Bureau, census block data	1:100,000	Carroll et al. (2001)
	Road density	roaden	Interpolated road density (negative)	US Geological Survey, Reston, Virginia	1:100,000	Carroll et al. (2001)
<i>Vegetative</i>	Forest edge habitat	edge	Amount of edge habitat within 300-m radius (positive)	National Land Cover Data (NLCD)	30 m	Homer et al. (2001)
	Conifer cover	cover	Amount of conifer cover within 300-m radius (positive)	National Land Cover Data (NLCD)	30 m	Homer et al. (2001)

variable (snow depth), two topographic variables (terrain ruggedness, and latitude-adjusted elevation), three anthropogenic variables (human population density, housing density, and road density), and two vegetative variables (forest edge, and conifer cover). Further information on the different data layers can be found in [Supplementary Data SD1](#).

We did not know a priori whether all of these landscape variables actually predict wolverine genetic structure, nor could we translate the landscape data into exact resistance values. However, we were able to hypothesize whether the relationship between the variable and genetic structure should be positive (increasing genetic connectivity with increasing variable values), or negative (decreasing connectivity with increasing variable values). For example, we hypothesized that if housing density predicts genetic structure in wolverines, this association most likely would be negative, with higher housing densities predicting reduced functional connectivity for wolverines. Thus, we rescaled the housing density layer so that highest values became 0 (lowest suitability for providing connectivity) and lowest values became 1 (highest suitability for providing connectivity). We rescaled the other anthropogenic data layers in the same way, thus hypothesizing that a potential relationship between these variables and wolverine functional connectivity would be negative. For all other data layers, we generally hypothesized that a potential relationship between the variable and connectivity would be positive (i.e., data were rescaled so that highest values became 1 and lowest values became 0). The only exception was the latitude-adjusted elevation index, which we used in two different ways. First, we calculated the absolute elevational difference between any cell in the landscape

and the elevation of the alpine timberline at the latitude of that cell and rescaled the data to range from 0 (highest elevational difference) to 1 (elevation of timberline, “absdiff” in [Table 1](#)). This rescaling assumes that wolverine dispersal and gene flow generally follow the alpine timberline, as the occurrence of the species is associated with high-elevation habitats along this line ([Squires et al. 2006](#); [Aubry et al. 2007](#); [Copeland et al. 2007](#); [Inman et al. 2012a](#)). However, this rescaling also assumes that even small differences between actual elevation and the elevation of the timberline reduce the suitability for successful wolverine dispersal, which is a rather restrictive assumption. We thus also rescaled latitude-adjusted elevation in an alternative way, where all areas within 300-m elevational difference to the timberline received a value of 1 (highest suitability for providing connectivity, “timberline” in [Table 1](#)). This use of elevation data for landscape resistance modeling is similar to that of other landscape genetic studies (e.g., [Cushman et al. 2006](#); [Shirk et al. 2010](#); [Wasserman et al. 2010](#)).

The rescaling approach only makes very general assumptions about the potential suitability of an area for providing connectivity via dispersal and ensures that the different variables are measured on the same scale (0 for lowest suitability, 1 for highest suitability), which is important for assessing their relative importance for predicting genetic structures.

*Converting landscape data into resistance models.*—Next, we converted the rescaled data into actual landscape resistance values, using the concept of weighted distances ([Singleton et al. 2002](#)). In this approach, each cell of the landscape is assigned a weighted distance value that reflects the hypothesized resistance of that cell to movement and gene flow. When

the landscape in a cell is not assumed to provide resistance, the weighted distance is simply the length of the shortest possible path through the entire cell. In our case, with a cell size of 1 km<sup>2</sup>, this shortest path has a length of 1,000 m. In contrast, when the landscape is assumed to have high resistance for successful dispersal, the weighted distances is 100 times greater than the shortest possible path through the cell (i.e., 100,000 m in our case). We used this approach to convert our rescaled landscape data into resistance values, so that landscape cells with low suitability for providing connectivity (rescaled data value = 0) received highest resistance values. In contrast, landscape cells with highest potential for successful dispersal (rescaled data value = 1) received lowest resistance values. Following [Mateo-Sánchez et al. \(2015\)](#), we used three different translations for this step.

First, we used a linear conversion to derive weighted distances from our rescaled data (see [Fig. 2](#), dashed graph):


$$WD = CS * (100 - 100 * DS) \quad (1)$$

where WD is the resistance assigned to a specific cell (i.e., the weighted distance), CS is the length of the shortest possible path through the cell ("cell size," i.e., 1,000 m), and DS is the value of the rescaled data ("dispersal suitability," range 0–1).

In addition to this linear translation, we also defined two exponential conversions which account for possible nonlinear relationships between landscape data values and actual resistance ([Fig. 2](#), dotted graph):

$$WD = CS * 100^{(1-DS)} \quad (2)$$

This second conversion assumes that decreased suitability for connectivity via dispersal increases landscape resistance most strongly when suitability is low. This is the kind of translation that has been recommended for deriving landscape resistances from habitat suitability values ([Keeley et al. 2016](#)), because animals often are still able and willing to move through nonhabitat areas, even if these areas are not suitable for settlement or breeding (e.g., [Mateo-Sánchez et al. 2015](#)).



**Fig. 2.**—Calculation of landscape resistances based on hypothesized dispersal suitability using linear (dashed line), upper exponential (solid line), and lower exponential (dotted line) conversions.

Finally, we used an inverse-symmetrical version of the previous formula, which assumes that reduced suitability for dispersal quickly results in higher landscape resistances ([Fig. 2](#), solid graph):

$$WD = CS * (100 - 100^{DS}) \quad (3)$$

*Estimating effective distances from landscape resistance models.*—Based on the resulting 27 resistance surfaces (9 rescaled landscape layers × 3 weighted distance translations), we calculated effective distances using two approaches. First, we calculated pairwise least-cost distances among individual wolverines with the ArcView extension PATHMATRIX ([Ray 2005](#)). Least-cost distances measure effective separation among all pairs of samples based on a single, optimal (i.e., least-costly) path through the resistance landscape ([Adriaensen et al. 2003](#)). Thus, the least-cost approach implicitly assumes that effective dispersal and resulting genetic structures are influenced by optimal animal choices that are based on substantial knowledge of the landscape.

In addition to least-cost paths, we also calculated pairwise effective resistances in program CIRCUITSCAPE 3.4 ([McRae and Shah 2013](#)). Effective resistances are based on circuit theory and estimate effective separation distances in terms of current flows across the resistance landscape ([McRae 2006](#); [McRae and Beier 2007](#)). This approach assumes that movement of individuals is not optimal and based on incomplete knowledge of the landscape, so that multiple paths contribute to effective dispersal. Since inferences derived from the two approaches can differ (e.g., [Avon and Bergès 2016](#); [McClure et al. 2016](#); [Marrotte and Bowman 2017](#)), we applied both approaches and used objective criteria to choose the best approach for our data (see next section).

*Model optimization and exclusion of spurious variables.*—We next conducted an exploratory analysis to assess how best to represent each variable (i.e., which resistance translation to use and whether to use least-cost distances or effective resistances), and to select variables for the final, multivariate analysis (i.e., excluding spurious variables). For this, we first used simple Mantel tests ([Mantel 1967](#)) to evaluate the correlation of each effective distance with the genetic distances. For each environmental variable, the representation giving highest correlation coefficients was chosen. Any variable that did not lead to significant correlations was excluded from further analyses. Next, we used partial Mantel tests ([Smouse et al. 1986](#)) to determine whether the best representation of each variable is significantly correlated with genetic distances after accounting for the effects of space (i.e., geographic distances). Thus, we controlled for the effects of geographic distance and then assessed whether effective distances based on environmental variables have an influence on genetic structure "beyond IBD." Thus, we essentially used IBD as a null model, which assumes that environmental heterogeneity does not impact genetic structure ([Jenkins et al. 2010](#)). Finally, simple Mantel tests among all effective distances were calculated, and if two effective distances showed a correlation

of  $r > 0.7$ , one of them was excluded from further analyses (i.e., the one that gave lower correlations with the genetic data—Dormann et al. 2013). Because each effective distance was used for multiple statistical tests, inferences were based on Bonferroni-corrected  $P$ -values. This approach ensures that only effective distances are retained for further analyses that are (1) significantly correlated with genetic distances “beyond IBD,” and (2) not significantly correlated with any other effective distances.

**Assessing relative variable importance.**—Simple and Mantel tests are appropriate for the exploratory data analysis described above, but they are not suitable to assess the relative importance of different landscape variables (Cushman et al. 2013a; Prunier et al. 2015). Hence, we used a combination of alternative methods to analyze the final, multivariate data set. All analyses were based on multiple regression on distance matrices (MRM), which is essentially a multivariate regression performed on the unfolded, lower-triangle distance matrices (Legendre 2000; Lichstein 2007). The method is one of the most effective ways for analyzing landscape genetic data (Balkenhol et al. 2009; Wang 2013), and ideally is combined with variance partitioning techniques to infer relative variable importance (Prunier et al. 2015, 2017). Here, we combined MRM with two types of variance partitioning. First, to distinguish IBD effects from actual landscape influences, we used the spatial variance partitioning approach of Borcard et al. (1992). In this approach, the variation explained by a regression model is partitioned into three different components. *Spatial variation* is the amount of variation that is solely due to endogenous factors, such as landscape-independent space-use or species-specific dispersal limitations, i.e., IBD, which is represented by straight-line geographic distances. *Environmental variation* is the component that can be attributed to purely exogenous factors, such as the heterogeneous landscape, i.e., IBR, which is represented by effective distances (i.e., least-cost paths or circuit-theoretic resistances). Finally, there is a spatial component of environmental factors, so that there also is a *shared component*, which cannot be separated into purely environmental versus purely spatial contributions (e.g., because of correlations between straight-line and effective distances). Separating these three components via spatial variance partitioning commonly is used in many other ecological disciplines (e.g., Wagner 2004; Wagner and Fortin 2005; Lichstein 2007). In landscape genetics, spatial variance partitioning can be used to answer the question: What is the relative importance of space (IBD) versus environment (IBR) for predicting spatial-genetic structure?

Second, to assess the relative importance of the different effective distances for predicting genetic distances, we used hierarchical partitioning (Chevan and Sutherland 1991; Mac Nally 2000, 2009), which is one of the most effective approaches for assessing relative variable importance in ecological studies (Murray and Conner 2009). The approach assumes that a hierarchy of models exists, ranging from the simplest, univariate models to the most complex model involving all variables. The method considers all possible models that can be constructed with the set of candidate variables, and estimates the increase

in model fit generated by a certain variable for each model in which the variable is included. Hierarchical partitioning can be seen as an alternative to model selection (e.g., based on AIC values), which cannot be applied to pairwise distance matrices (Franckowiak et al. 2017). In landscape genetics, hierarchical partitioning can be used to answer the question: What is the relative importance of different landscape variables for predicting spatial-genetic structure? Since we were particularly interested in assessing landscape-genetic relationships independently from any IBD effects, we conducted hierarchical partitioning with the original data, and with residuals from regressions of geographic distances on all other distances (i.e., partial regression—Wagner and Fortin 2016).

**Testing for scale effects.**—Finally, we tested for scale effects in the data set based on spatial autocorrelograms. We partitioned the data set into spatial lags, which included only those pairwise comparisons within a certain distance class or “bin.” We used Sturges’ rule to objectively determine the optimal number of bins (Sturges 1926), and assessed the correlation between genetic and geographic distances separately for each distance bin (Smouse et al. 1986; Goslee and Urban 2007). If spatial-genetic structure is homogeneous across scales, all bins should show a similar pattern, i.e., the correlogram should show a relatively straight line and little variation in spatial-genetic correlation coefficients. Because we found two distinct spatial scales in the wolverine data set where spatial-genetic correlation coefficients differed significantly (see “Results”), we repeated the spatial and hierarchical variance partitioning for these two distinct scales, thus assessing the influence of scale effects on our inferences.

All statistical analyses were conducted in the R software, using packages *ecodist* (Goslee and Urban 2007) and *hier.part* (Walsh and Mac Nally 2009). When significance tests were needed,  $P$ -values were estimated via 9,999 permutations, and 95% confidence intervals were based on 1,000 bootstraps (Goslee and Urban 2007).

## RESULTS

Our optimization approach produced models that performed well in predicting wolverine genetic structure, with Mantel correlations and amounts of explained variation in the multiple-matrix regressions being at the high end of values reported for individual-based studies (see, e.g., Wang et al. 2008). Our results were consistent across the different genetic distances we used, with all of them being significantly correlated to each other ( $r > 0.93$ ). We therefore only report results based on Miller’s  $D$ , as it gave slightly higher Mantel  $r$ -values than the other genetic distances.

**Model optimization and exclusion of spurious variables.**—All simple Mantel tests were highly significant ( $P = 0.0001$ ), and the highest Mantel correlations obtained for each variable are shown in Table 2. Effective resistances based on circuit theory generally led to higher correlation coefficients than least-cost distances, except for timberline elevation and conifer cover (Table 2; Supplementary Data SD2). Resistance models



**Table 2.**—Highest Mantel correlations obtained for each variable. \*‘‘Best’’ refers to the conversion and effective distance that led to the highest  $r$ -values. CT = effective resistances based on circuit theory; LCD = least-cost distances. Variable abbreviations as in Table 1, *geo* is the geographic (i.e., straight-line) distance. Note that in the partial Mantel tests, a Bonferroni-corrected  $P$ -value of 0.0056 is required for significance at  $\alpha = 0.05$ , meaning that *treecov* and *absdiff* are insignificant.

Variable	Best* conversion	Best* effective distance	Simple Mantel statistic				Partial Mantel statistic			
			$r$ -value	$P$ -value	Lower 95% CI	Upper 95% CI	$r$ -value	$P$ -value	Lower 95% CI	Upper 95% CI
snow	Upper exponential	CT	0.276	0.0001	0.25	0.304	0.238	0.0001	0.199	0.268
roaden	Upper exponential	CT	0.274	0.0001	0.248	0.301	0.189	0.0001	0.164	0.216
hden	Upper exponential	CT	0.261	0.0001	0.237	0.291	0.164	0.0001	0.137	0.185
edge	Lower exponential	CT	0.263	0.0001	0.236	0.287	0.165	0.0001	0.138	0.195
tri	Upper exponential	CT	0.243	0.0001	0.218	0.271	0.111	0.0030	0.078	0.154
timberline	Upper exponential	LCD	0.245	0.0001	0.222	0.273	0.136	0.0002	0.103	0.165
absdiff	Linear	CT	0.226	0.0001	0.202	0.252	0.106	0.0170	0.070	0.130
popden	Upper exponential	CT	0.247	0.0001	0.219	0.269	0.131	0.0001	0.109	0.154
treecov	Lower exponential	LCD	0.209	0.0001	0.189	0.238	0.08	0.0130	0.048	0.099
geo	NA	NA	0.218	0.0001	0.197	0.240	NA	NA	NA	NA

derived from the higher exponential translation led to slightly higher correlation coefficients for most variables (Table 2), but strong effects of the different translations only were apparent in three variables (i.e., road density, timberline, conifer cover; Supplementary Data SD2). In the partial Mantel tests, all variables were significant after Bonferroni correction, with the exception of conifer cover and absolute elevation difference (Table 2). These two variables were therefore dropped from further analyses.

Effective distances based on the remaining seven environmental variables showed no collinearity with each other (i.e., Mantel  $r < 0.7$ ,  $P \geq 0.05$ ), except for population density and housing density (Mantel  $r = 0.74$ ,  $P = 0.0001$ ). We chose to delete population density and retain housing density for further analyses, because the latter resulted in a higher correlation coefficient with genetic distances. Thus, we retained six environmental variables for our final analyses (i.e., elevation—timberline, terrain ruggedness—tri, snow depth—snow, housing density—hden, road density—roaden, and forest edge—edge), plus straight-line geographic distance (*geo*).

**Scale-dependent results.**—When using all pairwise comparisons, the MRM model explained 13.5% of the variation, and was highly significant ( $P < 0.001$ ). The spatial variance partitioning of the full MRM model showed that pure environmental factors had a strong impact on the explained variation (Supplementary Data SD3). About 64.7% of the explained variation can be attributed to pure environmental components (IBR), 23.9% to shared spatial-environmental components, and only 11.4% to pure spatial components (IBD). In terms of individual variable importance, hierarchical partitioning suggested that snow depth, terrain ruggedness, and housing density, best predict genetic distances across the 127 possible models (Supplementary Data SD4; maps of these three most influential variables are shown in Supplementary Data SD5). However, strong scale effects were detected in the genetic structure, as the Mantel correlogram (Fig. 3) suggested positive autocorrelation in the first five distance classes (i.e., up to ~230 km), while negative autocorrelation existed in the last five distance classes (i.e., beyond

distances of ~420 km). In between these distances, there is no significant spatial autocorrelation. Thus, wolverines were genetically more similar than expected under spatially random mating at small spatial scales (up to ~230 km distance), while they were less similar than expected at large spatial scales (beyond ~420 km distance). Analyzing these two spatial scales separately is particularly interesting, because different landscape-genetic relationships could exist within each scale.


Based on these results, we partitioned the data set into two groups: one including only pairwise comparisons belonging to the first five distance bins (‘‘small scale’’), and the other including only those distances belonging to the last five distance bins (‘‘broad scale’’). We then repeated the two variance partitioning analyses for small and broad scale separately.

The spatial variance partitioning analyses suggested substantial differences in the relative importance of spatial and environmental variables between the two scales (Fig. 4). While both MRM models explained similar percentages of variation (14.7% for small scale, and 13.5% for broad scale; Supplementary Data SD3), the combined effect of space and environmental heterogeneity were much more important at larger scales. Pure environmental components (IBR) were responsible for 78.9% of the explained variation at small scales, and 64.6% at broad scales. Pure space (IBD) accounted for 17.1% and 11.4% of the variation at small and broad scale, respectively. Shared components accounted for only 4.1% at fine scales, but for 24% at broad scales.


The relative importance of the different environmental variables also varied with scale (Fig. 5). At the small scale, snow depth was the most important variable, with a 42.9% contribution to the variance explained. Terrain ruggedness contributed slightly more than straight-line geographic distances (i.e., IBD) at this scale (16.6% and 13.2%, respectively), while the contribution of road density was similar to that of geographic distance (13.6% and 13.2%, respectively). All other variables contributed less than geographic distance to the genetic structure at this small scale.

At the broad scale, the contribution of snow depth to explained variance became much weaker (7.4%), while the





**Fig. 3.**—Spatial Mantel correlogram used to infer scale effects. The number and width of the equally sized spatial bins was determined using Sturges' rule ( $N = 15$ , width of bin = 47 km). Black dots indicate Mantel correlation coefficients ( $r$ ) that are significantly different from zero after Bonferroni correction; white dots indicate Mantel  $r$  values not different from zero.




**Fig. 4.**—Spatial variance partitioning for small and broad scales. Shown as percentages of explained variation.

importance of terrain ruggedness increased (24.9%). The most pronounced difference between the two scales occurred for housing density, which contributed only little at the small scale (12.1%), but was the most important variable at the broad scale (54.6%). Final resistance values for long-distance gene flow among wolverine in our study area are shown in Fig. 1, as inferred from the broad-scale model results. To create this final resistance layer, we added the best resistance layers of the six environmental variables, after weighting them by their relative importance at the broad scale (in % contribution to  $R^2$ ; Fig. 5).

## DISCUSSION

Dispersal is a key process in ecology and evolution (Clobert et al. 2012; Travis et al. 2013) and various approaches have been developed to study dispersal in the wild (i.e., Cayuela et al. 2018). Landscape genetic approaches are particularly useful for analyzing environmental influences on effective dispersal, i.e., dispersal leading to gene flow (Broquet and Petit



**Fig. 5.**—Relative variable importance for partial models at two distinct scales based on hierarchical partitioning. Variable importance is given in % contribution of each variable to overall variation explained. The black dotted line shows the contribution of straight-line geographical distance in the full models at the small scale (i.e., 13.2%; see Supplementary Data SD4). The gray dashed line shows the contribution of geographical distance in the full models at the broad scale (i.e., 12.7%; see Supplementary Data SD4).

2009; Robertson et al. 2018). Since dispersal can be impacted by multiple environmental drivers and their relative importance can vary across spatial scales (e.g., Morton et al. 2018), it is important to evaluate the effects of multiple environmental variables in landscape genetic studies and to quantitatively test and account for scale effects on landscape-genetic relationships.

Here we used a multivariate, multiscale approach for a detailed investigation of landscape-genetic relationships in wolverines inhabiting the contiguous United States. Several previous studies have demonstrated that wolverines in the contiguous United States exhibit lower levels of genetic diversity, and higher degrees of genetic structure than their conspecifics in Canada and Alaska (Kyle and Strobeck 2001, 2002; Cegelski

et al. 2003, 2006). While the homogeneous boreal and consistently snowy areas in the north are less fragmented and can support larger effective population sizes of the species (Schwartz et al. 2009), suitable wolverine habitat in the contiguous United States only can be found in several high-elevation mountain ranges, which are separated by low-elevation basins with limited amounts of snow and nonpersistent spring snow cover (Aubry et al. 2007; Inman et al. 2013). However, the comparably low levels of gene flow in the southernmost parts of the species' range sometimes also have been attributed to increased anthropogenic impacts, which potentially have led to additional habitat fragmentation and reduced wolverine densities in this area (Guillot et al. 2005; Cegelski et al. 2006).

Our results suggest that wolverine genetic structure in the region can indeed be predicted by a combination of bioclimatic, anthropogenic, and topographic, landscape characteristics, that these environmental factors are much more important for predicting wolverine gene flow than pure space (IBD), and that the relative influence of these variables is scale-dependent. Deriving this inferential detail only was possible through our multivariate and multiscale analyses, which greatly increased the ecological meaningfulness of our study.

*Methodological considerations for inferring landscape effects on gene flow.*—Using MRM and hierarchical partitioning allowed us to avoid the use of significance testing (e.g., based on Mantel tests) for our final inferences, and instead assess the relative importance of each individual variable in a multivariate context, as recommended by Prunier et al. (2015, 2017). From an ecological standpoint, this is arguably more informative than trying to find the single “best” landscape resistance model for predicting genetic structures, as often is done in landscape genetics. Furthermore, testing and accounting for scale effects proved vital for our analyses and led to a more detailed understanding of the potential processes underlying successful dispersal of wolverine within versus among habitat complexes. For the wolverines in our study area, our multiscale approach enabled us to differentiate between landscape influences on small-scale gene flow (likely reflecting typical dispersal and mating movements within habitat complexes) versus landscape influences on broad-scale gene flow (reflecting long-distance dispersal or multigenerational gene flow among habitat complexes). This kind of inferential detail would not have been derived from the analyses involving all scales, as the model involving all scales could not clearly disentangle the relative importance of individual variables for predicting genetic structure (Supplementary Data SD4). Our analytical framework thus can help to disentangle environmental impacts on the different processes leading to gene flow and results in a biologically meaningful distinction between habitat suitability and landscape resistance. These findings allow us to discuss possible causes of detected landscape-genetic relationships at fine and broad spatial scales.

*Fine-scale predictors of wolverine gene flow.*—In our fine-scale analyses (i.e., up to ~230 km), pure spatial and pure environmental components explained most of the variation in genetic distances. This scale is well within reported wolverine

dispersal distances of, e.g., up to 170–178 km (Vangen et al. 2001; Inman et al. 2012a). Accordingly, the strong impact of pure space at this small scale likely is not due to an actual, physical dispersal limitation (classical IBD—Wright 1943), but probably is caused by wolverine space-use and mating patterns (Dalerum et al. 2007). Wolverines show the typical space-use and polygamous mating patterns of mustelids, with male home ranges overlapping those of several females (Banci 1994; Dalerum et al. 2007; Persson et al. 2010). Also, wolverines may establish their home ranges in proximity of their natal range (Hedmark et al. 2007; Bischof et al. 2016; Aronsson and Persson 2018). These behavioral characteristics can lead to strong spatial-genetic correlations at fine spatial scales, because related individuals tend to be closer geographically than unrelated individuals (see, e.g., Wang et al. 2017; Kristensen et al. 2018, for other species). Because these are endogenous (i.e., landscape-independent) processes, they explain the relatively high percentage of pure spatial components in our small-scale analysis, despite the high mobility of the species.

In terms of landscape predictors of spatial-genetic structure, our small-scale results are congruent with those of Schwartz et al. (2009), who used data on the presence or absence of persistent spring snow cover to show that areas lacking such persistent snow cover have a resistance to gene flow approximately 20 times higher than areas with persistent spring snow cover. Because our weighted resistance approach for modeling resistance values required us to work with continuous variables, we used snow depth data in our study instead of categorical persistent spring snow cover data. Despite this difference in modeling snow-related landscape resistances, our results support the conclusion that deep, persistent snow cover predicts successful wolverine dispersal and resulting genetic structures. Indeed, snow was the most important predictor for wolverine gene flow across all scales, and especially up to distances of ca. 230 km. Most likely, gene flow at this scale is primarily determined by mating movements and natal dispersal of wolverine offspring, and these processes largely occur within primary habitat complexes, i.e., within mountain ranges. These complexes are typical wolverine habitats and often characterized by low housing densities and high topographic heterogeneity (e.g., May et al. 2006; Inman et al. 2013). Thus, at this spatial scale, wolverine gene flow cannot be predicted very well by housing and topography, as they simply are not “limiting factors” (sensu Cushman et al. 2013b) of dispersal within typical wolverine habitats. Instead, successful wolverine dispersal mainly is determined by the availability of high-quality breeding habitat, with deep, persistent spring snow cover as a very important predictor (Magoun and Copeland 1998; Copeland et al. 2010).

Other environmental variables that we investigated (i.e., conifer cover, forest edge habitat, elevation, and road densities) showed statistically significant, but relatively weak, relationships with genetic distances. However, these results do not contradict previous studies that identified these variables as important for wolverine occurrence and movement (Aubry et al. 2007; Copeland et al. 2007; Inman et al. 2013; Webb et al.

2016; Kortello et al. 2019; Sawaya et al. 2019), because these studies were conducted more locally or in different regions.

**Broad-scale predictors of wolverine gene flow.**—At scales beyond typical wolverine dispersal distances (i.e., > 420 km), housing density and terrain ruggedness were the best predictors of spatial-genetic structures in wolverines. These spatial distances likely are only covered by multigenerational gene flow or occasional long-distance dispersal events (Gardner et al. 1986; Moriarty et al. 2009). For example, an exceptional straight-line dispersal distance of 826 km for an adult male was reported by Packila et al. (2017), and male dispersal up to 500 km was detected using genetic means by Flagstad et al. (2004). To maintain connectivity over such large distances, wolverines eventually have to disperse outside of their typical, mountainous habitat, and potentially cross areas that are characterized by higher housing densities and areas that are topographically more homogeneous (e.g., Packila et al. 2017). Our results suggest that these factors strongly influence population genetic structure in wolverines at broad spatial scales. Given that anthropogenic features can affect wolverine space-use behavior, including habitat selection, at large spatial scales (e.g., Rowland et al. 2003; May et al. 2006; Stewart et al. 2016; Heim et al. 2017), it seems plausible that effective wolverine dispersal is higher across areas with low human housing density and associated human presence. In contrast, it is unlikely that wolverine dispersal actually is limited by topographic homogeneity per se, because wolverines show high levels of genetic connectivity in areas without much topographic variation (i.e., taiga and tundra—Wilson et al. 2000; Kyle and Strobeck 2001, 2002). Instead, the broad-scale relationship between genetic connectivity and topographic ruggedness observed in our study might be caused by comparably little human influences and associated infrastructures in areas with high topographic heterogeneity. For example, wolverines in Canada preferred more rugged terrain with less human development (Fisher et al. 2013), just as wolverines in Norway avoided areas with infrastructural development (May et al. 2006).

The increasing availability of suitable sites for food caches in more rugged terrain is an alternative or additional explanation for the importance of terrain ruggedness in predicting spatial-genetic structure in wolverines. Wolverines often cache food in cold, structured sites protected from competitors, such as bacteria, insects, and other scavengers, for example, under downed logs or large boulders (Magoun and Copeland 1998; Inman et al. 2012b). Inman et al. (2013) showed that terrain ruggedness was a strong predictor of wolverine resource selection and hypothesized that areas with increased terrain ruggedness offer more cliffs, boulders, and talus areas, and generally more topographic habitat structure than areas with low ruggedness, and showed that terrain ruggedness was indeed a strong predictor of wolverine resource selection. Thus, it seems possible that long-distance gene flow across topographically more variable areas is higher than across flat areas, because more rugged areas offer better caching opportunities, hence are more suitable for the species.

Overall, the relationship between terrain ruggedness and genetic connectivity detected in our study warrants further investigation such that potential mechanisms behind this landscape-genetic relationship can be clarified. Nevertheless, based on our results we can conclude that terrain ruggedness is a strong predictor of wolverine genetic structure at broad spatial scales, although not as strong as housing density.

**Conclusions for wolverine dispersal and population connectivity.**—Our study provides insights into gene flow among wolverine at different spatial scales, and improves our understanding of dispersal ecology and functional connectivity in the species. Our methodological approach allowed us to analyze effective dispersal (i.e., gene flow) in the species at distinct spatial scales, thereby enabling us to detect the different environmental predictors of genetic structure at small and broad scales. While previous studies had already shown the strong association between gene flow and deep, persistent snow (Schwartz et al. 2009), or suggested anthropogenic and topographic effects on population genetic structure without explicitly quantifying them (Cegelski et al. 2003, 2006; Guillot et al. 2005), we demonstrated the importance of terrain ruggedness and housing density for predicting broad-scale wolverine gene flow. Had we only analyzed the all-scale model, we would have concluded that snow depth is the best variable for predicting genetic structure. While this finding is not incorrect, it provides an incomplete picture, because deep snow in the region only occurs within typical wolverine habitat, where most gene flow occurs across usual dispersal and mating distances. In contrast, low housing density and terrain ruggedness occur outside of wolverine habitat and individuals only have to move through these areas during occasional long-distance dispersal. While not common, long-distance dispersal among habitat complexes is vital for maintaining meta-population viability and species persistence (Trakhtenbrot et al. 2005; Heinz et al. 2006). Our finding that housing density best predicts gene flow across broad spatial scales thus is of particular relevance for maintaining and improving connectivity among wolverine habitats. Specifically, our results suggest that connectivity among wolverine will be particularly sensitive to housing developments and other human impacts in rugged areas located between typical wolverine habitat. Even if these areas are not usually inhabited by wolverines, they are potentially used during dispersal and can therefore offer crucial pathways for gene flow across broad spatial scales. Efforts to identify optimal corridors among wolverine habitats in the region should accordingly consider housing density and terrain ruggedness as predictive variables of successful wolverine dispersal.

Our multivariate, multiscale analyses led to a more detailed understanding of wolverine dispersal in a heterogeneous environment. We suggest that the approach is generally useful for landscape genetic research in mammals, particularly for large scale studies involving highly mobile species, as such studies likely will capture gene flow at different spatial scales, and both within and between typical habitat of the study species.



## ACKNOWLEDGMENTS

During this work, NB was supported by a scholarship of the German-American Fulbright program and by a University of Idaho (UI) Doctoral Fellowship. LPW was supported by the College of Natural Resources at UI. We thank the Montana Department of Fish, Wildlife and Parks for sample collection, and Mark T. Packila for help with acquisition and preparation of landscape data.

## SUPPLEMENTARY DATA

Supplementary data are available at *Journal of Mammalogy* online.

**Supplementary Data SD1.**—Supplementary information on digital layers used for resistance modeling.

**Supplementary Data SD2.**—Mantel correlations obtained during model optimization.

**Supplementary Data SD3.**—Results for spatial variance partitioning for the global (all-scale), small-scale, and broad-scale models.

**Supplementary Data SD4.**—Relative variable importance for the full and partial models at three spatial scales based on hierarchical partitioning.

**Supplementary Data SD5.**—Maps showing the distribution of the three variables that best predicted genetic structure of wolverines in the study area.

## LITERATURE CITED

- ABRAHMS, B., S. C. SAWYER, N. R. JORDAN, J. W. MCNUTT, A. M. WILSON, AND J. BRASHARES. 2017. Does wildlife resource selection accurately inform corridor conservation? *Journal of Applied Ecology* 54:412–422.
- ADRIAENSEN, F., ET AL. 2003. The application of 'least-cost' modelling as a functional landscape model. *Landscape and Urban Planning* 64:233–247.
- ARONSSON, M., AND J. PERSSON. 2018. Female breeding dispersal in wolverines, a solitary carnivore with high territorial fidelity. *European Journal of Wildlife Research* 64:7.
- AUBRY, K. B., K. S. MCKELVEY, AND J. P. COPELAND. 2007. Distribution and broadscale habitat relations of the wolverine in the contiguous United States. *Journal of Wildlife Management* 71:2147–2158.
- AVON, C., AND L. BERGÈS. 2016. Prioritization of habitat patches for landscape connectivity conservation differs between least-cost and resistance distances. *Landscape Ecology* 31:1551–1565.
- BALKENHOL, N., L. P. WAITS, AND R. DEZZANI. 2009. Statistical approaches in landscape genetics: an evaluation of methods for linking landscape and genetic data. *Ecography* 32:818–830.
- BANCI, V. 1994. Wolverine. Pp. 99–127 in *The scientific basis for conserving forest carnivores: American marten, fisher, lynx, and wolverine in the western United States* (L. F. Ruggiero, K. B. Aubry, S. W. Buskirk, L. J. Lyon, and W. J. Zielinski, eds.). USDA Forest Service General Technical Report, RM-254. Fort Collins, Colorado.
- BARRETT, A. P. 2003. National Operational Hydrologic Remote Sensing Center Snow Data Assimilation System (SNODAS). National Snow and Ice Data Center Special Report 11. Boulder, CO.
- BARTOŃ, K. A., T. ZWIJACZ-KOZICA, F. ZIĘBA, A. SERGIEL, AND N. SELVA. 2019. Bears without borders: long-distance movement in human-dominated landscapes. *Global Ecology and Conservation* 17:e00541.
- BISCHOF, R., E. R. GREGERSEN, H. BRØSETH, H. ELLEGREN, AND Ø. FLAGSTAD. 2016. Noninvasive genetic sampling reveals intrasex territoriality in wolverines. *Ecology and Evolution* 6:1527–1536.
- BORCARD, D., P. LEGENDRE, AND P. DRAPEAU. 1992. Partialling out the spatial component of ecological variation. *Ecology* 73:1045–1055.
- BROCK, B. L., ET AL. 2007. Broad-scale wolverine habitat in the conterminous Rocky Mountain states. Pp. 21–53 in *Greater Yellowstone Wolverine Program. Cumulative Report May 2007* (R. M. Inman, M. L. Packila, K. H. Inman, R. S. Spence, and D. McCauley, eds.). Wildlife Conservation Society, North America Program. Ennis, Montana.
- BROCK, B. L., AND R. M. INMAN. 2006. Use of latitude-adjusted elevation in broad-scale species distribution models. *Intermountain Journal of Sciences* 12:12–17.
- BROQUET, T., AND E. J. PETIT. 2009. Molecular estimation of dispersal for ecology and population genetics. *Annual Review of Ecology, Evolution, and Systematics* 40:193–216.
- CARROLL, C., R. F. NOSS, AND P. C. PAQUET. 2001. Carnivores as focal species for conservation planning in the Rocky Mountain region. *Ecological Applications* 11:961–980.
- CARROLL, C., D. J. ROHLE, Y.-W. LI, B. HARTL, M. K. PHILLIPS, AND R. F. NOSS. 2015. Connectivity conservation and endangered species recovery: a study in the challenges of defining conservation-reliant species. *Conservation Letters* 8:132–138.
- CAYUELA, H., ET AL. 2018. Demographic and genetic approaches to study dispersal in wild animal populations: a methodological review. *Molecular Ecology* 27:3976–4010.
- CEGELSKI, C. C., L. P. WAITS, AND N. J. ANDERSON. 2003. Assessing population structure and gene flow in Montana wolverines (*Gulo gulo*) using assignment-based approaches. *Molecular Ecology* 12:2907–2918.
- CEGELSKI, C. C., L. WAITS, N. J. ANDERSON, O. FLAGSTAD, C. STROBECK, AND C. J. KYLE. 2006. Genetic diversity and population structure of wolverine (*Gulo gulo*) populations at the southern edge of their current distribution in North America with implications for genetic viability. *Conservation Genetics* 7:197–211.
- CHEVAN, A., AND M. SUTHERLAND. 1991. Hierarchical partitioning. *The American Statistician* 45:90–96.
- CLOBERT, J., M. BAGUETTE, T. G. BENTON, AND J. M. BULLOCK (EDS.). 2012. *Dispersal ecology and evolution*. Oxford University Press, Oxford, United Kingdom.
- COPELAND, J. P. 1996. *Biology of the wolverine in central Idaho*. M.S. thesis, University of Idaho. Moscow.
- COPELAND, J. P., ET AL. 2007. Seasonal habitat associations of the wolverine in Central Idaho. *Journal of Wildlife Management* 71:2201–2212.
- COPELAND, J. P., ET AL. 2010. The bioclimatic envelope of the wolverine: do environmental constraints limit their geographic distribution? *Canadian Journal of Zoology* 88:233–246.
- CUSHMAN, S. A., K. S. MCKELVEY, J. HAYDEN, AND M. K. SCHWARTZ. 2006. Gene flow in complex landscapes: testing multiple hypotheses with causal modeling. *The American Naturalist* 168:486–499.
- CUSHMAN, S. A., A. J. SHIRK, AND E. L. LANDGUTH. 2013b. Landscape genetics and limiting factors. *Conservation Genetics* 14:263–274.



- CUSHMAN, S. A., T. N. WASSERMAN, E. L. LANDGUTH, AND A. J. SHIRK. 2013a. Re-evaluating causal modeling with Mantel tests in landscape genetics. *Diversity* 5:51–72.
- DALERUM, F., J. LOXTERMAN, B. SHULTS, K. KUNKEL, AND J. A. COOK. 2007. Sex-specific dispersal patterns of wolverines: insights from microsatellite markers. *Journal of Mammalogy* 88:793–800.
- DORMANN, C. F., ET AL. 2013. Collinearity: a review of methods to deal with it and a simulation study evaluating their performance. *Ecography* 36:27–46.
- EDELMANN, F. B., AND J. P. COPELAND. 1999. Wolverine distributions in the northwestern United States and a survey in the Seven Devils mountains of Idaho. *Northwest Science* 73:295–300.
- FISHER, J. T., ET AL. 2013. Wolverines (*Gulo gulo luscus*) on the Rocky Mountain slopes: natural heterogeneity and landscape alteration as predictors of distribution. *Canadian Journal of Zoology* 91:706–716.
- FLAGSTAD, Ø., ET AL. 2004. Colonization history and noninvasive monitoring of a reestablished wolverine population. *Conservation Biology* 18:676–688.
- FORTUNE, L., J. P. PRUNIER, I. PAZ-VINAS, G. LOOT, C. VEYSSIÈRE, AND S. BLANCHET. 2018. Inferring causalities in landscape genetics: an extension of Wright's causal modeling to distance matrices. *The American Naturalist* 191:491–508.
- FRANCKOWIAK, R. P., ET AL. 2017. Model selection with multiple regression on distance matrices leads to incorrect inferences. *PLoS ONE* 12:e0175194.
- GARDNER, C. L., W. B. BALLARD, AND R. H. JESSUP. 1986. Long distance movement by an adult wolverine. *Journal of Mammalogy* 67:603.
- GOSLEE, S. C., AND D. L. URBAN. 2007. The ecodist package for dissimilarity-based analysis of ecological data. *Journal of Statistical Software* 22:7.
- GUILLOT, G., A. ESTOUP, F. MORTIER, AND J. F. COSSON. 2005. A spatial statistical model for landscape genetics. *Genetics* 170:1261–1280.
- HARDY, O. J., AND X. VEKEMANS. 2002. Spagedi: a versatile computer program to analyse spatial genetic structure at the individual or population levels. *Molecular Ecology Notes* 2:618–620.
- HEDMARK, E., J. PERSSON, P. SEGESTROEM, A. LANDA, AND H. ELLEGREN. 2007. Paternity and mating system in wolverines *Gulo gulo*. *Wildlife Biology* 13:13–30.
- HEIM, N., J. T. FISHER, A. CLEVINGER, J. PACZKOWSKI, AND J. VOLPE. 2017. Cumulative effects of climate and landscape change drive spatial distribution of Rocky Mountain wolverine (*Gulo gulo* L.). *Ecology and Evolution* 7:8903–8914.
- HEINZ, S. K., C. WISSEL, AND K. FRANK. 2006. The viability of metapopulations: individual dispersal behaviour matters. *Landscape Ecology* 21:77–89.
- HOMER, C., C. HUANG, L. LANG, B. WYLIE, AND M. COAN. 2001. Development of a 2001 national land-cover database for the United States. *Photogrammetric Engineering and Remote Sensing* 70:829–840.
- HORNOKER, M. G., AND H. S. HASH. 1981. Ecology of the wolverine in northwestern Montana. *Canadian Journal of Zoology* 59:1286–1301.
- INMAN, R. M., ET AL. 2012a. Spatial ecology of wolverines at the southern periphery of distribution. *Journal of Wildlife Management* 76:778–792.
- INMAN, R. M., ET AL. 2013. Developing priorities for metapopulation conservation at the landscape scale: wolverines in the western United States. *Biological Conservation* 166:276–286.
- INMAN, R. M., A. J. MAGOUN, J. PERSSON, AND J. MATTISSON. 2012b. The wolverine's niche: linking reproductive chronology, caching, competition, and climate. *Journal of Mammalogy* 93:634–644.
- JENKINS, D. G., ET AL. 2010. A meta-analysis of isolation by distance: relic or reference standard for landscape genetics? *Ecography* 33:315–320.
- KEELEY, A. T. H., P. BEIER, AND J. W. GAGNON. 2016. Estimating landscape resistance from habitat suitability: effects of data source and nonlinearities. *Landscape Ecology* 31:2151–2162.
- KORTELO, A., D. HAUSLEITNER, AND G. MOWAT. 2019. Mechanisms influencing the winter distribution of wolverine *Gulo gulo luscus* in the southern Columbia Mountains, Canada. *Wildlife Biology* 2019:wlb.00480.
- KRISTENSEN, T. V., ET AL. 2018. Spatial genetic structure in American black bears (*Ursus americanus*): female philopatry is variable and related to population history. *Heredity* 120:329–341.
- KYLE, C. J., AND C. STROBECK. 2001. Genetic structure of North American wolverine (*Gulo gulo*) populations. *Molecular Ecology* 10:337–347.
- KYLE, C. J., AND C. STROBECK. 2002. Connectivity of peripheral and core populations of North American wolverines. *Journal of Mammalogy* 83:1141–1150.
- LEGENDRE, P. 2000. Comparison of permutation methods for the partial correlations and partial Mantel tests. *Journal of Statistical Computation and Simulation* 67:37–73.
- LICHSTEIN, J. W. 2007. Multiple regression on distance matrices: a multivariate spatial analysis tool. *Plant Ecology* 188:117–131.
- MAC NALLY, R. 2000. Regression and model-building in conservation biology, biogeography and ecology: the distinction between - and reconciliation of - 'predictive' and 'explanatory' models. *Biodiversity and Conservation* 9:655–671.
- MAC NALLY, R. 2009. Hierarchical partitioning as an interpretative tool in multivariate inference. *Austral Ecology* 21:224–228.
- MAGOUN, A. J., AND J. P. COPELAND. 1998. Characteristics of wolverine reproductive den sites. *Journal of Wildlife Management* 62:1313–1320.
- MANTEL, N. 1967. The detection of disease clustering and a generalized regression approach. *Cancer Research* 27:209–220.
- MARROTTE, R. R., AND J. BOWMAN. 2017. The relationship between least-cost and resistance distance. *PLoS ONE* 12:e0174212.
- MATEO-SÁNCHEZ, M., N. BALKENHOL, S. A. CUSHMAN, T. PÉREZ, P. DOMÍNGUEZ, AND S. SAURA. 2015. A comparative framework to infer landscape effects on population genetic structure: are habitat suitability models effective in explaining gene flow? *Landscape Ecology* 8:1405–1420.
- MAY, R., A. LANDA, J. VAN DIJK, D. C. LINNELL, AND R. ANDERSEN. 2006. Impact of infrastructure on habitat selection of wolverines *Gulo gulo*. *Wildlife Biology* 12:285–295.
- MCCLURE, M. L., A. J. HANSEN, AND R. M. INMAN. 2016. Connecting models to movements: testing connectivity model predictions against empirical migration and dispersal. *Landscape Ecology* 31:1419–1432.
- MCKELVEY, K. S., ET AL. 2011. Climate change predicted to shift wolverine distributions, connectivity, and dispersal corridors. *Ecological Applications* 21:2882–2897.
- MCKELVEY, K. S., ET AL. 2014. Recovery of wolverines in the western United States: recent extirpation and recolonization or range retraction and expansion. *The Journal of Wildlife Management* 78:325–334.
- MCRAE, B. H. 2006. Isolation by resistance. *Evolution* 60:1551–1561.
- MCRAE, B. H., AND P. BEIER. 2007. Circuit theory predicts gene flow in plant and animal populations. *Proceedings of the*

- National Academy of Sciences of the United States of America 104:19885–19890.
- McRAE, B. H., AND V. B. SHAH. 2013. Circuitscape 3.4 user guide. The Nature Conservancy, Arlington, Virginia.
- MILLER, M. P. 2005. Alleles in space: computer software for the joint analysis of interindividual spatial and genetic information. *Journal of Heredity* 96:722–724.
- MORIARTY, K. M., ET AL. 2009. Wolverine confirmation in California after nearly a century: native or long-distance immigrant? *Northwest Science* 83:154–162.
- MORTON, E. R., ET AL. 2018. Dispersal: a matter of scale. *Ecology* 99:938–946.
- MURRAY, K., AND M. M. CONNER. 2009. Methods to quantify variable importance: implications for the analysis of noisy ecological data. *Ecology* 90:348–355.
- NEI, M., F. TAJIMA, AND Y. TATENO. 1983. Accuracy of estimated phylogenetic trees from molecular data. II. Gene frequency data. *Journal of Molecular Evolution* 19:153–170.
- OREGON WILD. 2019. Oregon's lonely wolverine. <https://oregonwild.org/about/blog/oregons-lonely-wolverine>. Accessed 9 March 2020.
- PACKILA, M. L., M. D. RILEY, R. S. SPENCE, AND R. M. INMAN. 2017. Long-distance wolverine dispersal from Wyoming to historic range in Colorado. *Northwest Science* 91:399–407.
- PEAKALL, R., AND P. E. SMOUSE. 2005. GenA1Ex 6: genetic analysis in Excel. Population genetic software for teaching and research. *Molecular Ecology Notes* 6:288–295.
- PERSSON, J., P. WEDHOLM, AND P. SEGERSTRÖM. 2010. Space-use and territoriality of wolverines (*Gulo gulo*) in northern Scandinavia. *European Journal of Wildlife Research* 56:49–57.
- PRUNIER, J. G., M. COLYN, X. LEGENDRE, AND M.-C. FLAMAND. 2017. Regression commonality analyses on hierarchical genetic distances. *Ecography* 40:1412–1425.
- PRUNIER, J. G., M. COLYN, X. LEGENDRE, K. F. NIMON, AND M. C. FLAMAND. 2015. Multicollinearity in spatial genetics: separating the wheat from the chaff using commonality analyses. *Molecular Ecology* 24:263–283.
- RADELOFF, V. C., R. B. HAMMER, S. I. STEWART, J. S. FRIED, S. S. HOLCOMB, AND J. F. MCKEEFRY. 2005. The wildland urban interface in the United States. *Ecological Applications* 15:799–805.
- RAY, N. 2005. PATHMATRIX: a GIS tool to compute effective distances among samples. *Molecular Ecology Notes* 5:177–180.
- RILEY, S. J., S. D. DEGLORIA, AND R. ELLIOT. 1999. A terrain ruggedness index that quantifies topographic heterogeneity. *Intermountain Journal of Sciences* 5:23–27.
- ROBERTSON, E. P., ET AL. 2018. Isolating the roles of movement and reproduction on effective connectivity alters conservation priorities for an endangered bird. *Proceedings of the National Academy of Sciences of the United States of America* 115:8591–8596.
- ROFFLER, G. H., ET AL. 2016. Identification of landscape features influencing gene flow: how useful are habitat selection models? *Evolutionary Applications* 9:805–817.
- ROUSSET, F. 2000. Genetic differentiation between individuals. *Journal of Evolutionary Biology* 13:58–62.
- ROWLAND, M. M., M. J. WISDOM, D. H. JOHNSON, B. C. WALES, J. P. COPELAND, AND F. B. EDELMANN. 2003. Evaluation of landscape models for wolverines in the interior Northwest, United States of America. *Journal of Mammalogy* 84:92–105.
- SAWAYA, M. A., A. P. CLEVINGER, AND M. K. SCHWARTZ. 2019. Demographic fragmentation of a protected wolverine population bisected by a major transportation corridor. *Biological Conservation* 236:16–625.
- SHIRK, A. J., E. L. LANDGUTH, AND S. A. CUSHMAN. 2017. A comparison of individual-based genetic distance metrics for landscape genetics. *Molecular Ecology Resources* 17:1308–1317.
- SHIRK, A. J., E. L. LANDGUTH, AND S. A. CUSHMAN. 2018. A comparison of regression methods for model selection in individual-based landscape genetic analysis. *Molecular Ecology Resources* 18:55–67.
- SCHWARTZ, M. K., ET AL. 2007. Inferring geographic isolation of wolverines in California using historical DNA. *Journal of Wildlife Management* 71:2170–2179.
- SCHWARTZ, M. K., ET AL. 2009. Wolverine gene flow across a narrow climatic niche. *Ecology* 90:3222–3232.
- SHIRK, A. J., D. O. WALLIN, S. A. CUSHMAN, C. G. RICE, AND K. I. WARHEIT. 2010. Inferring landscape effects on gene flow: a new model selection framework. *Molecular Ecology* 19:3603–3619.
- SINGLETON, P. H., W. L. GAINES, AND J. F. LEHMKUHL. 2002. Landscape permeability for large carnivores in Washington: a Geographic Information System weighted-distance and least-cost corridor assessment. USDA Research Paper PNW-RP-549. Pacific Northwest Research Station, Portland, Oregon.
- SMOUSE, P. E., J. C. LONG, AND R. R. SOKAL. 1986. Multiple regression and correlation extensions of the Mantel test of matrix correspondence. *Systematic Zoology* 35:627–632.
- SMOUSE, P. E., AND R. PEAKALL. 1999. Spatial autocorrelation analysis of individual multiallele and multilocus genetic structure. *Heredity* 82:561–573.
- SPEAR, S. F., N. BALKENHOL, M. J. FORTIN, B. H. McRAE, AND K. SCRIBNER. 2010. Use of resistance surfaces for landscape genetic studies: considerations for parameterization and analysis. *Molecular Ecology* 19:3576–3591.
- SPEAR, S. F., S. A. CUSHMAN, AND B. H. McRAE. 2016. Resistance surface modeling in landscape genetics. Pp. 129–148 in *Landscape genetics: concepts, methods, applications* (N. Balkenhol, S. A. Cushman, A. Storfer, and L. Waits, eds.). Wiley-Blackwell, Oxford, United Kingdom.
- SQUIRES, J. R., D. H. PLETCHER, T. J. ULIZIO, AND L. F. RUGGIERO. 2006. The association between landscape features and transportation corridors on movements and habitat-use patterns of wolverines. Final Report, MDT Project #8171. Montana Department of Transportation, Helena.
- STEWART, F. E., N. A. HEIM, A. P. CLEVINGER, J. PACZKOWSKI, J. P. VOLPE, AND J. T. FISHER. 2016. Wolverine behavior varies spatially with anthropogenic footprint: implications for conservation and inferences about declines. *Ecology and Evolution* 6:1493–1503.
- STORFER, A., ET AL. 2007. Putting the “landscape” in landscape genetics. *Heredity* 98:128–142.
- STURGES, H. 1926. The choice of a class-interval. *Journal of the American Statistical Association* 21:65–66.
- SUTHERLAND, G. D., A. S. HARESTAD, K. PRICE, AND K. P. LERTZMAN. 2000. Scaling of natal dispersal distances in terrestrial birds and mammals. *Conservation Ecology* 4:16.
- TRAKHTENBROT, A., R. NATHAN, G. PERRY, AND D. M. RICHARDSON. 2005. The importance of long-distance dispersal in biodiversity conservation. *Diversity and Distributions* 11:173–181.
- TRAVIS, J. M. J., ET AL. 2013. Dispersal and species' responses to climate change. *Oikos* 122:1532–1540.
- VANGEN, K. M., J. PERSSON, A. LANDA, R. ANDERSEN, AND P. SEGERSTRÖM. 2001. Characteristics of dispersal in wolverines. *Canadian Journal of Zoology* 79:1641–1649.

- WAGNER, H. H. 2004. Direct multiscale ordination with canonical correspondence analysis. *Ecology* 85:342–351.
- WAGNER, H. H., AND M.-J. FORTIN. 2005. Spatial analysis of landscapes: concepts and statistics. *Ecology* 86:1975–1987.
- WAGNER, H. H., AND M.-J. FORTIN. 2016. Basics of spatial data analysis: linking landscape and genetic data for landscape genetic studies. Pp. 77–98 in *Landscape genetics: concepts, methods, applications* (N. Balkenhol, S. A. Cushman, A. Storfer, and L. Waits, eds.). Wiley-Blackwell. Oxford, United Kingdom.
- WAITS, L. P., S. A. CUSHMAN, AND S. F. SPEAR. 2016. Applications of landscape genetics to connectivity research in terrestrial animals. Pp. 199–219 in *Landscape genetics: concepts, methods, applications* (N. Balkenhol, S. A. Cushman, A. Storfer, and L. Waits, eds.). Wiley-Blackwell. Oxford, United Kingdom.
- WALSH, C., AND R. MAC NALLY. 2009. The hier.part package for R. <http://cran.r-project.org/web/packages/hier.part/index.html>. Accessed 9 March 2020.
- WALTON, Z., ET AL. 2018. Long-distance dispersal in red foxes *Vulpes vulpes* revealed by GPS tracking. *European Journal of Wildlife Research* 64:64.
- WANG, I. J. 2013. Examining the full effects of landscape heterogeneity on spatial genetic variation: a multiple matrix regression approach for quantifying geographic and ecological isolation. *Evolution* 67:3403–3411.
- WANG, G., W. LIU, Y. WANG, X. WAN, AND W. ZHONG. 2017. Restricted dispersal determines fine-scale spatial genetic structure of Mongolian gerbils. *Current Zoology* 63:687–691.
- WANG, Y.-H., K.-C. YANG, C. L. GRIDGMAN, AND L.-K. LIN. 2008. Habitat suitability modelling to correlate gene flow with landscape connectivity. *Landscape Ecology* 23:989–1000.
- WASSERMAN, T. N., S. A. CUSHMAN, M. K. SCHWARTZ, AND D. O. WALLIN. 2010. Spatial scaling and multi-model inference in landscape genetics: *Martes americana* in northern Idaho. *Landscape Ecology* 25:1601–1612.
- WEBB, S. M., ET AL. 2016. Distribution of female wolverines relative to snow cover, Alberta, Canada. *The Journal of Wildlife Management* 80:1461–1470.
- WHITMEE, S., AND C. D. L. ORME. 2013. Predicting dispersal distance in mammals: a trait-based approach. *Journal of Animal Ecology* 82:211–221.
- WILSON, G. M., R. A. VAN DEN BUSSCHE, P. K. KENNEDY, A. GUNN, AND K. POOLE. 2000. Genetic variability of wolverines (*Gulo gulo*) from the Northwest Territories, Canada: conservation implications. *Journal of Mammalogy* 81:186–196.
- WRIGHT, S. 1943. Isolation by distance. *Genetics* 28:114–138.
- ZELLER, K. A., K. MCGARIGAL, AND A. R. WHITELEY. 2012. Estimating landscape resistance to movement: a review. *Landscape Ecology* 27:777–797.

Submitted 23 September 2019. Accepted 16 March 2020.

Associate Editor was Antoinette Piaggio