

## **Lin28a Is Dormant, Functional, and Dispensable During Mouse Oocyte-to-Embryo Transition 1**

Authors: Flemr, Matyas, Moravec, Martin, Libova, Veronika, Sedlacek, Radislav, and Svoboda, Petr

Source: Biology of Reproduction, 90(6)

Published By: Society for the Study of Reproduction

URL: <https://doi.org/10.1095/biolreprod.114.118703>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

# ***Lin28a* Is Dormant, Functional, and Dispensable During Mouse Oocyte-to-Embryo Transition<sup>1</sup>**

Matyas Flemr, Martin Moravec, Veronika Libova, Radislav Sedlacek, and Petr Svoboda<sup>2</sup>

*Institute of Molecular Genetics of the Academy of Sciences of the Czech Republic, Prague, Czech Republic*

## **ABSTRACT**

The oocyte-to-embryo transition (OET) denotes transformation of a highly differentiated oocyte into totipotent blastomeres of the early mammalian embryo. OET depends exclusively on maternal RNAs and proteins accumulated during oocyte growth, which implies importance of post-transcriptional control of gene expression. OET includes replacement of abundant maternal microRNAs (miRNAs), enriched also in differentiated cells and exemplified by the Let-7 family, with embryonic miRNAs common in pluripotent stem cells (the miR-290 family in the mouse). *Lin28a* and its homolog *Lin28b* encode RNA-binding proteins, which interfere with Let-7 maturation and facilitate reprogramming of induced pluripotent stem cells. Both *Lin28a* and *Lin28b* transcripts are abundant in mouse oocytes. To test the role of maternal expression of *Lin28a* and *Lin28b* during oocyte-to-zygote reprogramming, we generated mice with oocyte-specific knockdown of both genes by using transgenic RNA interference. *Lin28a* and *Lin28b* are dispensable during oocyte growth because their knockdown has no effect on Let-7a levels in fully grown germinal vesicle (GV)-intact oocytes. Furthermore, transgenic females were fertile and produced healthy offspring, and their overall breeding performance was comparable to that of wild-type mice. At the same time, 2-cell embryos derived from transgenic females showed up-regulation of mature Let-7, suggesting that maternally provided LIN28A and LIN28B function during zygotic genome activation. Consistent with this conclusion is increased translation of *Lin28a* transcripts upon resumption of meiosis. Our data imply dual repression of Let-7 during OET in the mouse model, the selective suppression of Let-7 biogenesis by *Lin28* homologs superimposed on previously reported global suppression of miRNA activity.

*early development, genome activation, guinea pigs, Let-7, LIN28, mice, microRNA, oocyte, rats, RNAi, rodents, transgenic/knockout model, voles, zygote*

## **INTRODUCTION**

Successful zygotic genome activation is entirely dependent on cytoplasmic factors, as demonstrated by cloning using

somatic nuclear transfer [1]. Post-transcriptional mechanisms engaging mRNAs and proteins produced in the oocyte during oogenesis are thus essential for establishment of correct zygotic gene expression. Oocyte maturation and fertilization induce major changes in translation and stability of cohorts of maternal transcripts. The selection of maternal mRNAs for different post-transcriptional events has been shown to be driven by *cis*-acting elements located in 3'-untranslated regions (3'UTRs), and *trans*-acting factors that bind to them [2–4]. The number, position, and combination of *cis*-acting elements and the presence of *trans*-acting factors offer a complex combinatorial system controlling mRNA localization, stability, and translation [3].

Dormancy is a classical example of post-transcriptional regulation of maternal mRNAs, where certain deadenylated and translationally inactive mRNAs are recruited for translation in response to developmental cues. Translation of dormant mRNAs is induced through cytoplasmic polyadenylation, which is mediated by a *cis*-acting 3'UTR element called cytoplasmic polyadenylation element (CPE) and its binding protein (CPEB) (reviewed in [5, 6]). CPEB is one of many RNA binding proteins which recognize either sequence motifs or secondary structures within 3'UTRs and regulate mRNA metabolism.

Another class of *trans*-acting factors binding *cis*-acting 3'UTR elements is represented by microRNAs (miRNAs), which act as sequence-specific guides, usually imperfectly basepairing with 3'UTR binding sites. MicroRNAs are the dominant small RNA class in mammalian cells. More than 1000 miRNAs have been identified and implicated in the regulation of a large fraction of mammalian genes and in many cellular processes [7]. MicroRNAs are essential also for pluripotency [8–10]. Two antagonistic miRNA families strongly contribute to control of developmental capacity in mammals. Highly conserved Let-7 miRNAs are typically found in differentiated animal tissues and are associated with negative regulation of pluripotency and tumor suppression in mammals [11]. Remarkably, Let-7 miRNAs are the most abundant maternal miRNAs although their expression is minimal in early embryos [12, 13]. In contrast, miR-290 family miRNAs are virtually absent in oocytes and become expressed during zygotic genome activation and early development [13, 14]. Notably, the miR-290 family is the most abundant and most active miRNA family in embryonic stem cells [14, 15]. Opposing roles of Let-7 and miR-290 miRNA families were demonstrated in pluripotent stem cells where Let-7 promotes differentiation [16], whereas miR-290-related miRNAs suppress senescence, promote rapid proliferation, and facilitate formation of induced pluripotent stem cells [14, 17, 18]. Interestingly, the ability of miRNAs to repress translation during oocyte-to-embryo transition (OET) is strongly reduced despite intact miRNA biogenesis [12, 19–21]. This unique observation is consistent with the impressive tolerance of mouse oocytes to loss of miRNA, which extends far into the early development [20] where the first functional requirement

<sup>1</sup>Supported by Czech Science Foundation grant GACR P305/12/G034, Czech Ministry of Education, Youth, and Sports (MEYS) project KONZAKT II LH13084 to P.S., MEYS project LM2011032 to R.S., and institutional support RVO 68378050 to I.M.G.

<sup>2</sup>Correspondence: Petr Svoboda, Institute of Molecular Genetics of the Academy of Sciences of the Czech Republic, Videnska 1083, 142 20 Prague 4, Czech Republic. E-mail: svobodap@img.cas.cz

Received: 20 February 2014.

First decision: 8 April 2014.

Accepted: 25 April 2014.

© 2014 by the Society for the Study of Reproduction, Inc.

This is an Open Access article, freely available through *Biology of Reproduction's* Authors' Choice option.

eISSN: 1529-7268 <http://www.biolreprod.org>

ISSN: 0006-3363




FIG. 1. *Lin28a* and *Lin28b* expression analysis. **A**) Microarray data from the BioGPS database (NCBI accession number GSE1133 [36, 37]) show that *Lin28a* and *Lin28b* are highly expressed in the oocyte but not in somatic tissues. Data for each tissue represent relative average ( $n = 2$ ) fluorescence intensity of gene-detecting probes on the GNF1M platform (Affymetrix). Data from microarrays were normalized using the GC content adjusted Robust Multi-Array (gcRMA) algorithm and global median scaling, where the median was set to 200. **B**) Temporal expression profile of *Lin28a* transcripts during early development. Relative abundance of *Lin28a* transcripts was determined by real-time PCR. Expression data (Ct $\Delta\Delta$  values) obtained from individual

for miRNAs was observed during implantation [22–24]. MicroRNA inactivity in mouse oocytes likely prevents maternal Let-7 miRNAs from suppressing transcripts necessary for maternal genome reprogramming and facilitates exchange of the two antagonistic miRNA families [25].

Global suppression of mature miRNA activity during OET might be aided by suppression of Let-7 biogenesis by LIN28A and LIN28B, which bind Let-7 precursors and block their processing into mature miRNAs [26–30]. LIN28A (also known as LIN28) and LIN28B are highly conserved mammalian homologs of *Lin28*, which was first identified as a regulator of developmental timing in *Caenorhabditis elegans* [31]. LIN28 proteins bind Let-7 pre-miRNAs as well as various mRNAs and have multiple roles in pluripotency, development, growth, and metabolism [32–34]. Remarkably, LIN28 was used together with OCT4, SOX2, and NANOG as a pluripotency reprogramming factor to reprogram human fibroblasts into induced pluripotent stem cells [35]. Both *Lin28a* and *Lin28b* mRNAs are highly expressed in oocytes, embryos, and embryonic stem cells [36, 37]. *Lin28a*<sup>−/−</sup> mice have reduced germ cell pool sizes at embryonic and adult stages. Consequently, *Lin28a*<sup>−/−</sup> female mice have smaller ovaries, produce smaller numbers of mature oocytes, and have reduced fertility (~30% smaller litter size) [38]. In contrast, postzygotic morpholino-based knockdown of *Lin28a* resulted in most early embryos arresting between 2- and 4-cell stages, and LIN28A has been implicated in nucleogenesis during early development [39]. Here we report results of LIN28A expression analysis during OET and functional analysis of *Lin28a* and *Lin28b*, using oocyte-specific transgenic RNA interference (RNAi) [40].

## MATERIALS AND METHODS

### Mouse Oocyte/Egg/Embryo Collection, Cell Culture, and Microinjection

Fully grown, germinal vesicle (GV)-intact oocytes, metaphase II (MII) eggs, and 1-cell and 2-cell embryos were collected as previously described [41, 42]. GV oocytes were cultured in Chatot-Ziomek-Bavister (CZB) medium containing 0.2 mM isobutylmethylxanthine (IBMX; Sigma) to inhibit GV breakdown. MII eggs were cultured in CZB medium [43] and fertilized eggs in simplex optimization method derived medium with increased K<sup>+</sup> concentration (KSOM) [44]. Two-cell embryos were isolated by flushing oviducts of superovulated and mated mice 42–44 h after human chorionic gonadotropin (hCG) injection. All animal experiments were approved by the Institutional Animal Use and Care Committee and were performed consistent with legal guidelines.

### DNA Constructs

The following cloning strategy was used to generate *Lin28a/Lin28b* (*Lin28a/b*) inverted repeat for transgenic maternal *Lin28a* and *Lin28b* knockdown. Two fragments of different length were amplified from *Lin28a* transcript by RT-PCR, using forward primers GCGGATCCAGGGAGGAA GAGGAAGAGAT (long fragment) and GCGGATCCAGGAGTTTAAAG

GAAAGAGGCA (short fragment) with a reverse primer GCAAGCTTGGTTTGACACTTGTTCGCT. *Lin28b* fragment was cloned with the forward primer GCAAGCTTCCTTTGATTTCAGAAACGGAA and the reverse primer GCTCTAGAGTGCCCTGACTCTTATGTGA. PCR products were digested with *Hind*III and ligated to form *Lin28a/b* hybrid arms. These were further digested with *Bam*HI and ligated together, and the resulting inverted repeat was cloned into the pZP3EGFP vector [45] by using a unique *Xba*I site to produce pZP3EGFP-*Lin28IR* for transgenic RNAi. Positions of *Lin28a* and *Lin28b* sequences used for transgenic RNAi are shown in Supplemental Figure S1 (all supplemental data are available online at [www.biolreprod.org](http://www.biolreprod.org)).

To generate the *Lin28a Renilla* luciferase reporters, the full-length sequence and the last 500 bp of *Lin28a* 3'UTR were amplified using the forward primer GCTCTAGAGGCCAGGAGTCAGGGTTATTC (full-length) and GCTCTAGATCAAAACCAATGTAATCTGTAC (500 bp) with a common reverse primer, AGCGGCCGAGTACCAACTCTGGAGTACCA. PCR products were digested with *Xba*I and *Not*I and cloned into *Xba*I and *Not*I opened phRL-SV40 plasmid (Promega). The control firefly luciferase reporter (pFL-SV40) was created by exchanging the *Renilla* luciferase-encoding sequence in phRL-SV40 with the firefly luciferase-encoding sequence.

### In Vitro Transcription

The phRL-SV40-*Lin28a* 3'UTR variants and the control pFL-SV40 were linearized by *Not*I digestion. Capped RNAs were synthesized by in vitro transcription using a T7 mMessage mMachine kit (Ambion) according to the manufacturer's instruction. Following in vitro transcription, template DNA was removed by Turbo DNase (Ambion), and the control firefly luciferase RNA was polyadenylated using poly(A) tailing kit (Ambion). RNAs were purified with RNeasy mini-kit columns (Qiagen) and eluted in RNase-free water.

### Immunocytochemistry and Immunoblotting

Oocyte or embryo samples were fixed in 4% paraformaldehyde for 1 h at room temperature as described previously [46]. The cells were then permeabilized for 15 min in PBS containing 0.1% Triton X-100, blocked in PBS containing 0.2% immunoglobulin G (IgG)-free BSA and 0.01% Tween-20 for 30 min (blocking solution), and then incubated with rabbit anti-LIN28 antibody (clone D1A1A; diluted 1:400 in blocking solution; Cell Signaling Technology) at 4°C overnight. After 4 10-min washes in blocking solution, samples were incubated for 1 h with Alexa 594-conjugated donkey-anti-rabbit secondary antibody (diluted 1:500 in blocking solution; Invitrogen). After an additional 4 10-min washes in blocking solution, the samples were mounted in mounting medium (Vectashield) containing DAPI (4',6-diamidino-2-phenylindole; Vector Laboratories). Images were captured with a TCS SP5 model laser-scanning confocal microscope (Leica) as described previously [47]. Antibody specificity for LIN28A was confirmed by transgenic RNAi (see Fig. 3).

### RNA Isolation, Reverse Transcription, and Real-Time PCR

Total RNA was isolated from 50 oocytes/zygotes/2-cell embryos (48 h post-hCG injection) and from 30 8-cell embryos/blastocysts (72 and 96 h post-hCG, respectively) using RNeasy spin columns (Qiagen) and RNeasy spin columns (Qiagen) and reverse-transcribed with Superscript III reverse transcriptase (Invitrogen) using random hexamers as primers. The cDNA equivalent of 0.4 oocyte/zygote/2-cell embryo and 0.15 8-cell embryo/blastocyst per reaction was subjected to quantitative real-time PCR (qPCR) using Maxima SYBR Green qPCR Master Mix (Thermo Scientific) with a Mx3000P system (Stratagene). PCR primers are listed in Supplemental Table S1. Obtained real-

GV-intact oocytes (GV, n = 11) and from 1-cell (1C, n = 6), 2-cell (2C, n = 9), and 8-cell (8C, n = 4) embryos were averaged, and *Lin28a* expression was calculated relative to that of the GV stage. Error bars are SEM. C) Immunofluorescence analysis shows accumulation of LIN28A during meiotic maturation and after fertilization. GV = fully grown oocytes; MII = metaphase II egg; 1-cell = 1-cell embryo; 2-cell = 2-cell embryo. In MII, the confocal plane was placed to visualize chromosomes. LIN28A signal is shown in red. DNA staining with DAPI is shown in blue. Bar = 10 μm. Inset in the 2-cell stage panel shows the absence of LIN28A signal in a 2-cell stage stained with the secondary antibody only. D) Schematic representation of *Lin28a* 3'UTR and *Renilla* luciferase (RL) reporter constructs. At the top is shown the 5' end of the *Lin28a* coding sequence (CDS) and 3'UTR fragments (full-length 3'UTR [2997 nucleotides, represented by the dark gray line] and 3' end [498-nucleotide fragment, represented by the black line]) used to make RL reporter constructs (shown below). E) Schema of the experimental design: a tested, capped, and deadenylated sample of *Renilla* reporter RNA was co-injected with capped and adenylated firefly luciferase (FL) into GV oocytes or 1-cell embryos. Luciferase activities were measured at the stages shown after 18–20 h in culture. F) Activity of luciferase reporters in GV oocytes, MII eggs, and 2-cell embryos. *Luc* reporter RNAs were injected into GV oocytes or 1-cell embryos. Luciferase activity was analyzed in MII eggs or embryos at 18 and 20 h after microinjection, respectively. Injected GV oocytes cultured in medium containing IBMX for 18 h served as controls. The experiment was performed three times, and data are means ± SEM.




FIG. 2. Down-regulation of *Lin28a* and *Lin28b* transcripts by transgenic RNAi. **A**) Design of the transgenic RNAi vector pZP3EGFP-Lin28IR. *Lin28a* and *Lin28b* sequences used to produce the inverted repeat originated from 3'UTR regions (Supplemental Fig. S1). Detailed description of the vector construction is described in *Materials and Methods*. **B**) Schematic depiction of simultaneous targeting of *Lin28a* and *Lin28b* by transgenic RNAi. **C**) Transgenic mice carrying the pZP3EGFP-Lin28IR transgene show uniform enhanced green fluorescent green protein (EGFP) expression in oocytes. A population of fully grown GV oocytes isolated from ovaries of a transgenic animal is shown, as seen with inverted fluorescence microscopy. Bar = 100  $\mu$ m. **D**) Oocytes from mice carrying the pZP3EGFP-Lin28IR transgene show strong down-regulation of *Lin28a* and *Lin28b* expression. RT-PCR analysis was carried out as described in *Materials and Methods*. PCR-amplified amplicons after 45 cycles are shown.

time PCR data were analyzed by the Ct $\Delta\Delta$  approach, using an in-house-designed software.

#### Fluidigm Analysis of *Lin28a* Expression

Single oocytes or early embryos were collected in 0.5-ml tubes containing 4  $\mu$ l of lysis solution (3.7  $\mu$ l of water and 0.3  $\mu$ l of RNase inhibitor [Ribolock; Thermo Scientific]). Samples were immediately frozen at  $-80^{\circ}\text{C}$ . Cell lysis was performed at  $85^{\circ}\text{C}$  for 10 min. Lysates were placed on ice for 2 min, and RNA was subjected to reverse transcription reaction. The reaction mixture consisted of 2  $\mu$ l of dNTPs (2.5 mM each), 2  $\mu$ l of 5 $\times$  reverse transcriptase (RT) buffer, 0.5  $\mu$ l of random hexamer primer (all Thermo Scientific), 1  $\mu$ l of 1 pg of rabbit globin as an external spike control (Sigma-Aldrich) and 0.5  $\mu$ l of RevertAid polymerase (Thermo Scientific). Complementary DNA was synthesized at  $42^{\circ}\text{C}$  for 50 min, followed by inactivation of the enzyme at  $70^{\circ}\text{C}$  for 10 min. Converted RNA was subjected to preamplification PCR reaction to increase the number of template molecules. Preamplification was carried out in a total

volume of 40  $\mu$ l, which consisted of 2  $\mu$ l of cDNA, 2  $\mu$ l of primer mixture (0.5  $\mu$ M each primer), 30  $\mu$ l of 2 $\times$  SYBR Green (Thermo Scientific), and 6  $\mu$ l of water. The original cDNA template was amplified by 18 cycles under the following conditions:  $95^{\circ}\text{C}$  for 10 min; denaturation of DNA strands was performed at  $95^{\circ}\text{C}$  for 15 sec, followed by annealing at  $57^{\circ}\text{C}$  for 4 min, and strand synthesis phase at  $72^{\circ}\text{C}$  for 20 sec. Two microliters of 10 $\times$ -diluted preamplification mixture was used for Fluidigm 48  $\times$  48 well dynamic array (Fluidigm), and the experiment was carried out according to the manufacturer's protocol.

#### Analysis of miRNA Expression

Twenty oocytes or embryos were collected in 5  $\mu$ l of water supplemented with RNase inhibitor (Ribolock; Thermo Scientific) and lysed by heating at  $95^{\circ}\text{C}$  for 5 min. Lysates were directly subjected to reverse transcription with miRCURY LNA universal cDNA synthesis kit (Exiqon). A cDNA equivalent of 0.25 oocyte or embryo was used for qPCR reaction with miRCURY LNA






FIG. 3. Transgenic RNAi prevents accumulation of LIN28A during oocyte-to-zygote transition. GV oocytes (GV), 1-cell embryos (1-cell), and 2-cell embryos (2-cell) from wild-type and transgenic females were stained with  $\alpha$ -LIN28A antibody (red signal). DNA was stained with DAPI (blue signal). Bar = 10  $\mu$ m.

SYBR Green Master Mix and for LNA-modified primers to amplify U6 small nuclear RNA (U6-snrRNA), Let-7a miRNA, and miR-30c miRNA (Exiqon). Real-time PCR was performed with an Mx3000P system (Stratagene) and analyzed by using the Ct $\Delta\Delta$  approach, using in-house-designed software.

### Luciferase Reporter Assay

Fully grown GV oocytes or zygotes (22 h post-hCG) were microinjected with ~5  $\mu$ l of a mixture of control polyadenylated firefly luciferase RNA (15 ng/ $\mu$ l) and RNA encoding *Renilla* luciferase fused to the last 500 bp of *Lin28a* 3'UTR (16 ng/ $\mu$ l) or to full-length *Lin28a* 3'UTR (39 ng/ $\mu$ l). Injected oocytes were cultured for 18 h in CZB medium with or without 0.2 mM IBMX to prevent or allow maturation to MII stage, respectively. Injected zygotes were cultured for 20 h in KSOM medium. Individual oocytes/eggs/embryos were lysed in 11  $\mu$ l of 1 $\times$  passive lysis buffer, and half of the lysate was assayed by using a dual-luciferase reporter assay system (Promega) with a Modulus microplate reader luminometer (Turner Biosystems). For signal normalization, the *Renilla* luciferase activity readout from noninjected oocytes/eggs was subtracted as background, and *Renilla* luciferase activity was then normalized to that of the co-injected control firefly luciferase.

## RESULTS

### *Lin28a* Encodes Dormant Maternal mRNA

We became interested in *Lin28a* because we found it among the RNA binding factors whose expression was highly enriched in mouse oocytes (Fig. 1A). Therefore, we investigated *Lin28a* expression during OET in more detail (Fig. 1, B and C). The temporal pattern of mRNA expression indicated that the maternal pool of *Lin28a* transcript is strongly depleted

by the time of the zygotic genome activation at the 2-cell stage. Immunofluorescent staining of oocytes and zygotes with LIN28A-specific antibody revealed mostly cytoplasmic signal, which was low in the fully grown GV oocyte whereas the signal strongly increased during meiotic maturation and zygotic genome activation (Fig. 1C and Supplemental Fig. S2).

The apparent anticorrelation between transcript and protein levels suggested that *Lin28a* encodes a dormant maternal mRNA. To examine regulation of *Lin28a* mRNA translation, we created reporters where *Renilla* luciferase coding sequence was fused with the entire *Lin28a* 3'UTR (RL-full) or the last 498 nucleotides of *Lin28a* 3'UTR (RL-500) (Fig. 1D). Capped, but non-adenylated, recombinant *Renilla* RNAs were co-injected with a capped and polyadenylated firefly luciferase, which served as an internal control for normalization. Microinjected GV oocytes were either matured for 18 h to MII or maintained for the same time in the presence of IBMX to prevent resumption of meiosis (Fig. 1E). To examine the control of *Lin28a* translation through its 3'UTR after fertilization, we microinjected reporters into 1-cell embryos, which were cultured until the 2-cell stage for 20 h (Fig. 1E). Our results showed that both of the *Lin28a* 3'UTR fragments supported increased translation of deadenylated transcripts during meiotic maturation compared to GV oocytes (Fig. 1F). Both of the *Lin28a* 3'UTR fragments showed up-regulation of luciferase activity, comparable to our previous experiments with *Ccnbl* and *Dcp2* 3'UTRs [48]. Interestingly, RL-500 showed the same accumulation of luciferase activity during meiotic maturation and in the zygote, whereas RL-full luciferase activity was approximately three times higher in early embryos, indicating distinct post-transcriptional control of *Lin28a* in early embryos enabling further accumulation of the protein product. Taken together, *Lin28a* encodes an abundant dormant maternal transcript translated during meiotic maturation and after fertilization, thus resulting in a robust accumulation of LIN28A at the time of the zygotic genome activation.

### Efficient Depletion of *Lin28a* and *Lin28b* by Transgenic RNAi

To address the role of accumulation of LIN28A during oocyte-to-zygote transition, we decided to deplete the maternal pool of *Lin28a* by transgenic RNAi, which allows studying the gene function in vivo during oocyte growth and OET. Because mouse oocytes also express *Lin28b*, a close homolog of *Lin28a*, which might potentially compensate for the loss of *Lin28a*, we decided to knock down both homologs simultaneously. In order to do that, we fused together 3'UTR fragments of *Lin28a* and *Lin28b* and created an inverted repeat, which was inserted into the transgenic RNAi vector (Fig. 2A). Upon transcription, which occurs during oocyte growth [49], this inverted repeat should be processed by RNase III Dicer into small interfering RNAs (siRNAs), which would target *Lin28a* and *Lin28b* mRNAs (Fig. 2B).

Upon pronuclear injection of the transgenic construct, we obtained one transgenic line where the transgene was expressed (Fig. 2C) and mediated highly efficient knockdown of both *Lin28* homologs (Fig. 2D). Immunostaining of transgenic and wild-type oocytes and early embryos also showed lack of LIN28A signal in 1-cell and 2-cell embryos, demonstrating that transgenic RNAi efficiently prevented accumulation of LIN28A during oocyte-to-zygote transition (Fig. 3).




FIG. 4. Transgenic RNAi had no effect on fertility but relieved Let-7a expression during zygotic genome activation. **A**) Litter sizes of transgenic maternal *Lin28a/b* knockdown C57Bl/6 females were compared with their wild-type siblings. In total, 5–7 litters were counted from 4 knockdown and 3 wild-type females. **B**) Frequencies of productive matings in maternal *Lin28a/b* knockdown were compared with those of wild-type C57Bl/6 females over a 25-wk period. **C**) Litter sizes of maternal *Lin28a/b* knockdown females compared with those of wild-type C57Bl/6 females, classified by different ages as young (14–22 wk old), middle aged (23–30 wk old), and old (31–40 wk old). **D**) RT-qPCR analyses of Let-7a and miR-30c miRNAs in maternal *Lin28a/b* knockdown (KD) oocytes and wild-type (WT) oocytes and preimplantation embryos. MicroRNA expression values were normalized to those of U6-sRNA control. Error bars represent SEM. \*Statistically significant differences (two-tailed *t*-test,  $P = 0.024$ ) between wild-type (WT) and KD samples, which were observed only in 2-cell embryos. Data were pooled from four independent experiments performed in duplicate. Comparison with published studies of Let-7a expression in oocytes and early embryos [13, 68, 69] suggests that *Lin28a/b* knockdown results in Let-7a levels in 2-cell embryos exceeded those in wild-type oocytes.

#### Effects of *Lin28a* and *Lin28b* Depletion in Preimplantation Embryos

First, we analyzed whether *Lin28a/b* knockdown had any effect on fertility. Analysis of litter sizes did not show any statistically significant difference between wild-type and




FIG. 5. Analysis of gene expression in early embryos from transgenic RNAi females. Gene expression in early embryos from transgenic females was analyzed by qPCR at 48, 72, and 96 h after hCG. Color coding indicates different categories of selected genes. Analyzed genes were (from left to right): *Lin28a* and *Lin28b*, *pri-miR-290* and *pri-Let-7a* (primary miRNA transcripts), *Oct4*, *Nanog*, *Klf4*, and *Sox2* (core pluripotency factors), *Trim71* and *Igf2bp1* (known targets of Let-7), *Rbl2* and *p21* (known targets of miR-290 miRNAs), and *Ccnb1* and *H2A* (LIN28-bound mRNAs). Data are expressed as the relative ratio (log<sub>2</sub>-fold change) of expression in embryos from transgenic (KD) to that in wild-type females (WT). Asterisks indicate statistical significance of the change (\* $P < 0.05$ , \*\* $P < 0.01$ , \*\*\* $P < 0.001$ ;  $n = 3$ ). Error bars = SEM.

transgenic mice (Fig. 4A). Likewise, mating efficiency of wild-type and transgenic mice was equal (Fig. 4B). We also tested whether there would be any age-dependent effect on fertility. Again, we did not observe any statistically significant differences in litter sizes between younger and older mice (Fig. 4C).

Next, we analyzed whether *Lin28a/b* knockdown had any effect on Let-7a miRNA levels. Real-time PCR analysis of Let-7a revealed a strong, statistically significant increase in Let-7a in 2-cell embryos isolated from transgenic females but not in GV oocytes (Fig. 4D). This result correlates with LIN28A levels and indicates that LIN28A (and presumably LIN28B) acts during the zygotic genome activation to block Let-7a biogenesis from newly transcribed precursors.

In addition to the suppression of Let-7 through binding Let-7 pre-miRNA, LIN28 proteins were shown to bind mRNAs [50–52]. In order to test whether depletion of maternally provided LIN28 proteins had any observable effects on gene expression, we analyzed gene expression of selected candidate genes during early development. We analyzed levels of miR-

290 and Let-7 primary miRNAs (pri-miRNAs), core pluripotency factors (*Oct4*, *Sox2*, *Nanog*, and *Klf4*), known targets of Let-7 (*Trim71* and *Igf2bp1*) and miR-290 miRNAs (*Rbl2* and *p21*), and two known LIN28-bound mRNAs (*Ccnb1* and *H2A*). We also monitored *Lin28a* and *Lin28b* levels (Fig. 5). Our results showed that ~4-fold reduction of *Lin28a/b* transcripts persisted in embryos from transgenic mice up to the 8-cell stage (72 h post-hCG). However, 96 h after hCG injection, there were essentially no differences between embryos from wild-type and those from transgenic mice. Generally, none of the examined genes but *Lin28a* and *Lin28b* showed a strong change in gene expression. Four additional statistically significant minor expression changes did not exhibit any clear pattern, and it is not clear whether they represented direct consequences of *Lin28a/b* knockdown and whether they are functionally significant at all (Fig. 5). At 48 h after hCG injection, we observed a mild reduction of pri-Let-7a ( $P < 0.05$ ), but it is unclear how LIN28 proteins could affect primary miRNA (pri-miRNA) transcript as they target the Let-7 at the next biogenesis stage, the precursor miRNA (pre-miRNA), which is released from pri-miRNA. Similarly, a mild upregulation of pri-miR-290 ( $P < 0.05$ ) at 72 h post-hCG is of unclear origin and significance.

## DISCUSSION

We report that *Lin28a* encodes a dormant maternal mRNA that is translated during oocyte-to-zygote transition in order to prevent zygotic expression of Let-7 during the zygotic genome activation. *Lin28a* mRNA expression analysis is in agreement with previous microarray analyses [53, 54] and RT-PCR profiling [39]. Altogether, these data demonstrate that *Lin28a* transcript is abundant in the oocyte and becomes degraded after fertilization, and zygotic transcripts start to accumulate after the 8-cell stage. The dormancy of LIN28 proteins is also consistent with published data, which show *Lin28a* recruitment to polysomes and increased protein abundance by immunofluorescent staining of LIN28A [2, 39]. However, there is an inconsistency concerning the localization of LIN28A. Previously, a strong nucleolar localization of LIN28A has been reported [39], whereas our immunofluorescent staining yielded a dominant cytoplasmic signal, which is consistent with other data obtained from mammalian cells, using various antibodies recognizing LIN28A [55–57]. Although we did not observe a distinct nucleolar enrichment (Supplemental Fig. S2), it is still possible that the antibodies used after the method described by Vogt et al. [39] and in our study recognize the same protein but the staining pattern differs due to variable epitope exposure. This would explain why both studies observed experimentally reduced levels of LIN28A despite a differing staining pattern (Fig. 3).

The functional study of *Lin28a* and *Lin28b* used a transgenic RNAi approach that offers several advantages over microinjection of siRNA, double-stranded RNA (dsRNA), or morpholino [58]. Transgenic RNAi allows for highly specific mRNA knockdown during oocyte growth without any off-targeting effects [59]. While insertion site effects could be a potential source of artifacts, it is extremely unlikely that the lack of developmental phenotype and post-zygotic up-regulation of Let-7 would be manifestations of insertion site effects (which would have to compensate Let-7 overexpression phenotype).

Unlike the *Lin28a* knock-out [38], transgenic RNAi knockdown of *Lin28a* and *Lin28b* did not result in reduced fertility. One explanation could be that the knockdown approach does not reduce *Lin28a* expression below the

threshold level necessary for the phenotype manifestation. However, there are several arguments that this is not the case. First, maternal *Lin28a* was strongly suppressed at both mRNA and protein levels. In addition, we also down-regulated *Lin28b*, which could be partially compensating down-regulation of *Lin28a* (notably, *Lin28b* expression was not sufficient to prevent the phenotype in knock-out females [38]). Second, we detect relieved repression of Let-7 expression during zygotic genome activation demonstrating that *Lin28a* and *Lin28b* knockdown efficiently suppressed their function. Third, reduced fertility in knock-out animals might originate from events occurring at the beginning of the oocyte growth or before. *Lin28a* knock-out females have smaller ovaries and reduced numbers of primordial follicles; it was suggested that the loss of *Lin28a* affects proliferation of primordial germ cells. In contrast, transgenic RNAi knockdown is initiated at the onset of oocyte growth [49].

A postzygotic knockdown of maternal *Lin28a* using a morpholino approach implied an essential function in preimplantation development because it caused 2- and 4-cell developmental arrest [39]. However, this result is in strong contrast with the knock-out phenotype, where maternal contribution of *Lin28a* was completely eliminated, yet it resulted in mildly reduced fertility (average litter size was reduced from ~10 to ~7 pups [38]), which might seemingly have originated in defects in oocyte development rather than in postzygotic events [38]. Similarly, our results do not provide any evidence for postzygotic developmental arrest. As mentioned above, the lack of strong postzygotic phenotype in the transgenic RNAi model could be caused by insufficient knockdown of *Lin28a* and/or *Lin28b*, where their transcript levels were not reduced below a threshold necessary for phenotype appearance. However, as discussed above, RT-PCR and immunofluorescent staining demonstrated strong knockdown that resulted in up-regulation of mature Let-7. Another explanation of the reported morpholino-induced 2- and 4-cell developmental arrest [39] is that it does not represent a *Lin28a*-specific phenotype. Although morpholinos are supposed to be highly specific, a morpholino may also inhibit a gene(s) other than the intended target [60]. Off-targeting that results in 2- and 4-cell developmental arrest was reported in morpholino experiments in early mouse embryos. For example, knockdown of *Oct4* using morpholino microinjection into zygotes caused 4-cell embryo arrest in ~60% of embryos (<10% in control) [61], whereas maternal zygotic knock-out embryos genetically lacking *Oct4* developed to the blastocyst stage [62]. Similarly, *Nanog* morpholino knockdown caused 4-cell arrest in ~50% embryos [61]; embryonic knock-out develops to the blastocyst stage [63] whereas *Nanog* does not seem to be maternally expressed [37, 64]. Because experimental groups were relatively small and the experiment did not include a rescue control [39], it should be considered that the observed 4-cell developmental arrest might represent an off-target effect.

Dormancy of *Lin28a* and the absence of phenotype in transgenic females, despite increased expression of Let-7 during zygotic genome activation, offers important insights into the role of LIN28 homologs in the mouse model (Fig. 6) and raises a number of interesting questions. First, LIN28 proteins have multiple roles in pluripotency, development, growth, and metabolism and, in addition to their role in Let-7 biogenesis, they bind various mRNAs [50–52]. However, none of these mRNA regulations appears to be functionally important during OET, where we observed increased cytoplasmic abundance of LIN28A.

Second, there is a strongly reduced ability of miRNAs to repress translation during OET despite generally intact miRNA






FIG. 6. Model of *Lin28* function during OET. The model integrates published data and results presented in this work. The maternal pool of *Lin28a* transcripts was translated during meiotic maturation and after fertilization (our data and those in ref. [39]). LIN28 does not target the mature Let-7 micro-RNA, but the precursor miRNA (pre-miRNA) released from the primary miRNA (pri-miRNA) transcript [26–30] suppresses zygotic expression of Let-7 (our data). The primary miRNA transcript (pri-miRNA) carrying miR-290-295 becomes expressed during the zygotic genome activation at the 2-cell stage, and miR-290-295 accumulate during early development [13, 54]. Superimposed on specific suppression of Let-7 is the global suppression of miRNA activity, which reduces the ability of mature miRNAs loaded on effector complexes to target imperfectly complementary target mRNAs [19, 20]. Likewise, oocytes and zygotes completely lacking canonical miRNAs (via maternal and maternal zygotic knock-out of *Dgcr8*, an essential miRNA biogenesis factor) develop to the blastocyst stage [20], further demonstrating that miRNAs are not important regulators of OET. Dynamics of P-bodies suggest that suppression of miRNA activity begins with the onset of oocyte growth and that miRNAs regain their function at approximately the 8-cell or morula stage [47].

biogenesis [12, 19–21]. This phenomenon should mask any phenotype caused by up-regulation of zygotic expression of Let-7. Thus, why would mouse OET use two functionally redundant but mechanistically different mechanisms to suppress Let-7 function? We propose two possible explanations for this phenomenon: first, *Lin28* homologs may function as a fail-safe mechanism, which would protect the developing embryo from Let-7 activity in case global suppression of miRNA activity was relieved. We do not know the mechanism responsible for reduced ability of miRNAs to suppress translation in the oocyte and during OET. It seems that the inhibition is provided maternally by ineffective formation of the miRNA-guided effector complexes, and it is unclear how embryos acquire ability to produce fully active miRNA-guided effector complexes. Thus, maternally provided *Lin28* homologs may assure Let-7 repression in the case of premature reactivation of the miRNA pathway. Second, Let-7 suppression by LIN28 during embryonic development is an evolutionarily conserved mechanism. Thus, *Lin28a* dormancy and activity during zygotic genome activation may be a vestigial mechanism that became replaced by global miRNA suppression. Whether the same global miRNA repression exists in other mammalian oocytes is not known. In fact, there is some evidence that miRNAs are active in porcine oocytes and [65] bovine early embryos [66]. Furthermore, Dicer function appears to be unique in mouse oocytes [67]. Therefore, the lack of requirement for maternally provided LIN28 homologs during OET in mice might represent another unique example of small RNA regulation rather than a common mammalian case. Consequently, there is an open possibility that the role of maternally provided LIN28 homologs is significant in other mammals during their OET.

## ACKNOWLEDGMENT

We thank Inken Beck (Institute of Molecular Genetics, AS CR) for technical assistance and advice and Radek Malik (Institute of Molecular Genetics, AS CR) for help with manuscript preparation.

## REFERENCES

- Wilmot I, Schnieke AE, McWhir J, Kind AJ, Campbell KH. Viable offspring derived from fetal and adult mammalian cells. *Nature* 1997; 385: 810–813.
- Chen J, Melton C, Suh N, Oh JS, Horner K, Xie F, Sette C, Blleloch R, Conti M. Genome-wide analysis of translation reveals a critical role for deleted in azoospermia-like (*Dazl*) at the oocyte-to-zygote transition. *Genes Dev* 2011; 25:755–766.
- Pique M, Lopez JM, Foissac S, Guigo R, Mendez R. A combinatorial code for CPE-mediated translational control. *Cell* 2008; 132:434–448.
- Potiredy S, Midic U, Liang CG, Obradovic Z, Latham KE. Positive and negative cis-regulatory elements directing postfertilization maternal mRNA translational control in mouse embryos. *Am J Physiol Cell Physiol* 2010; 299:C818–827.
- Richter JD. CPEB: a life in translation. *Trends Biochem Sci* 2007; 32: 279–285.
- Villalba A, Coll O, Gebauer F. Cytoplasmic polyadenylation and translational control. *Curr Opin Genet Dev* 2011; 21:452–457.
- Bartel DP. MicroRNAs: target recognition and regulatory functions. *Cell* 2009; 136:215–233.
- Kanellopoulou C, Muljo SA, Kung AL, Ganesan S, Drapkin R, Jenuwein T, Livingston DM, Rajewsky K. Dicer-deficient mouse embryonic stem cells are defective in differentiation and centromeric silencing. *Genes Dev* 2005; 19:489–501.
- Murchison EP, Partridge JF, Tam OH, Cheloufi S, Hannon GJ. Characterization of Dicer-deficient murine embryonic stem cells. *Proc Natl Acad Sci U S A* 2005; 102:12135–12140.
- Wang Y, Medvid R, Melton C, Jaenisch R, Blleloch R. DGCR8 is essential for microRNA biogenesis and silencing of embryonic stem cell self-renewal. *Nat Genet* 2007; 39:380–385.
- Bussing I, Slack FJ, Grosshans H. Let-7 microRNAs in development, stem cells and cancer. *Trends Mol Med* 2008; 14:400–409.
- Tam OH, Aravin AA, Stein P, Girard A, Murchison EP, Cheloufi S, Hodges E, Anger M, Sachidanandam R, Schultz RM, Hannon GJ. Pseudogene-derived small interfering RNAs regulate gene expression in mouse oocytes. *Nature* 2008; 453:534–538.
- Tang F, Kaneda M, O'Carroll D, Hajkova P, Barton SC, Sun YA, Lee C, Tarakhovsky A, Lao K, Surani MA. Maternal microRNAs are essential for mouse zygotic development. *Genes Dev* 2007; 21:644–648.
- Sinkkonen L, Huguenschmidt T, Berninger P, Gaidatzis D, Mohn F, Artus-Revel CG, Zavolan M, Svoboda P, Filipowicz W. MicroRNAs control de novo DNA methylation through regulation of transcriptional repressors in mouse embryonic stem cells. *Nat Struct Mol Biol* 2008; 15:259–267.
- Marson A, Levine SS, Cole MF, Frampton GM, Brambrink T, Johnstone S, Guenther MG, Johnston WK, Wernig M, Newman J, Calabrese JM, Dennis LM, et al. Connecting microRNA genes to the core transcriptional regulatory circuitry of embryonic stem cells. *Cell* 2008; 134:521–533.
- Melton C, Judson RL, Blleloch R. Opposing microRNA families regulate self-renewal in mouse embryonic stem cells. *Nature* 2010; 463:621–626.
- Wang Y, Baskerville S, Shenoy A, Babiarz JE, Baehner L, Blleloch R. Embryonic stem cell-specific microRNAs regulate the G1-S transition and promote rapid proliferation. *Nat Genet* 2008; 40:1478–1483.
- Banito A, Rashid ST, Acosta JC, Li S, Pereira CF, Geti I, Pinho S, Silva JC, Azuara V, Walsh M, Vallier L, Gil J. Senescence impairs successful reprogramming to pluripotent stem cells. *Genes Dev* 2009; 23:2134–2139.
- Ma J, Flemr M, Stein P, Berninger P, Malik R, Zavolan M, Svoboda P, Schultz RM. MicroRNA activity is suppressed in mouse oocytes. *Curr Biol* 2010; 20:265–270.
- Suh N, Baehner L, Moltzahn F, Melton C, Shenoy A, Chen J, Blleloch R. MicroRNA function is globally suppressed in mouse oocytes and early embryos. *Curr Biol* 2010; 20:271–277.
- Watanabe T, Totoki Y, Toyoda A, Kaneda M, Kuramochi-Miyagawa S, Obata Y, Chiba H, Kohara Y, Kono T, Nakano T, Surani MA, Sakaki Y, et al. Endogenous siRNAs from naturally formed dsRNAs regulate transcripts in mouse oocytes. *Nature* 2008; 453:539–543.
- Bernstein E, Kim SY, Carmell MA, Murchison EP, Alcorn H, Li MZ, Mills AA, Elledge SJ, Anderson KV, Hannon GJ. Dicer is essential for mouse development. *Nat Genet* 2003; 35:215–217.
- Foshay KM, Gallicano GL. miR-17 family miRNAs are expressed during early mammalian development and regulate stem cell differentiation. *Dev Biol* 2009; 326:431–443.
- Pernaute B, Spruce T, Rodriguez TA, Manzanares M. MiRNA-mediated

- regulation of cell signaling and homeostasis in the early mouse embryo. *Cell Cycle* 2011; 10:584–591.
25. Svoboda P, Flemr M. The role of miRNAs and endogenous siRNAs in maternal-to-zygotic reprogramming and the establishment of pluripotency. *EMBO Rep* 2010; 11:590–597.
  26. Heo I, Joo C, Cho J, Ha M, Han J, Kim VN. Lin28 mediates the terminal uridylation of let-7 precursor MicroRNA. *Mol Cell* 2008; 32:276–284.
  27. Newman MA, Thomson JM, Hammond SM. Lin-28 interaction with the Let-7 precursor loop mediates regulated microRNA processing. *RNA* 2008; 14:1539–1549.
  28. Piskounova E, Viswanathan SR, Janas M, LaPierre RJ, Daley GQ, Sliz P, Gregory RI. Determinants of microRNA processing inhibition by the developmentally regulated RNA-binding protein Lin28. *J Biol Chem* 2008; 283:21310–21314.
  29. Rybak A, Fuchs H, Smirnova L, Brandt C, Pohl EE, Nitsch R, Wulczyn FG. A feedback loop comprising lin-28 and let-7 controls pre-let-7 maturation during neural stem-cell commitment. *Nat Cell Biol* 2008; 10: 987–993.
  30. Viswanathan SR, Daley GQ, Gregory RI. Selective blockade of micro-RNA processing by Lin28. *Science* 2008; 320:97–100.
  31. Moss EG, Lee RC, Ambros V. The cold shock domain protein LIN-28 controls developmental timing in *C. elegans* and is regulated by the lin-4 RNA. *Cell* 1997; 88:637–646.
  32. Huang Y. A mirror of two faces: Lin28 as a master regulator of both miRNA and mRNA. *Wiley Interdiscip Rev RNA* 2012; 3:483–494.
  33. Shyh-Chang N, Daley GQ. Lin28: primal regulator of growth and metabolism in stem cells. *Cell Stem Cell* 2013; 12:395–406.
  34. Viswanathan SR, Daley GQ. Lin28: A microRNA regulator with a macro role. *Cell* 2010; 140:445–449.
  35. Yu J, Vodyanik MA, Smuga-Otto K, Antosiewicz-Bourget J, Frane JL, Tian S, Nie J, Jonsdottir GA, Ruotti V, Stewart R, Slukvin II, Thomson JA. Induced pluripotent stem cell lines derived from human somatic cells. *Science* 2007; 318:1917–1920.
  36. Su AI, Cooke MP, Ching KA, Hakak Y, Walker JR, Wiltshire T, Orth AP, Vega RG, Sapinoso LM, Moqrich A, Patapoutian A, Hampton GM, et al. Large-scale analysis of the human and mouse transcriptomes. *Proc Natl Acad Sci U S A* 2002; 99:4465–4470.
  37. Su AI, Wiltshire T, Batalov S, Lapp H, Ching KA, Block D, Zhang J, Soden R, Hayakawa M, Kreiman G, Cooke MP, Walker JR, et al. A gene atlas of the mouse and human protein-encoding transcriptomes. *Proc Natl Acad Sci U S A* 2004; 101:6062–6067.
  38. Shinoda G, De Soysa TY, Seligson MT, Yabuuchi A, Fujiwara Y, Huang PY, Hagan JP, Gregory RI, Moss EG, Daley GQ. Lin28a regulates germ cell pool size and fertility. *Stem Cells* 2013; 31:1001–1009.
  39. Vogt EJ, Meglicki M, Hartung KI, Borsuk E, Behr R. Importance of the pluripotency factor LIN28 in the mammalian nucleolus during early embryonic development. *Development* 2012; 139:4514–4523.
  40. Stein P, Svoboda P, Schultz RM. Transgenic RNAi in mouse oocytes: a simple and fast approach to study gene function. *Dev Biol* 2003; 256: 187–193.
  41. Stein P, Svoboda P. Collection of early mouse embryos for RNAi. *CSH Protoc* 2006; 2006.
  42. Stein P, Svoboda P. Collection of mouse oocytes for RNAi. *CSH Protoc* 2006; 2006.
  43. Chatot CL, Ziomek CA, Bavister BD, Lewis JL, Torres I. An improved culture medium supports development of random-bred 1-cell mouse embryos in vitro. *J Reprod Fertil* 1989; 86:679–688.
  44. Erbach GT, Lawitts JA, Papaioannou VE, Biggers JD. Differential growth of the mouse preimplantation embryo in chemically defined media. *Biol Reprod* 1994; 50:1027–1033.
  45. Svoboda P. Long dsRNA and silent genes strike back: RNAi in mouse oocytes and early embryos. *Cytogenet Genome Res* 2004; 105:422–434.
  46. Flemr M, Svoboda P. Ribonucleoprotein localization in mouse oocytes. *Methods* 2011; 53:136–141.
  47. Flemr M, Ma J, Schultz RM, Svoboda P. P-body loss is concomitant with formation of a messenger RNA storage domain in mouse oocytes. *Biol Reprod* 2010; 82:1008–1017.
  48. Ma J, Flemr M, Strnad H, Svoboda P, Schultz RM. Maternally recruited DCP1A and DCP2 contribute to messenger RNA degradation during oocyte maturation and genome activation in mouse. *Biol Reprod* 2013; 88: 11.
  49. Philpott CC, Ringuette MJ, Dean J. Oocyte-specific expression and developmental regulation of ZP3, the sperm receptor of the mouse zona pellucida. *Dev Biol* 1987; 121:568–575.
  50. Hafner M, Max KE, Bandaru P, Morozov P, Gerstberger S, Brown M, Molina H, Tuschl T. Identification of mRNAs bound and regulated by human LIN28 proteins and molecular requirements for RNA recognition. *RNA* 2013; 19:613–626.
  51. Wilbert ML, Huelga SC, Kapeli K, Stark TJ, Liang TY, Chen SX, Yan BY, Nathanson JL, Hutt KR, Lovci MT, Kazan H, Vu AQ, et al. LIN28 binds messenger RNAs at GGAGA motifs and regulates splicing factor abundance. *Mol Cell* 2012; 48:195–206.
  52. Graf R, Munschauer M, Mastrobuoni G, Mayr F, Heinemann U, Kempa S, Rajewsky N, Landthaler M. Identification of LIN28B-bound mRNAs reveals features of target recognition and regulation. *RNA Biol* 2013; 10: 1146–1159.
  53. Zeng F, Baldwin DA, Schultz RM. Transcript profiling during preimplantation mouse development. *Dev Biol* 2004; 272:483–496.
  54. Zeng F, Schultz RM. RNA transcript profiling during zygotic gene activation in the preimplantation mouse embryo. *Dev Biol* 2005; 283: 40–57.
  55. Balzer E, Moss EG. Localization of the developmental timing regulator Lin28 to mRNP complexes, P-bodies and stress granules. *RNA Biol* 2007; 4:16–25.
  56. Moss EG, Tang L. Conservation of the heterochronic regulator Lin-28, its developmental expression and microRNA complementary sites. *Dev Biol* 2003; 258:432–442.
  57. Zheng K, Wu X, Kaestner KH, Wang PJ. The pluripotency factor LIN28 marks undifferentiated spermatogonia in mouse. *BMC Dev Biol* 2009; 9: 38.
  58. Malik R, Svoboda P. Transgenic RNAi in mouse oocytes: the first decade. *Anim Reprod Sci* 2012; 134:64–68.
  59. Stein P, Zeng F, Pan H, Schultz RM. Absence of non-specific effects of RNA interference triggered by long double-stranded RNA in mouse oocytes. *Dev Biol* 2005; 286:464–471.
  60. Eisen JS, Smith JC. Controlling morpholino experiments: don't stop making antisense. *Development* 2008; 135:1735–1743.
  61. Tan MH, Au KF, Leong DE, Foygel K, Wong WH, Yao MW. An Oct4-Sall4-Nanog network controls developmental progression in the pre-implantation mouse embryo. *Mol Syst Biol* 2013; 9:632.
  62. Wu G, Han D, Gong Y, Sebastiano V, Gentile L, Singhal N, Adachi K, Fischedick G, Ortmeier C, Sinn M, Radstaak M, Tomilin A, et al. Establishment of totipotency does not depend on Oct4A. *Nat Cell Biol* 2013; 15:1089–1097.
  63. Mitsui K, Tokuzawa Y, Itoh H, Segawa K, Murakami M, Takahashi K, Maruyama M, Maeda M, Yamanaka S. The homeoprotein Nanog is required for maintenance of pluripotency in mouse epiblast and ES cells. *Cell* 2003; 113:631–642.
  64. Smallwood SA, Tomizawa S, Krueger F, Ruf N, Carli N, Segonds-Pichon A, Sato S, Hata K, Andrews SR, Kelsey G. Dynamic CpG island methylation landscape in oocytes and preimplantation embryos. *Nat Genet* 2011; 43:811–814.
  65. Chen L, Hu X, Dai Y, Li Q, Wang X, Li Q, Xue K, Li Y, Liang J, Wang Y, Liu X, Li N. MicroRNA-27a activity is not suppressed in porcine oocytes. *Front Biosci (Elite Ed)* 2012; 4:2679–2685.
  66. Tripurani SK, Wee G, Lee KB, Smith GW, Wang L, Jianboyao. MicroRNA-212 post-transcriptionally regulates oocyte-specific basic-helix-loop-helix transcription factor, factor in the germline alpha (FIGLA), during bovine early embryogenesis. *PLoS One* 2013; 8:e76114.
  67. Flemr M, Malik R, Franke V, Nejepinska J, Sedlacek R, Vlahovick K, Svoboda P. A retrotransposon-driven dicer isoform directs endogenous small interfering RNA production in mouse oocytes. *Cell* 2013; 155: 807–816.
  68. Garcia-Lopez J, del Mazo J. Expression dynamics of microRNA biogenesis during preimplantation mouse development. *Biochim Biophys Acta* 2012; 1819:847–854.
  69. Ohnishi Y, Totoki Y, Toyoda A, Watanabe T, Yamamoto Y, Tokunaga K, Sakaki Y, Sasaki H, Hohjoh H. Small RNA class transition from siRNA/piRNA to miRNA during pre-implantation mouse development. *Nucleic Acids Res* 2010; 38:5141–5151.