



## **New Zealand Tenuipalpidae (Acari: Trombidiformes): A new species of Acaricis from Cyperaceae and its ontogenetic patterns in chaetotaxy**

Authors: Xu, Yun, and Zhang, Zhi-Qiang

Source: Systematic and Applied Acarology, 18(4) : 357-388

Published By: Systematic and Applied Acarology Society

URL: <https://doi.org/10.11158/saa.18.4.6>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non-commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

## New Zealand Tenuipalpidae (Acari: Trombidiformes): A new species of *Acaricis* from Cyperaceae and its ontogenetic patterns in chaetotaxy

YUN XU<sup>1</sup> & ZHI-QIANG ZHANG<sup>2,3,4</sup>

<sup>1</sup> Key Laboratory of Integrated Pest Management for Fujian-Taiwan Crops, Ministry of Agriculture; Fujian Provincial Key Laboratory of Insect Ecology; Fujian Agricultural and Forestry University, Fuzhou 350002, China

<sup>2</sup> Landcare Research, Private Bag 92170, Auckland, New Zealand

<sup>3</sup> Center for Biodiversity and Biosecurity, School of Biological Sciences, University of Auckland, Tamaki Campus, Auckland, New Zealand

<sup>4</sup> Corresponding author: Zhi-Qiang Zhang; [ZhangZ@landcareresearch.co.nz](mailto:ZhangZ@landcareresearch.co.nz)

### Abstract

The genus *Acaricis* (Acari: Tenuipalpidae) was represented by two species from Australian sedges prior to this study. A new species, *Acaricis urigersoni* **sp. nov.**, is here described and illustrated from leaves of *Gahnia lacera* (Cyperaceae) in Auckland, New Zealand. In this paper, we presented the ontogenetic changes in idiosomal and leg chaetotaxy from larva to adult. A key to the world species of *Acaricis* is also proposed.

**Key words:** Flat mite, false spider mite, new species, Cyperaceae, *Gahnia lacera*, ontogenetic changes

### Introduction

The Tenuipalpidae, commonly known as false spider mites or flat mites, has been considered as an under-explored family of plant-feeding mites of great economic importance (Gerson 2008). Recent increase in the interest in these mites was attributed to their biosecurity importance (as invasive species as well as vectors of virus diseases) and their use as model organisms to explore basic evolutionary questions such as sex determination and microorganism-mite interactions (Zhang 2010). Recent counts of the number of described species included 891 species (Mesa *et al.* 2009) and 895 species (Zhang *et al.* 2011), with several new species being described every year. In New Zealand, Spain and Luxton (1971) listed 11 described species, Zhang and Rhode (2003) estimated 23 species, and most recently, Sirvid *et al.* (2010) included 27 species of the Tenuipalpidae in the *New Zealand Inventory of Biodiversity*. There have been few papers on the Tenuiplapidae of New Zealand in the last decade (Zhang & Fan 2004; Berry & Fan 2012). This paper represents one of a series of papers to describe new species of this family and revise poorly described species from New Zealand.

The genus *Acaricis* was erected by Beard and Gerson (2009) with only two species: the type species *Acaricis plana* collected on *Scleria sphaelata* from Queensland, and *Acaricis danutae* collected on *Carex* sp. from New South Wales, Australia. Both host plant species are of plant family Cyperaceae. Currently, only 9 species belonging to 5 genera of the Tenuipalpidae have been described from Cyperaceae in the world. Wang (1983) described a new species *Tenuipalpus obvelatus* on Cyperaceae from China. Beard, Fan and Walter (2005) reported a new genus *Prolixus* with two new species collected on Cyperaceae in Australia. Beard and Ochoa (2011) added two new

genera *Gahniacarus* and *Cyperacarus* (each genus with two species from Australia) to the list. Prior to this study, *Acaricis* was believed to be endemic to Australia. In this paper, a new species, *Acaricis urigersoni* **sp. nov.**, is described from *Gahnia lacera* (Cyperaceae) in New Zealand and its ontogenetic development in body and leg chaetotaxy is studied. A key to world species of *Acaricis* is also proposed.

## Material and methods

Mites were mounted in Hoyer's medium, and examined at 1000 times with a DIC Nikon E800 microscope. All measurements were made from slide-mounted specimens using a stage-calibrated ocular ruler and are given in micrometers ( $\mu\text{m}$ ) (Zhang & Fan 2004). Measurements of the holotype are presented, followed by a range of paratypes in brackets. Setae were measured from the centre of the setal base to the tip of the seta; distances between setae were measured as the distance from the centre of one setal base to the other. Legs were measured from the basal end of trochanter to the distal end of tarsus. Terminology follows that applied to the Tetranychidae by Lindquist (1985).

## Family Tenuipalpidae Berlese, 1913

### Genus *Acaricis* Beard & Gerson, 2009

*Acaricis* Beard & Gerson, 2009: 2; Beard & Ochoa, 2011: 32.

Type species: *Acaricis plana* Beard & Gerson, 2009

**Diagnosis.** Body elongate, more than twice as long as wide; palpus usually 4-segmented. Prodorsum with a rostral shield and 2 or 3 pairs of setae ( $sc_1$  and  $sc_2$  or  $v_2$ ,  $sc_1$  and  $sc_2$ ); setae  $sc_2$  lanceolate. Opisthosoma with 9 pairs of dorsal setae; dorsosublateral setae absent; dorsocentral setae  $c_1$  absent,  $d_1$  and  $e_1$  present; humeral setae  $c_3$  and dorsolateral setae 6 pairs ( $d_3$ ,  $e_3$ ,  $f_2$ ,  $f_3$ ,  $h_2$  and  $h_1$ ); setae  $h_2$  long and attenuate. Genua IV nude.

The original concept of *Acaricis* was based on two Australian species (Beard & Gerson, 2009). The new species from New Zealand with new morphological features allows revision of the generic diagnosis as presented above.

### *Acaricis urigersoni* **sp. nov.**


(Figs. 1–24)

**Diagnosis.** The adult female of *Acaricis urigersoni* **sp. nov.** is easily distinguishable from those of the two other species by the absence of the first pair of prodorsal setae  $v_2$ , setae  $sc_1$  bifurcated, setae  $e_3$  lanceolate, cylinder-shaped spermathecal vesicle, and chaetotaxy (legs I–IV): femora 4-4-2-1, genua 2-2-0-0, tibiae 5-5-3-2.


The adult male of *Acaricis urigersoni* **sp. nov.** is differentiated from those of the two other species by the absence of the first pair of prodorsal setae  $v_2$ , setae  $e_3$  lanceolate, setae  $4a_2$  elongate and flagelliform, and Chaetotaxy (legs I–IV): femora 4-4-2-1, genua 2-2-0-0, tibiae 5-5-3-2, tarsi  $7+2\omega-7+2\omega-5+\omega-5+\omega$ .

**Type specimens.** **Holotype** ♀. **New Zealand**, Auckland, Kepa Reserve, 5 Aug, 2013, by Nicholas A. Martin, ex. *Gahnia lacera* (Cyperaceae). **Paratypes.** 6 females, 5 males, 7 deutonymphs, 4 protonymphs, 3 larvae, same data as holotype. 1 female, 1 male, 2 protonymphs, 3


larvae, Wenderholm Regional Park, Auckland, New Zealand, 27 Sep, 2013, ex. *Gahnia lacera* (Cyperaceae), by Nicholas A. Martin. 8 females, 4 males, 4 deutonymphs, 2 protonymphs, Kapa Reserve, Auckland, New Zealand, 9 Oct, 2013, ex. *Gahnia lacera* (Cyperaceae), by Nicholas A. Martin. The holotype and 39 paratypes (12 females, 8 males, 9 deutonymphs, 6 protonymphs, 4 larvae) will be deposited in the New Zealand Arthropod Collection, Landcare Research, Auckland, New Zealand (NZAC); 10 paratypes (2 females, 2 males, 2 deutonymphs, 2 protonymphs, and 2 larvae) will be deposited in the Natural History Museum, London (BMNH).




**FIGURE 1.** *Acaricis urigersoni* sp. nov. (adult female). A, dorsal view of idiosoma; B, ventral view of idiosoma.




**FIGURE 2.** *Acaricis urigersoni* sp. nov. (adult female). A, dorsal setae; B, ventral setae; C, chelicerae and collar.



**FIGURE 3.** *Acaricis urigersoni* sp. nov. (adult female). A, subcapitulum; B, genital-anal plate.



**FIGURE 4.** *Acaricis urigersoni* sp. nov. (adult female). A, leg I; B, leg II.



**FIGURE 5.** *Acaricis urigersoni* sp. nov. (adult female). A, leg III; B, leg IV.



#### Adult Female (n=5)

**Gnathosoma.** (Fig. 3A) Rostrum reaching proximal end of femur I, subcapitular seta *m* setiform,  $m=16-18$  (18),  $m-m=16-18$  (18). Palp 4-segmented, setal formula: 0, 0, 2, 2; tarsus with 2 eupathidia 4, 4.

**Idiosoma.** (Figs. 1A, 2A, 20A, 21A) 445–470 (470) long, 195–205 (200) wide. Rostral shield pitted. Idiosoma widest at posterior margin of propodosoma, narrowing abruptly at base of opisthosoma, then expanding posteriorly. PRODORSUM covered with weak longitudinal sculpturing and faint broad band of transverse striations between setae  $sc_1$  and  $d_1$  (Figs. 1A, 20A); bearing 2 pairs of setae ( $sc_1$  and  $sc_2$ ), setae  $sc_1$  bifurcated,  $sc_2$  lanceolate, barbed; lengths:  $sc_1$  5–6 (5),  $sc_2$  31–36 (35); distances:  $sc_1-sc_1$  70–74 (73),  $sc_1-sc_2$  40–49 (43),  $sc_2-sc_2$  155–165 (155). HYSTEOSOMA smooth, covered with oblique and longitudinal wrinkles as shown in Fig. 21A, a pair of distinct pores anterior to  $e_1$ ; with 1 pair of humeral setae  $c_3$ , 2 pairs of dorsocentral setae  $d_1$  and  $e_1$ , and 6 pairs of dorsolateral setae. Setae  $c_3$ ,  $d_1$ ,  $d_3$  and  $e_1$  setiform, dorsolateral setae  $e_3$ ,  $f_2$ ,  $f_3$  and  $h_1$  lanceolate, barbed, and  $h_2$  elongate, ending in a minute club. Lengths:  $d_1$  4–6 (4),  $e_1$  4,  $c_3$  4,  $d_3$  3–4 (3),  $e_3$  15–20 (20),  $f_2$  23–25 (25),  $f_3$  23–25 (25),  $h_2$  117–142 (117),  $h_1$  21–25 (21); distances:  $d_1-d_1$  24–28 (24),  $e_1-e_1$  10–20 (17),  $c_3-c_3$  160–170 (165),  $d_3-d_3$  120–130 (125),  $d_3-e_3$  110–130 (125),  $e_3-e_3$  120–135 (125),  $e_3-f_2$  23–24 (24),  $f_2-f_2$  115–125 (115),  $f_2-f_3$  19–22 (21),  $f_3-f_3$  95–105 (95),  $f_3-h_2$  21–27 (22),  $h_2-h_2$  (34),  $h_2-h_1$  13–14 (13),  $h_1-h_1$  33–42 (34).

**Venter.** (Figs. 1B, 2B, 3B, 22A, 23A, B) Venter with few broken oblique striae laterally and fine transverse striations between coxae II and IV. All coxal setae setiform. Setae  $1a$  and posterior medioventral setae  $4a_1$  and  $4a_2$  flagelliform, anterior medioventral setae  $3a$  setiform; setae  $4a_1$  and  $4a_2$  subequal, more than 8 times as long as  $3a$  (Fig. 2B). Lengths:  $1a$  130–145 (135),  $1b$  10–11 (11),  $1c$  10–12 (11),  $2b$  17–19 (18),  $2c$  18–20 (18),  $3a$  13–15 (13),  $3b$  16–20 (18),  $4a_1$  120–130 (125),  $4a_2$  115–125 (115),  $4b$  15–20 (18). Distances:  $1a-1a$  25–33 (30),  $3a-3a$  30–32 (30),  $4a_1-4a_1$  9–13 (9),  $4a_1-4a_2$  11–15 (12),  $4a_2-4a_2$  29–39 (29). Genital and ventral plates with flaps of cuticle as shown in Fig. 1B, bearing 1 pair of aggenital setae  $ag$  and 2 pairs of genital setae ( $g_1$  and  $g_2$ ), setiform; setae  $g_1$  and  $g_2$  subequal. Anal plate with 2 pairs of pseudanal setae ( $ps_1$  and  $ps_2$ ). Seta lengths:  $ag$  13–16 (14),  $g_1$  15–18 (16),  $g_2$  15–18 (16),  $ps_1$  12–14 (12),  $ps_2$  10–12 (10). Distances:  $ag-ag$  22–28 (22),  $g_1-g_1$  20–29 (24),  $g_1-g_2$  10–12 (11),  $g_2-g_2$  40–44 (40),  $ps_1-ps_2$  36–37 (37).

**Spermatheca.** (Figs. 3B, 22A, 23B) A short, narrow, unsclerotised tube extending from genital opening (mesad setae  $ps_2$ ) and ending in a cylinder-shaped vesicle.


**Legs.** (Figs. 4, 5) Lengths of legs I–IV: 125–135 (130), 100–110 (105), 82–87 (85), 80–84 (84). Chaetotaxy: coxae 2-2-1-1; trochanters 1-1-1-0; femora 4-4-2-1; genua 2-2-0-0, tibiae 5-5-3-2, tarsi 7+ $\omega$ -7+ $\omega$ -5-5. Most dorsal and lateral setae on femora, genua and tibiae lanceolate, lateral setae  $l'$  on tibiae pectinate; ventral setae  $bv''$  on femur I pectinate and on femur II lanceolate; seta  $ev'$  on femur III setiform; ventral setae  $v$  pectinate. Setae  $ft''$  on tarsi I–II and  $ft''\zeta$  on tarsi III–IV flagelliform; unguinal setae  $u$  pectinate and equal in length; proral setae  $p'\zeta$  and  $p''\zeta$  eupathidial; tectal setae  $tc$  setiform except on tarsus IV pectinate. Lengths of solenidia: I  $\omega''$  4–5 (4), II  $\omega''$  4–5 (4). Claws developed with tenent hairs on each side.

#### Male (n=5)


**Gnathosoma.** (Fig. 7C) Rostrum reaching proximal end of femur I, and rostral shield pitted, subcapitular setae  $m$  setiform,  $m=14-18$ ,  $m-m=14-18$ . Palp 4-segmented, setal formula: 0, 0, 2, 2; tibia with 2 bare setae, tarsus with 2 eupathidia, lengths 3–4, 3–4.

**Idiosoma.** (Figs. 6A, 7A, 20B, 21B) 325–340 long, 140–155 wide. Rostral shield pitted. Idiosoma widest at posterior margin of propodosoma, narrowing abruptly at base of opisthosoma, then expanding posteriorly. PRODORSUM covered with faint broad band of transverse striations between setae  $sc_1$  and  $d_1$  (Figs. 6A, 20B); bearing 2 pairs of setae ( $sc_1$  and  $sc_2$ ), setae  $sc_1$  setiform,  $sc_2$  lanceolate, barbed; lengths:  $sc_1$  4–5,  $sc_2$  25–30; distances:  $sc_1-sc_1$  56–65,  $sc_1-sc_2$  34–38,  $sc_2-sc_2$


125–135. HYSTEROSOMA divided into metapodosoma and opisthosoma by narrow band of horizontal striations (Figs. 6A, 21B); with 1 pair of humeral setae  $c_3$ , 2 pairs of dorsocentral setae  $d_1$  and  $e_1$ , and 6 pairs of dorsolateral setae. Setae  $c_3$ ,  $d_1$ ,  $d_3$  and  $e_1$  setiform, dorsolateral setae  $e_3$ ,  $f_2$ ,  $f_3$  and  $h_1$  lanceolate, barbed, and  $h_2$  elongate, ending in a minute club. Lengths:  $d_1$  3–4,  $e_1$  3–4,  $c_3$  3–4,  $d_3$  3–4,  $e_3$  7–10,  $f_2$  18–19,  $f_3$  19–21,  $h_2$  117–122,  $h_1$  18–21; distances:  $d_1$ – $d_1$  15–22,  $e_1$ – $e_1$  10–13,  $c_3$ – $c_3$  120–130,  $d_3$ – $d_3$  80–91,  $d_3$ – $e_3$  90–100,  $e_3$ – $e_3$  70–81,  $e_3$ – $f_2$  15–20,  $f_2$ – $f_2$  70–82,  $f_2$ – $f_3$  12–18,  $f_3$ – $f_3$  66–78,  $f_3$ – $h_2$  12–18,  $h_2$ – $h_2$  45–53,  $h_2$ – $h_1$  10,  $h_1$ – $h_1$  24–36.




**FIGURE 6.** *Acaricis urigersoni* sp. nov. (adult male). A, dorsal view of idiosoma; B, ventral view of idiosoma.




**FIGURE 7.** *Acaricis urigersoni* sp. nov. (adult male). A, dorsal setae; B, ventral setae; C, subcapitulum; D, chelicerae and collar.



**FIGURE 8.** *Acaricis urigersoni* sp. nov. (adult male). A, genital-anal plate.



**FIGURE 9.** *Acaricis urigersoni* sp. nov. (adult male). A, leg I; B, leg II.



**FIGURE 10.** *Acaricis urigersoni* sp. nov. (adult male). A, leg III; B, leg IV.

**Venter.** (Figs. 6B, 7B, 8A, 22B, 23D) Venter with fine transverse striation between coxae II and IV, longitudinal striations in the genital area. All coxal setae setiform. Setae *1a* and posterior medioventral setae *4a<sub>1</sub>* and *4a<sub>2</sub>* flagelliform, anterior medioventral setae *3a* setiform; setae *4a<sub>1</sub>* and *4a<sub>2</sub>* subequal, more than 9 times as long as *3a* (Fig. 7B). Lengths: *1a* 135–150, *1b* 10–11, *1c* 9–12, *2b* 15–20, *2c* 16–20, *3a* 10–12, *3b* 18–20, *4a<sub>1</sub>* 100–110, *4a<sub>2</sub>* 105–110, *4b* 14–17. Distances: *1a–1a* 25–30, *3a–3a* 23–28, *4a<sub>1</sub>–4a<sub>1</sub>* 9–11, *4a<sub>1</sub>–4a<sub>2</sub>* 8–10, *4a<sub>2</sub>–4a<sub>2</sub>* 25–30. Genital and ventral plates with an aedeagus and seminal vesicle, bearing one pair of aggenital setae *ag* and 2 pairs of genital setae (*g<sub>1</sub>* and *g<sub>2</sub>*), setiform; setae *g<sub>1</sub>* and *g<sub>2</sub>* subequal. Anal plate with 2 pairs of pseudanal setae (*ps<sub>1</sub>* and *ps<sub>2</sub>*). Seta lengths: *ag* 15–18, *g<sub>1</sub>* 27–29, *g<sub>2</sub>* 27–30, *ps<sub>1</sub>* 11–12, *ps<sub>2</sub>* 7–8. Distances: *ag–ag* 11–18, *g<sub>1</sub>–g<sub>1</sub>* 15–19, *g<sub>1</sub>–g<sub>2</sub>* 10–11, *g<sub>2</sub>–g<sub>2</sub>* 35–38, *ps<sub>1</sub>–ps<sub>2</sub>* 7–14.

**Aedeagus.** (Figs. 8, 22B, 23D) A narrow, elongate, sclerotised aedeagus tapering to a point posteriorly (at genital opening); membranous duct running from sclerotised aedeagus to flared, lightly sclerotised, cone-shaped cup distally, appearing to open into a soft membranous vesicle.

**Legs.** (Figs. 9, 10) Lengths of legs I–IV: 110–120, 90–96, 72–85, 72–83. Chaetotaxy: coxae 2-2-1-1; trochanters 1-1-1-0; femora 4-4-2-1; genua 2-2-0-0, tibiae 5-5-3-2, tarsi 7+2 $\omega$ -7+2 $\omega$ -5+ $\omega$ -5+ $\omega$ . Most dorsal and lateral setae on femora, genua and tibiae lanceolate, lateral setae *l'* on tibiae I and II setiform; ventral setae *bv''* on femur I pectinate and on femur II lanceolate; seta *ev'* on femur III setiform; ventral setae *v* pectinate except on trochanter II setiform. Setae *ft''* on tarsi I–II and *ft''* $\zeta$  on tarsi III–IV flagelliform; unguinal setae *u* pectinate and equal in length; proral setae *p'\zeta* and *p''\zeta* eupathidial; tectal setae *tc* setiform except *tc'* on tarsi I and II bifurcated. Lengths of solenidia: I  $\omega'$  8–9,  $\omega''$  6–8, II  $\omega'$  7–8,  $\omega''$  6, III  $\omega'$  7, IV  $\omega'$  7. Claws developed with tenent hairs on each side.

#### Deutonymph (n=5)


**Gnathosoma.** (Fig. 12A) Rostrum reaching proximal end of femur I, subcapitular seta *m* setiform, *m*=13–15, *m–m*=13–15. Palp 4-segmented, setal formula: 0, 0, 2, 2; tibia with 2 bare setae, tarsus with 2 eupathidia 3–4, 3–4.

**Idiosoma.** (Figs. 11A, 12C, 20C, 21C) 310–365 long, 140–160 wide. Rostral shield pitted. Idiosoma widest at posterior margin of propodosoma, narrowing abruptly at base of opisthosoma, then expanding posteriorly. PRODORSUM covered with faint broad band of transverse striations between setae *sc<sub>1</sub>* and *d<sub>1</sub>* (Figs. 11A, 20C), bearing 2 pairs of setae, setae *sc<sub>1</sub>* bifurcated, *sc<sub>2</sub>* lanceolate, barbed; lengths: *sc<sub>1</sub>* 3, *sc<sub>2</sub>* 21–28; distances: *sc<sub>1</sub>–sc<sub>1</sub>* 56–63, *sc<sub>1</sub>–sc<sub>2</sub>* 32–39, *sc<sub>2</sub>–sc<sub>2</sub>* 115–130. HYSTEROSOMA smooth, covered with oblique and longitudinal wrinkles as shown in Fig. 11A. Setae *c<sub>3</sub>*, *d<sub>1</sub>*, *d<sub>3</sub>* and *e<sub>1</sub>* setiform, subequal, dorsolateral setae *e<sub>3</sub>*, *f<sub>2</sub>*, *f<sub>3</sub>* and *h<sub>1</sub>* lanceolate, barbed, and *h<sub>2</sub>* elongate, ending in minute club. Lengths: *d<sub>1</sub>* 2–3, *e<sub>1</sub>* 2–4, *c<sub>3</sub>* 2–3, *d<sub>3</sub>* 2–3, *e<sub>3</sub>* 5–7, *f<sub>2</sub>* 12–16, *f<sub>3</sub>* 12–17, *h<sub>2</sub>* 86–105, *h<sub>1</sub>* 12–17; distances: *d<sub>1</sub>–d<sub>1</sub>* 15–21, *e<sub>1</sub>–e<sub>1</sub>* 6–10, *c<sub>3</sub>–c<sub>3</sub>* 110–125, *d<sub>3</sub>–d<sub>3</sub>* 81–100, *d<sub>3</sub>–e<sub>3</sub>* 57–86, *e<sub>3</sub>–e<sub>3</sub>* 73–95, *e<sub>3</sub>–f<sub>2</sub>* 11–16, *f<sub>2</sub>–f<sub>2</sub>* 72–95, *f<sub>2</sub>–f<sub>3</sub>* 11–16, *f<sub>3</sub>–f<sub>3</sub>* 60–85, *f<sub>3</sub>–h<sub>2</sub>* 12–20, *h<sub>2</sub>–h<sub>2</sub>* 40–51, *h<sub>2</sub>–h<sub>1</sub>* 10–13, *h<sub>1</sub>–h<sub>1</sub>* 19–26.

**Venter.** (Figs. 11B, 12C, 14C, 22C) Venter with fine transverse striations between coxae II and IV. All coxal setae setiform. Setae *1a* and posterior medioventral setae *4a<sub>1</sub>* flagelliform, anterior medioventral setae *3a* setiform; setae *4a<sub>1</sub>* about 6 times as long as *3a* (Fig. 12C). Lengths: *1a* 125–140, *1b* 8–10, *1c* 8–10, *2b* 9–13, *2c* 11–14, *3a* 10–14, *3b* 14–16, *4a<sub>1</sub>* 79–90, *4b* 12–13. Distances: *1a–1a* 21–23, *3a–3a* 21–24, *4a<sub>1</sub>–4a<sub>1</sub>* 12–19. Genital and ventral plates with oblique cuticles anterior to *ag* and transverse cuticles between *ag* and *g<sub>1</sub>*, and oblique striations laterad seta *g<sub>1</sub>* as shown in Fig. 22C, bearing one pair of aggenital setae *ag* and one pair of genital setae *g<sub>1</sub>*, setiform. Anal plate with 2 pairs of pseudanal setae (*ps<sub>1</sub>* and *ps<sub>2</sub>*). Seta lengths: *ag* 10–14, *g<sub>1</sub>* 11–14, *ps<sub>1</sub>* 8–9, *ps<sub>2</sub>* 5–7. Distances: *ag–ag* 14–23, *g<sub>1</sub>–g<sub>1</sub>* 13–23, *ps<sub>1</sub>–ps<sub>2</sub>* 18–30.


**Legs.** (Figs. 13, 14A, B) Lengths of legs I–IV: 84–96, 66–77, 53–61, 48–55. Chaetotaxy: coxae 2-2-1-1; trochanters 1-1-1-0; femora 4-4-2-1; genua 2-2-0-0, tibiae 5-5-3-2, tarsi 7+ $\omega$ -7+ $\omega$ -5-5. Most dorsal and lateral setae on femora, genua and tibiae lanceolate, lateral setae *l''* on genua I

pectinate setiform, lateral setae  $l'$  on tibiae I and II setiform. Ventral seta  $bv''$  on femur I setiform and on femur II lanceolate; ventral seta  $v$  and  $ev'$  setiform. Setae  $ft''$  on tarsi I–II and  $ft'\zeta$  on tarsi III–IV flagelliform; unguinal setae  $u$  setiform and equal in length; proral setae  $p'\zeta$  and  $p''\zeta$  eupathidial; tectal setae  $tc$  setiform. Lengths of solenidia: I  $\omega''$  4, II  $\omega''$  4. Claws developed with tenent hairs on each side.




**FIGURE 11.** *Acaricis urigersoni* sp. nov. (deutonymph). A, dorsal view of idiosoma; B, ventral view of idiosoma.






**FIGURE 12.** *Acaricis urigersoni* sp. nov. (deutonymph). A, subcapitulum; B, ventral view of idiosoma; C, dorsal view of idiosoma; D, chelicerae and collar.




**FIGURE 13.** *Acaricis urigersoni* sp. nov. (deutonymph). A, leg I; B, leg II.




**FIGURE 14.** *Acaricis urigersoni* sp. nov. (deutonymph). A, leg III; B, leg IV; C, genital-anal plate.

**Protonymph** (n=4)


**Gnathosoma.** (Fig. 16D) Rostrum reaching proximal end of femur I, and rostral shield pitted, subcapitular seta *m* setiform, *m*=14–16, *m*-*m*=14–16. Palp 4-segmented, setal formula: 0, 0, 2, 2; tibia with 2 bare setae, tarsus with 2 eupathidia 3–4, 3–4.



**FIGURE 15.** *Acaricis urigersoni* sp. nov. (protonymph). A, dorsal view of idiosoma; B, ventral view of idiosoma.



**FIGURE 16.** *Acaricis urigersoni* sp. nov. (protonymph). A, ventral setae; B, dorsal setae; C, chelicerae and collar; D, subcapitulum; E, genital-anal plate.



**FIGURE 17.** *Acaricis urigersoni* sp. nov. (protonymph). A, leg I; B, leg II; C, leg III; D, leg IV.

**Idiosoma.** (Figs. 15A, 16B, 20D, 21D) 279–305 long, 120–130 wide. Rostral shield pitted. Idiosoma widest at posterior margin of propodosoma, narrowing abruptly at base of opisthosoma, then expanding posteriorly. **PRODORSUM** covered with faint broad band of transverse striations between setae  $sc_1$  and  $d_1$  (Figs. 16B), bearing 2 pairs of setae, setae  $sc_1$  setiform,  $sc_2$  lanceolate, barbed; lengths:  $sc_1$  3,  $sc_2$  20–21; distances:  $sc_1$ – $sc_1$  45–57,  $sc_1$ – $sc_2$  28–33,  $sc_2$ – $sc_2$  105–115. **HYSTEOSOMA** smooth (Figs. 15A, 21D), with 1 pair of humeral seta  $c_3$ , 2 pairs of dorsocentral setae  $d_1$  and  $e_1$ , and 6 pairs of dorsolateral setae. Setae  $c_3$ ,  $d_1$ ,  $d_3$  and  $e_1$  setiform, subequal,

dorsolateral setae  $e_3$ ,  $f_2$ ,  $f_3$  and  $h_1$  lanceolate, barbed, and  $h_2$  elongate, ending in minute club. Lengths:  $d_1$  2,  $e_1$  2–4,  $c_3$  2,  $d_3$  2,  $e_3$  3–5,  $f_2$  9–13,  $f_3$  10–12,  $h_2$  68–82,  $h_1$  9–11; distances:  $d_1$ – $d_1$  15–24,  $e_1$ – $e_1$  6–9,  $c_3$ – $c_3$  93–99,  $d_3$ – $d_3$  68–75,  $d_3$ – $e_3$  55–58,  $e_3$ – $e_3$  63–73,  $e_3$ – $f_2$  8–15,  $f_2$ – $f_2$  59–66,  $f_2$ – $f_3$  9–13,  $f_3$ – $f_3$  46–54,  $f_3$ – $h_2$  8–14,  $h_2$ – $h_2$  29–37,  $h_2$ – $h_1$  9–10,  $h_1$ – $h_1$  11–20.

**Venter.** (Figs. 15B, 16A, E, 22D) Venter similar to deutonymph. All coxal setae setiform. Setae *Ia* flagelliform and anterior medioventral setae *3a* setiform (Fig. 16A). Lengths: *Ia* 98–105, *Ib* 8–10, *Ic* 8–9, *2c* 11–12, *3a* 9–10, *3b* 11–14. Distances: *Ia*–*Ia* 19–22, *3a*–*3a* 21–25. Similar to deutonymph, bearing one pair of aggenital setae *ag*, setiform. Anal plate with 2 pairs of pseudanal setae (*ps*<sub>1</sub> and *ps*<sub>2</sub>). Setae lengths: *ag* 7–12, *ps*<sub>1</sub> 6–7, *ps*<sub>2</sub> 5–6. Distances: *ag*–*ag* 15–16, *ps*<sub>1</sub>–*ps*<sub>2</sub> 14–17.

**Legs.** (Fig. 17) Lengths of legs I–IV: 67–74, 55–62, 48–50, 41–44. Chaetotaxy: coxae 2-1-1-0; trochanters 0-0-1-0; femora 3-3-2-1; genua 1-1-0-0, tibiae 5-5-3-2, tarsi 7+ $\omega$ -7+ $\omega$ -5-3. Dorsal setae on femora, genua and tibiae lanceolate; ventral seta *bv*'' on femur I setiform and on femur II lanceolate; lateral setae *l*' on tibiae I and II setiform and *l*'' on tibiae I and II lanceolate; ventral seta *v* setiform. Setae *ft*'' on tarsi I–II and *ft*' $\zeta$  on tarsi III–IV flagelliform; unguinal seta *u* and tectal seta *tc* setiform; proral setae *p*' $\zeta$  and *p*'' $\zeta$  eupathidial. Lengths of solenidia: I  $\omega$ '' 3, II  $\omega$ '' 3. Claws developed with tenent hairs on each side.

**Larva** (n=5)

**Gnathosoma.** (Fig. 19D) Rostrum reaching middle of femur I, subcapitular seta *m* setiform, *m*=14–17, *m*–*m*=14–17. Palp 4-segmented, setal formula: 0, 0, 2, 2; tibia with 2 bare setae, tarsus with 2 eupathidia 3–4, 3–4.

**Idiosoma.** (Figs. 18A, 19F, 20E, 21E) 240–275 long, 85–110 wide. PRODORSUM covered with faint broad band of transverse striations between setae *sc*<sub>1</sub> and *d*<sub>1</sub> (Figs. 18A, 20E), bearing 2 pairs of setae, setae *sc*<sub>1</sub> setiform, *sc*<sub>2</sub> lanceolate, barbed; lengths: *sc*<sub>1</sub> 2–4, *sc*<sub>2</sub> 12–16; distances: *sc*<sub>1</sub>–*sc*<sub>1</sub> 45–47, *sc*<sub>1</sub>–*sc*<sub>2</sub> 20–25, *sc*<sub>2</sub>–*sc*<sub>2</sub> 82–96. HYSTEOSOMA smooth (Figs. 18A, 21E), with 1 pair of humeral seta *c*<sub>3</sub>, 2 pairs of dorsocentral setae *d*<sub>1</sub> and *e*<sub>1</sub>, and 6 pairs of dorsolateral setae. Setae *c*<sub>3</sub>, *d*<sub>1</sub>, *d*<sub>3</sub>, *e*<sub>1</sub> and *e*<sub>3</sub> setiform, subequal, dorsolateral setae *f*<sub>2</sub>, *f*<sub>3</sub> and *h*<sub>1</sub> lanceolate, barbed, and *h*<sub>2</sub> elongate, ending in minute club. Lengths: *d*<sub>1</sub> 3–4, *e*<sub>1</sub> 2, *c*<sub>3</sub> 2, *d*<sub>3</sub> 2, *e*<sub>3</sub> 4, *f*<sub>2</sub> 8–9, *f*<sub>3</sub> 8–9, *h*<sub>2</sub> 89–98, *h*<sub>1</sub> 8; distances: *d*<sub>1</sub>–*d*<sub>1</sub> 15–18, *e*<sub>1</sub>–*e*<sub>1</sub> 6–8, *c*<sub>3</sub>–*c*<sub>3</sub> 70–82, *d*<sub>3</sub>–*d*<sub>3</sub> 45–53, *d*<sub>3</sub>–*e*<sub>3</sub> 35–43, *e*<sub>3</sub>–*e*<sub>3</sub> 50–60, *e*<sub>3</sub>–*f*<sub>2</sub> 7–9, *f*<sub>2</sub>–*f*<sub>2</sub> 45–55, *f*<sub>2</sub>–*f*<sub>3</sub> 7–8, *f*<sub>3</sub>–*f*<sub>3</sub> 35–44, *f*<sub>3</sub>–*h*<sub>2</sub> 8–10, *h*<sub>2</sub>–*h*<sub>2</sub> 22–27, *h*<sub>2</sub>–*h*<sub>1</sub> 7–8, *h*<sub>1</sub>–*h*<sub>1</sub> 8–11.

**Venter.** (Figs. 18B, 19E, 22E) Venter similar to deutonymph. All coxal setae setiform. Setae *Ia* flagelliform and anterior medioventral setae *3a* setiform (Fig. 19E). Lengths: *Ia* 82–93, *Ib* 7–8, *3a* 8–9. Distances: *Ia*–*Ia* 18–25, *3a*–*3a* 21–26. Anal plate with 2 pairs of pseudanal setae (*ps*<sub>1</sub> and *ps*<sub>2</sub>). Setae lengths: *ps*<sub>1</sub> 5–6, *ps*<sub>2</sub> 4–5. Distances: *ps*<sub>1</sub>–*ps*<sub>2</sub> 8–10.

**Legs.** (Figs. 19A, B, C) Lengths of legs I–III: 50–55, 44–48, 41–44. Chaetotaxy: coxae 1-0-0; trochanters 0-0-0; femora 3-3-2; genua 1-1-0, tibiae 5-5-3, tarsi 5+ $\omega$ -5+ $\omega$ -3. Most dorsal setae on femora, genua and tibiae lanceolate except on tibiae I and II nude. Ventral setae *bv*'' on femora and lateral seta *l*' on tibiae setiform; lateral seta *l*'' on tibiae I and II lanceolate; ventral seta *v* setiform. Setae *ft*'' on tarsi I–II and *ft*' $\zeta$  on tarsus III flagelliform; unguinal seta *u* and tectal seta *tc* setiform; proral setae *p*' $\zeta$  and *p*'' $\zeta$  eupathidial. Lengths of solenidia: I  $\omega$ '' 2, II  $\omega$ '' 2. Claws developed with tenent hairs on each side.

**Etymology.** The species is named after Prof Uri Gerson, who is an acarologist from Israel and has made significant contributions to the systematics and biology of the Tenuipalpidae.

**Remarks.** In live specimens, the body of this new species looks glossy with a translucent appearance, and is yellow-greenish with a series of minute black spots; legs I–II are obviously orange, and legs III–IV are pale orange; the eyes are red (Fig. 24). In one adult male, one dorsolateral setae *h*<sub>2</sub> was duplicated (*h*<sub>2</sub> and *h*<sub>2x</sub> in Fig. 23C), and shortened by three times; the other seta *h*<sub>2</sub> was normal.






FIGURE 18. *Acaricis urigersoni* sp. nov. (larva). A, dorsal view of idiosoma; B, ventral view of idiosoma.






**FIGURE 19.** *Acaricis urigersoni* sp. nov. (larva). A, leg I; B, leg II; C, leg III; D, subcapitulum; E, ventral setae; F, dorsal setae.




**FIGURE 20.** *Acaricis urigersoni* sp. nov. (prodorsum). A, female; B, male; C, deutonymph; D, protonymph; E, larva.






**FIGURE 21.** *Acaricis urigersoni* sp. nov. (hysterosoma). A, female; B, male; C, deutonymph; D, protonymph; E, larva.






**FIGURE 22.** *Acaricis urigersoni* sp. nov. (genital-anal plate). A, female; B, male; C, deutonymph; D, protonymph; E, larva.





**FIGURE 23.** *Acaricis urigersoni* sp. nov. (adult). A, female genital plate; B, spermatheca; C, hysterosoma with 2  $h_2$ ; D, Aedeagus.





**FIGURE 24.** *Acaricis urigersoni* **sp. nov.** (dorsal view of idiosoma). A, female; B, male; C, deutonymph; D, protonymph; E, larva; F, egg.

The ontogeny of *Acaricis* was reported by Beard and Gerson (2009). Based on the observation of *Acaricis plana* and *Acaricis urigersoni* **sp. nov.**, the dorsal idiosomal setae ( $sc_1$ ,  $sc_2$ ,  $d_1$ ,  $e_1$ ,  $c_3$ ,  $d_3$ ,  $e_3$ ,  $f_2$ ,  $f_3$ ,  $h_2$  and  $h_1$ ) and pseudanal setae ( $ps_1$  and  $ps_2$ ) are present throughout all stages.

**TABLE 1.** Ontogeny of leg chaetotaxy in *Acaricis plana* (Ap) and *A. urigersoni* (Au). Setae are indicated where they are first added. Setae in parentheses represent pairs. Hyphen indicates no additions.

	Coxae	Trochanters	Femora	Genua	Tibiae	Tarsi
<b>Leg I</b>						
Larva—Ap	<i>1b</i>	-	<i>d, v', bv''</i>	<i>d</i>	<i>d, (l), (v)</i>	<i>(u), p'ζ, p''ζ, ft'', ω''</i>
Au	<i>1b</i>	-	<i>d, v', bv''</i>	<i>d</i>	<i>d, (l), (v)</i>	<i>(u), p'ζ, p''ζ, ft'', ω''</i>
Protonymph—Ap	<i>1c</i>	-	-	-	-	<i>(tc)</i>
Au	<i>1c</i>	-	-	-	-	<i>(tc)</i>
Deutonymph—Ap	-	<i>v'</i>	<i>l'</i>	<i>l''</i>	-	-
Au	-	<i>v'</i>	<i>l'</i>	<i>l''</i>	-	-
Female—Ap	-	-	-	-	-	-
Au	-	-	-	-	-	-
Male—Ap	-	-	-	-	-	<i>ω'</i>
Au	-	-	-	-	-	<i>ω'</i>
<b>Leg II</b>						
Larva—Ap	-	-	<i>d, v', bv''</i>	<i>d</i>	<i>d, (l), (v)</i>	<i>(u), p'ζ, p''ζ, ft'', ω''</i>
Au	-	-	<i>d, v', bv''</i>	<i>d</i>	<i>d, (l), (v)</i>	<i>(u), p'ζ, p''ζ, ft'', ω''</i>
Protonymph—Ap	<i>2c</i>	-	-	-	-	<i>(tc)</i>
Au	<i>2c</i>	-	-	-	-	<i>(tc)</i>
Deutonymph—Ap	<i>2b</i>	<i>v'</i>	<i>l'</i>	<i>l''</i>	-	-
Au	<i>2b</i>	<i>v'</i>	<i>l'</i>	<i>l''</i>	-	-
Female—Ap	-	-	-	-	-	-
Au	-	-	-	-	-	-
Male—Ap	-	-	-	-	-	<i>ω'</i>
Au	-	-	-	-	-	<i>ω'</i>
<b>Leg III</b>						
Larva—Ap	-	-	<i>d, ev'</i>	-	<i>d, (v)</i>	<i>(u), ft' ζ</i>
Au	-	-	<i>d, ev'</i>	-	<i>d, (v)</i>	<i>(u), ft' ζ</i>
Protonymph—Ap	<i>3b</i>	<i>l'</i>	-	-	-	<i>(tc)</i>
Au	<i>3b</i>	<i>l'</i>	-	-	-	<i>(tc)</i>
Deutonymph—Ap	-	-	-	-	-	-
Au	-	-	-	-	-	-
Female—Ap	-	-	-	-	-	-
Au	-	-	-	-	-	-
Male—Ap	-	-	-	-	-	<i>ω'</i>
Au	-	-	-	-	-	<i>ω'</i>
<b>Leg IV</b>						
Protonymph—Ap	-	-	<i>d, ev'</i>	-	<i>d, (v)</i>	<i>(u), ft' ζ</i>
Au	-	-	<i>d</i>	-	<i>(v)</i>	<i>(u), ft' ζ</i>
Deutonymph—Ap	<i>4b</i>	-	-	-	-	<i>(tc)</i>
Au	<i>4b</i>	-	-	-	-	<i>(tc)</i>
Female—Ap	-	-	-	-	-	-
Au	-	-	-	-	-	-
Male—Ap	-	-	-	-	-	<i>ω'</i>
Au	-	-	-	-	-	<i>ω'</i>

\* Leg IV absent in larva.

Ventral setae  $1c$ ,  $2c$  and  $3b$  are added in the protonymph,  $2b$ ,  $4a_1$  and  $4b$  added in the deutonymph, and  $4a_2$  added in the adult. Aggenital seta  $ag$  emerges in the protonymph; the first pair of genital setae  $g_1$  are formed in the deutonymph and the second pair appear in the adult. The palpal chaetotaxy (0, 0, 2, 2) remains unchanged from larva to adult. The changes in the chaetotaxy of leg segments are listed in Table 1. All dorsal setae  $d$  of legs lanceolate except on tibiae I and II in larva nude; ventral setae  $v$  in larva, protonymph and deutonymph setiform, in female pectinate, in male on leg I–II pectinate and on leg III–IV setiform; ventral seta  $ev'$  setiform; ventral seta  $bv'$  in adult pectinate and in all immature stages setiform; lateral setae  $l''$  in adult lanceolate, in larva, protonymph and deutonymph lanceolate except on genua I in deutonymph pectinate; unguinal seta  $u$  setiform in immature stages and pectinate in adult; tectal seta  $tc$  setiform and prorals setae  $p'\zeta$  and  $p''\zeta$  eupathidial.

### Key to species of *Acaricis* (based on adult females)

1. Setae  $v_2$  absent;  $sc_1$  bifurcate;  $e_3$  lanceolate; spermathecal vesicle cylinder-shaped; femora 4-4-2-1; genua 2-2-0-0; tibiae 5-5-3-2 . . . . . *A. urigersoni* **sp. nov.**
- Setae  $v_2$  present;  $sc_1$  setiform;  $e_3$  setiform or barbed; spermathecal vesicle dome-shaped; femora 4-4-2-2; genua 2-2-1-0; tibiae 5-5-3-3 . . . . . 2
2. Setae  $e_3$  setiform,  $sc_2$  lanceolate; dorsal seta  $d$  on femur III lanceolate . . . . . *A. plana* Beard & Gerson
- Setae  $e_3$  barbed,  $sc_2$  short and barbed; dorsal seta  $d$  on femur III short and stout . . . . . *A. danutae* Beard & Gerson

### Acknowledgements

We are grateful to Dr Nicholas A. Martin (Landcare Research, Auckland, New Zealand) for collecting the specimens of this new species and to Dr Qing-Hai Fan (Ministry for Primary Industries, Auckland, New Zealand) for assistance in taking photographs used in Figures 20–23. While this paper was prepared, Zhi-Qiang Zhang was supported by Core funding for Crown Research Institutes from the Ministry of Business, Innovation and Employment's Science and Innovation Group.

### References

- Beard, J.J., Fan, Q.-H. & Walter, D.E. (2005) A new genus and two new species of Tenuipalpidae (Prostigmata: Tetranychioidea) from an Australian sedge. *Acarologia*, 45, 161–181.
- Beard, J.J. & Gerson, U. (2009) A new flat mite genus, *Acaricis* (Prostigmata: Tenuipalpidae), from Australian sedges (Cyperaceae). *Zootaxa*, 2073, 31–44.
- Beard, J.J. & Ochoa, R. (2011) New flat mite genera (Acari: Trombidiformes: Tenuipalpidae) associated with Australian sedges (Cyperaceae). *Zootaxa*, 2941, 1–37.
- Berry, J.A. & Fan, Q.-H. (2012) Biological notes and risk status of *Brevipalpus phoenicis* (Geijskes, 1939) (Acari: Tenuipalpidae) in New Zealand. *Systematic & Applied Acarology*, 17(2), 224–230. <http://dx.doi.org/10.11158/saa.17.2.9>
- Gerson, U. (2008) The Tenuipalpidae: An under-explored family of plant-feeding mites. *Systematic & Applied Acarology*, 13, 83–101.
- Lindquist, E.E. (1985) External anatomy. In: Helle, W. & Sabelis, M.W. (Eds), *Spider Mites: Their Biology, Natural Enemies and Control*. Vol. 1A. Elsevier, Amsterdam, pp. 3–28.
- Mesa, N.C., Ochoa, R., Welbourn, W.C., Evans, G.A. & de Moraes, G.J. (2009) A catalog of the Tenuipalpidae (Acari) of the World with a key to genera. *Zootaxa*, 2098, 1–185.



- Sirvid, P.J., Zhang, Z.-Q., Harvey, M.S., Rhode, B.E., Cook, D.R., Bartsch, I. & Staples, D.A. (2010) Chapter Six: Phylum Arthropoda, Chelicerata. Horseshoe crabs, arachnids, sea spiders. *In*: Gordon, D P (Ed.), New Zealand inventory of biodiversity. Volume 2. Kingdom Animalia. Chaetognatha, Ecdysozoa, Ichno-fossils. Canterbury University Press, Christchurch, New Zealand, pp. 50–89.
- Spain, A.V. & Luxton, M. (1971) Catalog and bibliography of the Acari of the New Zealand subregion. *Pacific Insects Monograph*, 25, 179–226.
- Wang, H.-F. (1983) New species of the genus *Tenuipalpus* from China (Acarina: Tenuipalpidae). *Acta Zootaxonomica Sinica*, 8(1), 51–62.
- Zhang, Z.-Q. (2010) Tenuipalpidae of China: a review of progress. *Zoosymposia*, 4, 151–157
- Zhang, Z.-Q. & Fan, Q.-H. (2004) Redescription of *Dolichotetranychus ancistrus* Baker & Pritchard (Acari: Tenuipalpidae) from New Zealand. *Systematic & Applied Acarology*, 9, 111–131.
- Zhang, Z.-Q., Fan, Q.-H., Pesic, V., Smit, H., Bochkov, A.V., Khaustov, A.A., Baker, A., Wohltmann, A., Wen, T.-H., Amrine, J.W., Beron, P., Lin, J.-Z., Gabrys, G. & Husband, R. (2011) Order Trombidiformes Reuter, 1909. *In*: Zhang, Z.-Q. (Ed.) *Animal biodiversity: an outline of higher-level classification and survey of taxonomic richness*. *Zootaxa*, 3148, 129–138.

*Accepted by Qing-Hai Fan: 4 Dec. 2013; published 24 Dec. 2013*