

Tenuipalpus mansonculus (Acari: Tenuipalpidae) of Australia: redescription of the holotype

Authors: Xu, Yun, and Zhang, Zhi-Qiang

Source: Systematic and Applied Acarology, 19(3) : 322-328

Published By: Systematic and Applied Acarology Society

URL: <https://doi.org/10.11158/saa.19.3.6>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

***Tenuipalpus mansonculus* (Acari: Tenuipalpidae) of Australia: redescription of the holotype**

YUN XU¹ & ZHI-QIANG ZHANG^{2, 3, 4}

¹ Key Laboratory of Integrated Pest Management for Fujian-Taiwan Crops, Ministry of Agriculture; Fujian Provincial Key Laboratory of Insect Ecology; Fujian Agriculture and Forestry University, Fuzhou 350002, China.

² Landcare Research, 231 Morrin Road, Auckland, New Zealand.

³ Centre for Biodiversity & Biosecurity, School of Biological Sciences, University of Auckland, Auckland, New Zealand

⁴ Corresponding author: zhangz@landcareresearch.co.nz

Abstract

The holotype female of *Tenuipalpus mansonculus* Ghai & Shenhmar, 1984 (replacement name of *Colopalpus masoni* Collyer, 1973) is described and illustrated. The nomenclatural history of this species is documented and errors in the original description are corrected.

Introduction

Collyer (1973) described two new species of *Colopalpus*, including *Colopalpus masoni* from leaves of *Pomaderris* sp. imported from Australia. *Colopalpus* was synonymised with *Tenuipalpus* by Meyer (1979). As a result, *T. masoni* (Collyer 1973) became a homonym of *Tenuipalpus masoni* De Leon, 1965, and was later replaced with *Tenuipalpus mansonculus* Ghai & Shenhmar, 1984.

Tenuipalpus mansonculus is one of the six species of *Tenuipalpus* known from Australia (Smiley & Gerson 1995; Halliday 1998), but it is the least known species among these six species. Since its original description by Collyer (1973), no further information is available. In a recent study of the Tenuipalpidae from Australasia, we reexamined the holotype of this species. Since the original description by Collyer (1973) is brief and contains some errors, we herein redescribe the holotype to enable correct identification of this species in the future.

The specimen was examined at 1000 times using DIC Nikon E800 microscope. All measurements were made from the slide-mounted specimen using a stage-calibrated ocular ruler and are given in micrometers (µm) (Zhang & Fan 2004). Body length was measured from the anterior margin of the rostral shield to the posterior end of the idiosoma, and the body width was measured as the greatest width between legs I and II. Setae were measured from the centre of setal base to the tip of the seta; distances between setae were measured from the centre of one setal base to that of the other. Legs were measured from the base of the trochanter to the distal end of tarsus (excluding pretarsus). Terminology follows that of Lindquist (1985).

***Tenuipalpus mansonculus* Ghai & Shenhmar, 1984 (Figs. 1–4)**

Colopalpus masoni Collyer, 1973: 531, figs. 2–3.

Tenuipalpus masoni: Meyer, 1979: 5; Gutierrez & Schicha, 1982: 141.

Tenuipalpus mansoniculus Ghai & Shenhmar, 1985: 117; Smiley & Gerson, 1995: 41; Halliday, 1998: 84.

Material examined

Holotype female. Intercepted in New Zealand quarantine from *Pomaderris* sp. imported from Australia, 29 May, 1970, forwarded by C. A. F. Jaques to D. C. M. Manson. Originally deposited in Collection of Entomology Division, Department of Scientific and Industrial Research, Nelson, New Zealand, and now in New Zealand Arthropod Collection (NZAC), Landcare Research, Auckland, New Zealand.

Redescription of the holotype female

Gnathosoma. (Figs. 1B, 2) Rostrum reaching middle of femur I, subcapitular setae *m* setiform, $m=15$, $m-m=15$. Palp 1-segmented, tarsus with 1 spine-like seta, 18 long.

Idiosoma. (Fig. 1A) 245 long, 125 wide. Rostral shield pitted, with 1 median conical projection. PRODORSUM covered with irregular wrinkles mesally and laterally and a transverse sclerotized bar-like basally, bearing 3 pairs of setiform setae (v_2 , sc_1 and sc_2), sc_1 about twice as long as v_2 , and sc_2 about 3 times as long as v_2 ; lengths: v_2 5, sc_1 11, sc_2 15; distances: v_2-v_2 38, v_2-sc_1 21, sc_1-sc_1 73, sc_1-sc_2 23, sc_2-sc_2 110. HYSTEROSOMA covered with irregular wrinkles mesally and laterally; bearing 1 pair of pores, 1 pair of humeral setae (c_3), 3 pairs of dorsocentral setae (c_1 , d_1 and e_1), and 5 pairs of dorsolateral setae (d_3 , e_3 , f_3 , h_2 and h_1). All setae setiform except flagelliform h_2 . Lengths: c_1 6, d_1 5, e_1 7, c_3 7, d_3 4, e_3 9, f_3 11, h_2 205, h_1 8; distances: c_1-c_1 35, d_1-d_1 16, e_1-e_1 8, c_3-c_3 105, d_3-d_3 95, e_3-e_3 74, e_3-f_3 12, f_3-f_3 60, f_3-h_2 18, h_2-h_2 33, h_2-h_1 9, h_1-h_1 20.

Venter. (Fig. 2) Venter covered with oblique striae between coxae II–IV and transversal striations posterior to setae *1a*; oblique striations between setae *3a–3a* V-shaped; oblique striations between *3a* and *4a* also V-shaped. All coxal setae setiform. Setae *1a* flagelliform, middle medioventral setae *3a* and posterior medioventral setae *4a* setiform. Setae *3a* and *4a* subequal in length, *1a* longest, about 5 times as long as *3a*. Lengths: *1a* 96, *1b* 20, *1c* 14, *2b* 20, *2c* 17, *3a* 18, *3b* 17, *4a* 20, *4b* 18. Distances: *1a–1a* 26, *3a–3a* 41, *4a–4a* 31. Genital and ventral area with transversal striae mesally and oblique laterally as shown in Fig. 2, bearing 1 pair of aggenital setae (*ag*) and 2 pairs of genital setae (g_1 and g_2), setiform. Anal area with 2 pairs of pseudanal setae (ps_1 and ps_2), ps_1 about twice as long as ps_2 . Setae lengths: *ag* 14, g_1 19, g_2 16, ps_1 20, ps_2 11. Distances: *ag–ag* 16, g_1-g_1 11, g_1-g_2 8, g_2-g_2 23, ps_1-ps_2 6.

Legs. (Figs. 3–4) Lengths of legs I–IV: 115, 100, 92, 105. Chaetotaxy: coxae 2-2-1-1; trochanters 1-1-1-1; femora 4-4-2-1; genua 3-3-1-1; tibiae 5-5-3-3; tarsus 8+ ω -8+ ω -5-5. Most dorsal and lateral setae on trochanters, femora, genua and tibiae barbed, lateral setae *l'* and *l''* on femura I–II, genu I and tibiae I–II spine-like; ventral setae v' , ev' and bv'' on trochanters and femora setiform, setae v' and v'' on tibiae I–IV pectinate; Setae *ft'* on tarsi I–IV flagelliform, *ft''* barbed; ungual setae *u'* and *u''* pectinate and equal in length; tectal seta *tc'* and *tc''* on tarsus I–IV spine-like. Solenidion ω'' and eupathidium $p'\zeta$ and $p''\zeta$ on tarsi I–II rod-like. Lengths of solenidia: I ω'' 7, II ω'' 7. Claws and empodium reduced, pad-like, each with tenent hairs on each side.

Remarks

Collyer (1973) reported that only tarsus II in the female has a solenidion, but actually tarsi I and II each bears a solenidion. Collyer (1973) also mentioned four paratype females and two paratype males (three females sent to other institutions), but now only the holotype is preserved in NZAC. Collyer (1973) noted that males are similar to females but lack “the sclerotized bar” in the propodosomal area and smaller than females. She observed a solenidion each on tarsi I and II in males—this agrees with the pattern in the holotype female.




FIGURE 1. *Tenuipalpus mansonculus* Ghai & Shenhmar (female). A, dorsal view of idiosoma; B, subcapitulum.




FIGURE 2. *Tenuipalpus mansonicus* Ghai & Shenhmar (female). Ventral view of idiosoma.




FIGURE 3. *Tenuipalpus mansonculus* Ghai & Shenhmar (female). A, leg I; B, leg II.




FIGURE 4. *Tenuipalpus mansonculus* Ghai & Shenhmar (female). A, leg III; B, leg IV.

Acknowledgements

We thank Dr Qing-Hai Fan (Ministry for Primary Industries, Auckland, New Zealand) for help with references. While this paper was prepared, Zhi-Qiang Zhang was supported by Core funding for Crown Research Institutes from the Ministry of Business, Innovation and Employment's Science and Innovation Group.

References

- Collyer, E. (1973) Two new species of the genus *Colopalpus* (Acarina: Tenuipalpidae). *New Zealand Journal of Science*, 16, 529–532.
- De Leon, D. (1965) False spider mites of Jamaica and the Dominican Republic (Acarina: Tenuipalpidae). *Annals of the Entomological Society of America*, 58(4), 517–523.
- Ghai, S. & Shenhmar, M. (1984) A review of the world fauna of Tenuipalpidae (Acarina: Tetranychoidae). *Oriental Insects*, 18, 99–172.
<http://dx.doi.org/10.1080/00305316.1984.10432200>
- Gutierrez, J. & Schicha, E. (1982). Two new species of *Tenuipalpus* Donnadieu from New South Wales (Acari: Tenuipalpidae). *Journal of the Australian Entomological Society*, 21, 137–141.
<http://dx.doi.org/10.1111/j.1440-6055.1982.tb01781.x>
- Halliday, R.B. (1998) *Mites of Australia: a checklist and bibliography*. CSIRO Publishing, Melbourne, Australia, 317 pp.
- Meyer, M.K.P. (1979) The Tenuipalpidae (Acari) of Africa with keys to the world fauna. *Entomology Memoir, Department of Agriculture Republic South Africa, Pretoria*, 50, 1–133.
- Lindquist, E.E. (1985) External anatomy. In: Helle, W. & Sabelis, M.W. (Eds), *Spider Mites: Their Biology, Natural Enemies and Control*. Vol. 1a. Elsevier, Amsterdam, pp. 3–28.
- Smiley, R.L. & Gerson, U. (1995) A review of the Tenuipalpidae (Acari: Prostigmata) of Australia with descriptions of two new genera and four new species. *International Journal of Acarology*, 21(1), 33–45.
<http://dx.doi.org/10.1080/01647959508684041>
- Zhang, Z.-Q. & Fan, Q.-H (2004) Redescription of *Dolichotetranychus ancistrus* Baker & Pritchard (Acari: Tenuipalpidae) from New Zealand. *Systematic & Applied Acarology*, 9, 111–131.

Accepted by Qing-Hai Fan: 28 Aug. 2014; published 8 Sept. 2014