

## **Higher Ant Diversity in Native Vegetation Than in Stands of the Invasive Arundo, *Arundo donax* L., Along the Rio Grande Basin in Texas, USA**

Authors: Osbrink, Weste LA, Goolsby, John A, Thomas, Don B, Mejorado, Alba, Showler, Allan T, et al.

Source: International Journal of Insect Science, 9(1)

Published By: SAGE Publishing

URL: <https://doi.org/10.1177/1179543317724756>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

# Higher Ant Diversity in Native Vegetation Than in Stands of the Invasive *Arundo*, *Arundo donax* L., Along the Rio Grande Basin in Texas, USA

Weste LA Osbrink<sup>1</sup>, John A Goolsby<sup>2</sup>, Don B Thomas<sup>2</sup>, Alba Mejorado<sup>2</sup>, Allan T Showler<sup>1</sup> and Adalberto Pérez De León<sup>1</sup>

<sup>1</sup>Knipling-Bushland U.S. Livestock Insects Research Laboratory, USDA-ARS-SPA, Kerrville, TX, USA. <sup>2</sup>Cattle Fever Tick Research Lab, USDA-ARS, Edinburg, TX, USA.

International Journal of Insect Science  
Volume 9: 1–9  
© The Author(s) 2017  
Reprints and permissions:  
sagepub.co.uk/journalsPermissions.nav  
DOI: 10.1177/1179543317724756



**ABSTRACT:** Our hypothesis was that there will be greater ant biodiversity in heterogeneous native vegetation compared with *Arundo* stands. Changes in ant biodiversity due to *Arundo* invasion may be one of the ecological changes in the landscape that facilitates the invasion of cattle fever ticks from Mexico where they are endemic. Ants collected in pitfall traps were identified and compared between native vegetation and stands of *Arundo*, *Arundo donax* L., monthly for a year at 10 locations. A total of 82 752 ants representing 28 genera and 76 species were collected. More ants were collected in the native vegetation which also had greater species richness and biological diversity than ants collected from *Arundo* stands. It is suggested that the greater heterogeneous nature of native vegetation provided greater and more predictable nourishment in the form of nectars and more abundant arthropod prey when compared with *Arundo* stands.

**KEYWORDS:** Ant diversity, extrafloral nectaries, pitfall traps

**RECEIVED:** March 3, 2017. **ACCEPTED:** July 1, 2017.

**PEER REVIEW:** Five peer reviewers contributed to the peer review report. Reviewers' reports totaled 2484 words, excluding any confidential comments to the academic editor.

**TYPE:** Original Research

**FUNDING:** The author(s) received no financial support for the research, authorship, and/or publication of this article.

**DECLARATION OF CONFLICTING INTERESTS:** The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

**CORRESPONDING AUTHOR:** Weste LA Osbrink, Knipling-Bushland U.S. Livestock Insects Research Laboratory, USDA-ARS-SPA, 2700 Fredericksburg Road, Kerrville, TX 78028-9184, USA. Email: weste.osbrink@ars.usda.gov

## Introduction

Incidence of cattle fever ticks, *Rhipicephalus* (= *Boophilus*) spp, breaching the US quarantine zone along the Rio Grande is on the increase motivating studies of ecological changes occurring concurrent to the establishment of the invasive *Arundo donax* L. (Poales: Poaceae). *Arundo*, also known as giant reed or carrizo cane, is native to the Mediterranean coasts of Europe and North Africa to south Asia. *Arundo donax* is an invasive weed of riparian habitats of the southwestern United States.<sup>1–3</sup> Classified as an invasive perennial species, it spread widely in riparian zones of Texas where it has altered wildlife habitats, created fire hazards, compromised water conservation efforts, affected flood control, and reduced visibility for law enforcement officers along the international border with Mexico. *Arundo* might also facilitate cattle fever tick, *Rhipicephalus* (= *Boophilus*) spp, invasion into the permanent quarantine zone along the Rio Grande between Del Rio and Brownsville, TX, by harboring known mammalian host such as white-tailed deer, *Odocoileus virginianus* (Zimmermann).<sup>4–6</sup>

Ants were chosen as a survey taxon because they are known predators of ticks<sup>7–10</sup> and are often represented by multiple species demonstrating high diversity.<sup>11,12</sup> The diversity and abundance of ants in native vegetation and *Arundo* along the Rio Grande basin in Texas have not been previously studied. The objective of this study was to compare and contrast ant diversity between *Arundo* and native vegetation along the Rio Grande River at the Texas-Mexico border.

The hypothesis was that there will be greater ant biodiversity in heterogeneous native vegetation compared with *Arundo*

stands. Native vegetation containing a variety of plant species provides for greater occurrences of various edges and niches increasing opportunities for various ant species to have survival advantages over competitors including predator avoidance and opportunities for specialized relationships with myrmecophytic plants when contrasted with a monoculture.<sup>11</sup> Changes in ant biodiversity due to *Arundo* invasion may be one of the ecological changes in the landscape that facilitates the invasion of cattle fever ticks from Mexico where they are endemic.<sup>13</sup>

## Materials and Methods

### Study sites

Ant samples were collected from 10 Texas research sites located just north of the Rio Grande along the Texas-Mexico border (Table 1, Figure 1).

### Ant sampling

Ants were sampled using pitfall traps (Figure 2). A trap comprised 470-mL polypropylene container (Ball, Fishers, IN, USA) 10.5 cm × 9.9 cm × 7.0 cm (height × top diameter × bottom diameter). A plywood shelter (30.5 cm × 30.5 cm) was supported ≈1.3 cm above the trap. Each trap contained a 50:50 mixture of propylene glycol and water. Trapping intervals were ≈30 d. Four traps were located 10 m apart within an *Arundo* stand, and 4 traps were similarly placed in native vegetation at each of the 10 locations.



Creative Commons Non Commercial CC BY-NC: This article is distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 License (<http://www.creativecommons.org/licenses/by-nc/4.0/>) which permits non-commercial use, reproduction and distribution of the work without further permission provided the original work is attributed as specified on the SAGE and Open Access pages (<https://us.sagepub.com/en-us/nam/open-access-at-sage>).

**Table 1.** Ant sample study site locations along the Rio Grande, TX, USA.

LOCATION	COUNTY	LATITUDE, LONGITUDE
Los Indios	Cameron	26.05, -97.74
North American Butterfly Association (NABA)	Hidalgo	26.180243, -98.364973
Bentsen-Rio Grande State Park	Hidalgo	26.1731300, -98.3825200
San Ygnacio	Zapata	27.048175, -99.430788
Laredo Community College (LCC)	Webb	27.5084, -99.5214
La Bota Ranch	Webb	27.6161258, -99.5569872
Comanche Ranch	Maverick	28.643901, -100.444024
Rosita Ranch	Maverick	28.643901, -100.444024
Sycamore Creek	Kinney	29.4410659, -100.1228475
Del Rio	Val Verde	29.3709, -100.8959



**Figure 1.** Study sites in the cattle fever tick quarantine zone along the Texas-Mexico border.

Monthly pitfall trap collections were conducted during January to December 2014 at each site. Ant identification was conducted with a stereoscopic microscope using taxonomic ant keys.<sup>14–19</sup> A survey of plants and the presence of extrafloral nectaries was conducted at each study site.

*Statistical analysis*

Diversity profile estimations were calculated using Spade (Species Prediction and Diversity Estimation).<sup>20</sup> Continuous diversity profile including species richness, Shannon entropy, Simpson index, and Chao2 as well as their effective numbers of species based on incidence data were calculated using

species richness prediction and diversity estimation. Shannon entropy provides expected mean which increases with a species richness and evenness. Simpson index values increase as diversity decreases, whereas the inverse Simpson index values increase as diversity increases providing a dominance index giving more weight to common or dominant species. Chao2 estimates true species diversity of a sample using incidence data. Value differences were considered significant when their confidence intervals did not overlap. The program iNEXT<sup>21–23</sup> was used to plot sample size–based rarefaction and extrapolation sampling curves where this curve plots the species richness estimates for a rarefied and extrapolated sample with respect to sample size following bootstrapping 100 times



**Figure 2.** Standard pitfall trap.

(Figure 4).<sup>22</sup> iNEXT was used to calculate coverage-based rarefaction and extrapolation where species richness estimates for rarefied and extrapolated samples for sample coverage.<sup>21</sup> Using incidence data, iNEXT was used to calculate sample completeness curve where this curve plots the sample completeness (as measured by sample coverage) with respect to sample size. The social nature of ants promotes clumping as members of a colony will recruit numerous nest mates to a particular location making analysis of incidence data (presence or absence) most appropriate.<sup>24,25</sup>

Calculated Hill number unifies the reported biological diversity parameter such that the Hill numbers 0, near 1, and 2 provide the communities' species richness, exponent of Shannon index, and inverse of Simpson index, respectively. More weight is given to dominant species as the Hill number increases.<sup>23</sup> Coverage-based rarefaction and extrapolation sampling curve provide species richness estimates for rarefied sample and extrapolated sample with sample coverage up to double the reference sample size (Figure 4).

## Results

After 12 months, a total of 82 752 ants representing 28 genera and 76 species were captured (Table 2). Only 7 species were captured from all 10 locations (see Table 2). Twenty species were captured at single locations (Table 2). Seven of the ant species were found only in the southern most test sites, Los Indios, North American Butterfly Association (NABA), or Bentsen R.G. In addition, 11 species were found only in the northernmost test sites, Comanche Ranch, Rosita Ranch, Sycamore Creek, or Del Rio. *Pheidole* was the most diverse genus represented by 21 species (Table 2).

Excluding the NABA site, 2-fold to 15.5-fold greater abundances of ants were captured, and more species were encountered, in the native vegetation than in *Arundo* at each test site (Table 3). Ant dominance (number of traps with ants  $\times$  number of species captured at the respective locations) was higher in

native vegetation (Table 3). With the exception of Bentsen R.G., all calculated estimates of biodiversity were greater in native vegetation than in *Arundo* (Table 4). Chao2 estimated that species richness was greater in at least 70% of the test sites. Diversity indicated by the exponential of Shannon index was greater in native vegetation compared with *Arundo* at 40% of the sites: Laredo Community College, Comanche Ranch, Sycamore Creek, and Del Rio (Table 4). Biodiversity as quantified using the inverse of the Simpson index was also greater in native vegetation than in *Arundo* at 40% of the sites. The number of estimated species shared between native vegetation and *Arundo* at each study site ranged from 40.9% to 23.7% (Table 3).

Total ants collected (mean  $\pm$  SEM) were greater in native vegetation ( $6793.3 \pm 1593.8$ ) compared with *Arundo* ( $1421.2 \pm 374.2$ ;  $F = 10.768$ ;  $df = 1, 19$ ;  $P = .004$ ). Ant dominance (mean  $\pm$  SEM) was greater in native vegetation ( $216.0 \pm 18.1$ ) compared with *Arundo* ( $113.9 \pm 11.6$ ;  $F = 22.476$ ;  $df = 1, 19$ ;  $P < .001$ ). Total species collected (mean  $\pm$  SEM) was greater in native vegetation ( $27.4 \pm 1.9$ ) compared with *Arundo* ( $19.2 \pm 1.4$ ;  $F = 11.913$ ;  $df = 1, 19$ ;  $P = .003$ ). Native vegetation has greater diversity than the *Arundo* with both the empirical and extrapolated calculations (Table 4, Figure 3). Greater species richness is found in natural vegetation than is found in *Arundo* indicated by nonoverlap of confidence limits (Table 4, Figure 4). More than 76% of the native vegetation possessed extrafloral nectaries (Table 5).

## Discussion

Pitfall trapping of ants has been recognized as an effective monitoring technique.<sup>26,27</sup> The magnitude and diversity of ant assemblages reflect the responses of individual species to environmental conditions of native vegetation and *Arundo* stands. *Arundo* stands represent a new environmental setting different from native vegetation where food resources for ants differ. Thus, species distribution appears driven by resources resulting in the observed ant species assemblages.<sup>28</sup>

In the case of Los Indios, for example, the native vegetation has greater diversity than the *Arundo* with both the empirical and extrapolated calculations (Figure 3). Greater species richness in Los Indios natural vegetation is than found in *Arundo* indicated by nonoverlap of confidence limits (Figure 4). For Los Indios, the sample completeness curve indicates that the number of samples provided is adequate coverage of study area as indicated by the plateauing of the curve (Figure 4). The completeness curve provides a bridge between sample size-based and coverage-based rarefaction and extrapolation. Similar comparisons were conducted for all study sites as summarized in Table 4.

Presence of renewable and predictable food sources on vegetation will support greater populations of ants.<sup>29–31</sup> Composition of ant fauna differs between heterogeneous native vegetation and *Arundo* stands with the former supplying a heterogeneous supply of extrafloral nectaries and a greater

**Table 2.** Presence/absence of ants captured in pitfall traps in tick quarantine zone.

SPECIES	SUBFAMILY	LI	NABA	BRG	SY	LCC	LB	CR	RR	SC	DR
1. <i>Aphaenogaster texana texana</i>	Myrmicinae	+	+	0	0	0	0	0	0	0	0
2. <i>Atta texana</i>	Myrmicinae	0	0	+	+	0	0	0	+	+	0
3. <i>Brachymyrmex depilis</i>	Formicinae	0	0	0	0	+	+	0	0	0	+
4. <i>Brachymyrmex patagonicus</i>	Formicinae	+	+	+	+	+	+	+	+	+	+
5. <i>Camponotus festinatus</i>	Formicinae	0	+	0	0	0	+	+	0	+	0
6. <i>Camponotus floridanus</i>	Formicinae	+	+	+	+	0	0	0	0	0	0
7. <i>Camponotus planatus</i>	Formicinae	+	+	+	0	0	0	0	0	0	0
8. <i>Camponotus sayi</i>	Formicinae	+	+	0	0	0	0	0	0	+	0
9. <i>Camponotus texanus</i>	Formicinae	+	+	+	+	+	0	0	0	+	0
10. <i>Cardiocondyla emeryi</i>	Myrmicinae	0	0	0	0	+	0	0	0	0	0
11. <i>Crematogaster crinosa</i>	Myrmicinae	0	+	0	0	0	0	0	0	0	0
12. <i>Crematogaster laeviuscula</i>	Myrmicinae	+	+	+	+	+	+	+	+	+	+
13. <i>Crematogaster lineolata</i>	Myrmicinae	0	+	0	0	0	0	0	0	0	0
14. <i>Crematogaster torosa</i>	Myrmicinae	+	+	0	0	0	0	0	0	0	0
15. <i>Cyphomyrmex rimosus</i>	Myrmicinae	+	+	+	+	+	+	+	0	+	0
16. <i>Dorymyrmex bicolor</i>	Dolichoderinae	0	0	0	0	0	+	0	0	0	+
17. <i>Dorymyrmex flavus</i>	Dolichoderinae	+	+	+	0	+	+	+	+	+	+
18. <i>Dorymyrmex insanus</i>	Dolichoderinae	0	0	0	0	+	+	0	+	+	+
19. <i>Forelius mccooki</i>	Dolichoderinae	+	+	+	+	+	+	+	+	+	+
20. <i>Forelius pruinosus</i>	Dolichoderinae	+	+	0	0	+	+	0	+	+	+
21. <i>Formica pallidefulva</i>	Formicinae	0	0	0	0	0	+	0	0	+	0
22. <i>Hypoponera opaciceps</i>	Ponerinae	0	0	0	0	0	0	+	0	0	0
23. <i>Hypoponera opacior</i>	Ponerinae	0	+	0	0	0	+	+	+	+	0
24. <i>Hypoponera punctatissima</i>	Ponerinae	0	0	0	+	0	0	0	+	0	0
25. <i>Labidus coecus</i>	Dorylinae	+	+	+	+	+	0	0	+	+	+
26. <i>Leptogenys elongata</i>	Ponerinae	+	+	+	0	0	0	+	+	+	+
27. <i>Monomorium minimum</i>	Myrmicinae	+	0	0	+	0	0	+	+	+	0
28. <i>Monomorium pharoanis</i>	Myrmicinae	0	0	0	0	0	0	+	0	0	0
29. —	Mutillidae (Family)	+	+	+	+	+	+	+	+	+	+
30. <i>Myrmecocystus mendax</i>	Formicinae	0	0	0	0	0	0	0	0	+	0
31. <i>Neivamyrmex nigrescens</i>	Dorylinae	0	+	+	0	0	0	0	+	0	0
32. <i>Neivamyrmex swainsonii</i>	Dorylinae	0	+	+	+	0	+	+	+	0	+
33. <i>Neivamyrmex texanus</i>	Dorylinae	0	0	0	0	0	0	0	+	0	0
34. <i>Odontomachus clarus</i>	Ponerinae	0	0	0	0	0	0	0	+	+	+
35. <i>Pachycondyla harpax</i>	Ponerinae	0	0	0	+	0	+	0	+	0	0
36. <i>Paratrechina terricola</i>	Formicinae	+	+	+	0	+	+	+	+	+	+
37. <i>Pheidole bicarinata</i>	Myrmicinae	+	+	+	0	+	+	+	+	+	+
38. <i>Pheidole cockerelli</i>	Myrmicinae	0	0	+	0	+	0	0	0	0	0
39. <i>Pheidole dentata</i>	Myrmicinae	+	+	+	+	+	+	+	+	+	+

Table 2. (Continued)

SPECIES	SUBFAMILY	LI	NABA	BRG	SY	LCC	LB	CR	RR	SC	DR
40. <i>Pheidole flavens</i>	Myrmicinae	0	0	0	0	0	0	0	0	+	0
41. <i>Pheidole floridana constipata</i>	Myrmicinae	+	+	+	+	+	+	+	+	+	+
42. <i>Pheidole floridana floridana</i>	Myrmicinae	+	+	0	0	+	+	0	0	+	+
43. <i>Pheidole humeralis</i>	Myrmicinae	0	0	0	0	0	+	0	0	0	0
44. <i>Pheidole hyatti</i>	Myrmicinae	0	0	0	0	0	0	0	+	+	0
45. <i>Pheidole lamia</i>	Myrmicinae	0	0	0	0	0	0	0	+	+	0
46. <i>Pheidole mera</i>	Myrmicinae	+	0	0	0	0	+	0	+	+	0
47. <i>Pheidole metallescens</i>	Myrmicinae	0	0	+	0	0	+	0	+	+	0
48. <i>Pheidole moerens</i>	Myrmicinae	0	0	0	0	0	0	0	0	+	0
49. <i>Pheidole nukuliceps</i>	Myrmicinae	0	0	0	0	0	+	0	0	0	0
50. <i>Pheidole pelor</i>	Myrmicinae	0	0	0	0	0	0	0	0	+	0
51. <i>Pheidole porcula</i>	Myrmicinae	+	0	+	+	+	+	0	+	+	+
52. <i>Pheidole sciara</i>	Myrmicinae	0	0	0	+	+	0	+	+	+	0
53. <i>Pheidole spadonia</i>	Myrmicinae	+	0	0	0	0	0	0	0	0	0
54. <i>Pheidole tetra</i>	Myrmicinae	0	0	+	0	+	+	+	0	+	+
55. <i>Pheidole texana</i>	Myrmicinae	0	0	0	0	0	0	0	+	0	0
56. <i>Pheidole tysoni</i>	Myrmicinae	0	0	0	0	0	0	0	0	0	+
57. <i>Pheidole vallicola</i>	Myrmicinae	0	0	0	0	0	0	+	+	0	0
58. <i>Pogonomyrmex barbatus</i>	Myrmicinae	+	0	0	0	0	+	0	0	0	+
59. <i>Pogonomyrmex rugosus</i>	Myrmicinae	0	0	0	0	0	+	0	0	0	0
60. <i>Pseudomyrmex gracilis</i>	Pseudomyrmecinae	0	+	+	0	0	0	0	0	0	0
61. <i>Pseudomyrmex pallidus</i>	Pseudomyrmecinae	0	+	0	0	0	0	0	0	0	0
62. <i>Solenopsis aurea</i>	Myrmicinae	0	+	0	0	0	0	0	+	0	0
63. <i>Solenopsis geminata</i>	Myrmicinae	0	0	+	0	+	+	+	+	0	0
64. <i>Solenopsis invicta</i>	Myrmicinae	+	+	+	+	+	+	+	+	+	+
65. <i>Solenopsis molesta</i>	Myrmicinae	+	+	0	+	0	0	0	0	+	0
66. <i>Solenopsis texana</i>	Myrmicinae	+	+	+	+	+	+	+	+	+	+
67. <i>Strumigenys boneti</i>	Myrmicinae	+	0	0	0	0	0	0	0	0	0
68. <i>Strumigenys louisianae</i>	Myrmicinae	0	+	0	0	+	+	0	+	+	+
69. <i>Strumigenys membranifera</i>	Myrmicinae	0	0	0	0	+	0	0	0	0	0
70. <i>Tapinoma litorale</i>	Dolichoderinae	0	0	+	0	0	0	0	0	0	0
71. <i>Temnothorax subditivus</i>	Myrmicinae	+	+	+	+	0	0	+	+	+	+
72. <i>Temnothorax pergandei</i>	Myrmicinae	+	0	0	0	+	0	+	+	+	0
73. <i>Tetramorium bicarinatum</i>	Myrmicinae	0	+	0	0	0	+	0	0	0	0
74. <i>Tetramorium caldarium</i>	Myrmicinae	0	0	0	0	0	+	+	0	0	0
75. <i>Tetramorium lanuginosum</i>	Myrmicinae	0	0	+	0	+	0	+	0	0	0
76. <i>Tetramorium spinosum</i>	Myrmicinae	+	0	+	+	+	+	+	0	+	0
77. <i>Trachymyrmex turrifex</i>	Myrmicinae	0	+	0	0	+	+	+	+	+	+

Abbreviations: BRG, Bentsen R.G.; CR, Comanche Ranch; DR, Del Rio; LB, La Bota; LCC, Laredo Community College; LI, Los Indios; NABA, North American Butterfly Association; RR, Rosita Ranch; SC, Sycamore Creek; SY, San Ygnacio.

**Table 3.** Enumeration of ants captured in pitfall traps at 10 sample sites along the Rio Grande in Texas January to December 2014.

	LI	NABA	BRG	SY	LCC	LB	CR	RR	SC	DR
<b>Arundo</b>										
Total	573	1045	377	214	2059	1063	1089	4310	1737	1745
Ant dominance	137	168	83	62	150	117	84	154	103	81
No. of species	23	25	23	14	18	18	15	23	21	12
<b>Native vegetation</b>										
Total	1607	1099	738	3317	11461	5149	14467	8583	11884	9628
Ant dominance	248	276	111	173	238	149	257	224	287	197
No. of species	29	30	24	20	26	28	23	33	40	21


Abbreviations: BRG, Bentsen R.G.; CR, Comanche Ranch; DR, Del Rio; LB, La Bota; LCC, Laredo Community College; LI, Los Indios; NABA, North American Butterfly Association; RR, Rosita Ranch; SC, Sycamore Creek; SY, San Ygnacio.  
Ant dominance = traps with ants x species.

**Table 4.** Ant biodiversity in *Arundo* stands and mixed native vegetation ( $\pm$ SE).


LOCATION	NO. OF OBSERVED SPECIES	CHAO2 ESTIMATED SPECIES RICHNESS	EXPONENTIAL OF SHANNON INDEX	INVERSE OF SIMPSON INDEX	CHAO2 ESTIMATED SHARED SPECIES (OBSERVED)
<b>Los Indios</b>					19.3 $\pm$ 0.9 (19)
<i>Arundo</i>	23	24.6 $\pm$ 2.1a	19.0 $\pm$ 1.5a	15.0 $\pm$ 1.7a	
Native	29	33.1 $\pm$ 4.8b	19.7 $\pm$ 1.1a	14.8 $\pm$ 0.9a	
<b>NABA</b>					28.3 $\pm$ 12.0 (19)
<i>Arundo</i>	25	29.4 $\pm$ 4.7a	14.8 $\pm$ 1.3a	9.5 $\pm$ 0.7a	
Native	30	46.3 $\pm$ 14.5a	14.9 $\pm$ 1.1a	9.4 $\pm$ 0.5a	
<b>Bentsen R.G.</b>					24.9 $\pm$ 8.1 (18)
<i>Arundo</i>	24	73.6 $\pm$ 59.1a	14.1 $\pm$ 2.2a	6.4 $\pm$ 1.1a	
Native	24	27.5 $\pm$ 3.8a	19.2 $\pm$ 1.4a	15.1 $\pm$ 1.4b	
<b>San Ygnacio</b>					12.0 $\pm$ 2.5 (11)
<i>Arundo</i>	14	16.2 $\pm$ 3.3a	11.4 $\pm$ 1.5a	8.1 $\pm$ 1.3a	
Native	20	28.8 $\pm$ 10.0b	13.1 $\pm$ 0.9a	10.4 $\pm$ 0.6a	
<b>LCC</b>					14.5 $\pm$ 1.4 (14)
<i>Arundo</i>	18	20.6 $\pm$ 3.4a	11.1 $\pm$ 0.9a	8.2 $\pm$ 0.6a	
Native	26	32.0 $\pm$ 6.0b	14.6 $\pm$ 0.8b	10.0 $\pm$ 0.5a	
<b>La Bota</b>					18.3 $\pm$ 7.8 (11)
<i>Arundo</i>	20	51.8 $\pm$ 39.4a	10.7 $\pm$ 1.2a	6.6 $\pm$ 0.7a	
Native	22	34.2 $\pm$ 10.5a	11.5 $\pm$ 1.0a	7.2 $\pm$ 0.6a	
<b>Comanche Ranch</b>					10.5 $\pm$ 2.5 (9)
<i>Arundo</i>	15	24.5 $\pm$ 2.2b	8.3 $\pm$ 1.5a	4.7 $\pm$ 0.5a	
Native	23	28.7 $\pm$ 1.1a	13.2 $\pm$ 0.6b	9.6 $\pm$ 0.5b	
<b>Rosita Ranch</b>					27.3 $\pm$ 10.7 (18)
<i>Arundo</i>	31	54.5 $\pm$ 19.8a	17.9 $\pm$ 1.3a	11.2 $\pm$ 0.8a	
Native	33	44.8 $\pm$ 9.5b	19.1 $\pm$ 1.5a	12.4 $\pm$ 0.8a	
<b>Sycamore Creek</b>					32.4 $\pm$ 11.6 (19)
<i>Arundo</i>	23	36.2 $\pm$ 12.2a	14.4 $\pm$ 1.5a	8.2 $\pm$ 0.9a	
Native	40	47.4 $\pm$ 5.85b	22.2 $\pm$ 1.2b	13.9 $\pm$ 0.8b	
<b>Del Rio</b>					15.3 $\pm$ 11.8 (8)
<i>Arundo</i>	12	19.3 $\pm$ 8.0b	6.2 $\pm$ 1.1a	3.4 $\pm$ 0.4a	
Native	21	26.1 $\pm$ 5.3a	11.7 $\pm$ 0.7b	8.0 $\pm$ 0.6b	

Abbreviations: LCC, Laredo Community College; NABA, North American Butterfly Association.

Chao2 estimation of species richness is equivalent of diversity of order 0; exponential of Shannon index is equivalent of diversity of order 1; inverse of Simpson index is equivalent of diversity of order 2 (Chao et al., 2015). <sup>a,b</sup>Mean values in a column from the same location pair followed by the same letter are not significantly different as indicated by overlap of 95% confidence intervals.



**Figure 3.** Los Indios ant diversity index Hill number plots, native vegetation (left), and *Arundo* (right) (Chao et al., 2015).



**Figure 4.** Los Indios ant species richness and diversity comparing *Arundo* and natural habitat. Chao2 (Chao and Jost, 2012) used as estimator of species richness and suggested estimator of sample coverage. (A) Sample size–based rarefaction and extrapolation sampling curve: species richness estimates for a rarefied and extrapolated sample with sample size up to double the reference sample size. (B) Sample completeness curve: sample completeness (as measured by sample coverage) with respect to sample size. This curve provides a bridge between sample size–based and coverage–based rarefaction and extrapolation. (C) Coverage–based rarefaction and extrapolation sampling curve: species richness estimates for rarefied sample and extrapolated sample with sample coverage up to double the reference sample size.

**Table 5.** Presence/absence of plants and EFNs at 10 sample sites along the Rio Grande in Texas January to December 2014.

SPECIES	EFN	COMMON NAME	LI	NABA	BRG	SY	LCC	LB	CR	RR	SC	DR
<i>Prosopis glandulosa</i>	+	Mesquite	+	0	0	+	+	+	+	+	+	+
<i>Celtis laevigata</i>	0	Hackberry	0	+	+	0	0	+	+	+	+	0
<i>Ulmus crassifolia</i>	+	Cedar elm	+	+	+	+	+	+	+	+	+	+
<i>Acacia</i> spp	+	Acacia	0	+	+	0	0	+	+	0	+	0
<i>Acacia rigidula</i>	+	Blackbrush	+	+	+	+	+	+	+	+	+	+
<i>Celtis pallida</i>	+	spiny hackberry	+	0	+	+	0	+	+	+	0	0
<i>Acacia smallii</i>	+	Huisache	+	+	+	+	+	+	+	+	+	+
<i>Mimosa pigra</i>	0	Mimosa	0	0	+	+	0	+	+	+	+	0
<i>Opuntia lindheimeri</i>	+	Prickly pear cactus	+	+	+	+	+	+	+	+	+	+
<i>Triadica sebifera</i>	+	Chinese tallow	+	+	+	0	+	+	+	0	+	+
<i>Ricinus communis</i>	0	Castor bean		0	+	+	0	0	+	0	+	0
<i>Cenchrus ciliaris</i>	+	Buffel grass		+	+	+	+	+	+	+	+	+
<i>Panicum maximum</i>	+	Guinea grass		+	+	+	+	+	+	+	+	+
<i>Triadica sebifera</i>	+	Chinese tallow	+	+	+	0	+	+	+	0	+	+
<i>Ricinus communis</i>	0	Castor bean	0	+	+	0	0	+	0	+	0	0
<i>Cenchrus ciliaris</i>	+	Buffel grass	+	+	+	+	+	+	+	+	+	+
<i>Panicum maximum</i>	+	Guinea grass	+	+	+	+	+	+	+	+	+	+

Abbreviations: BRG, Bentsen R.G.; CR, Comanche Ranch; DR, Del Rio; EFNs, extrafloral nectaries; LB, La Bota; LCC, Laredo Community College; LI, Los Indios; NABA, North American Butterfly Association; RR, Rosita Ranch; SC, Sycamore Creek; SY, San Ygnacio.

reservoir of arthropods.<sup>31–33</sup> Many species of the native flora have extrafloral nectaries including trees and shrubs (Table 5). Reducing the dominance of *Arundo* and the subsequent re-establishment of native vegetation could increase the diversity and abundance of ant communities within the cattle fever tick permanent quarantine zone.

Extrafloral nectar is important for ant survival and growth and vitality of ant colonies. Ant richness and abundance will be higher where the availability of resources is greatest from water to extrafloral nectar to flower nectar being particularly important during dry periods.<sup>34–36</sup> Ants are obligated to search for other alternate sources of food and water including arthropod plant pests as prey where acquisition of resources by generalists with ant presence is influenced by availability of resources.<sup>34,35</sup>

*Arundo* stands provide a simpler environment with less resources when compared with native vegetation which results in repressed ant communities with less biodiversity.

## Acknowledgements

The authors thank reviewers for meaningful improvements of this manuscript.

## Author Contributions

All authors contributed equally.

## REFERENCES

- DiTomaso JM, Healy EA. *Aquatic and Riparian Weeds of the West* (ANR Publication 3241). Berkeley, CA: University of California Press; 2003.
- Yang C, Goolsby JA, Everitt JH. Using QuickBird satellite imagery to estimate giant reed infestations in the Rio Grande Basin of Mexico. *J Appl Remote Sens*. 2009;3:033530.
- Yang C, Everitt JH, Goolsby JA. Using aerial photography for mapping giant reed infestations along the Texas-Mexico portion of the Rio Grande. *Invas Plant Sci Mana*. 2011;4:402–410.
- Goolsby JA, Moran PJ, Adamczyk JA, et al. Host range of the European, rhizome-stem feeding scale *Rhizaspidiotus donacis* (Hemiptera: Diaspididae), a candidate biological control agent for giant reed, *Arundo donax* (Poales: Poaceae) in North America. *Biocontrol Sci Techn*. 2010;19:899–918.
- Moran PJ, Goolsby JA. Biology of the armored scale *Rhizaspidiotus donacis* (Hemiptera: Diaspididae), a candidate agent for biological control of giant reed. *Ann Entomol Soc Am*. 2010;103:252–263.
- Seawright EK, Rister M, Laceywell R, et al. Economic implications for the biological control of *Arundo donax* in the Rio Grande Basin. *Southwest Entomol*. 2010;34:377–394.
- Butler JF, Camino ML, Perez TO. *Boophilus microplus* and the fire ant *Solenopsis geninata*. In: Rodriguez JG (ed.) *Recent Advances in Acarology*. New York: Academic Press; 1979: 469–475.
- Robertson H. Egg predation by ants as a partial explanation of the differences in performance of *Cactoblastis cactorum* on cactus weeds in South Africa and Australia. In: Proceedings of the 6th International Symposium Biological Control of Weeds (ed ES Delfosse); August 19–25, 1984: 83–88; Vancouver, BC, Canada; 1985.
- Showler AT, Reagan TE. Ecological interactions of the red imported fire ant in the southeastern United States. *J Entomol Sci* 1987;1:52–64.
- Velez-Bonner A, Osbrink WLA, Summy KR, et al. Mitigating predatory ants promotes establishment of biological control of *Arundo* by *Arundo* scale in the cattle fever tick quarantine zone. *Subtropical Plant Sci*. 2013;65:38–44.
- Longino JT, Colwell RK. Density compensation, species composition, and richness of ants on a neotropical elevational gradient. *Ecosphere*. 2011;2:1–20.

12. Longino JT, Coddington J, Colwell RK. The ant fauna of a tropical rain forest: estimating species richness three different ways. *Ecol.* 2002;83:289–702.
13. Esteve-Gassent MD, Pérez de León AA, Romero-Salas D, et al. Pathogenic landscape of transboundary zoonotic diseases in the Mexico-US border along the Rio Grande. *Front Public Health.* 2014;2:1–23.
14. Bolton B. *Identification Guide to the Ant Genera of the World.* Cambridge, MA: Harvard University Press; 1994.
15. Cook JL, O'Keefe S, Vinson SB, Drees BM. *Texas Pest Ant Identification: An Illustrated Key to Common Pest Ants and Fire Ant Species.* College Station, TX: Texas A&M AgriLife Extension Service, The Texas A&M University System; 2016.
16. Creighton W. The ants of North America. *Bull Mus Comp Zool.* 1950;104:1–585 + 57 plates.
17. Holldobler B, Wilson EO. *The Ants.* Cambridge, MA: Belknap Press of Harvard University Press; 1990: 732.
18. Mueller UG. Mueller Lab's "ant identification key," University of Texas at Austin. <http://www.sbs.utexas.edu/muelleru/AntOutreach/>. Published 2016.
19. Vinson SB, O'Keefe S, Cook J. *The Common Ant Genera of Texas: B-6138.* College Station, TX: Texas A&M AgriLife Extension Service and Texas A&M AgriLife Research, The Texas A&M University System; 2003: 44. [http://fireant.tamu.edu/files/2014/03/ENTO\\_001.pdf](http://fireant.tamu.edu/files/2014/03/ENTO_001.pdf).
20. Chao A, Ma KH, Hsieh TC, Chiu CH. Online Program SpadeR (Species-richness Prediction and Diversity Estimation in R): Program and User's Guide. [http://chao.stat.nthu.edu.tw/wordpress/software\\_download/](http://chao.stat.nthu.edu.tw/wordpress/software_download/). Published 2015.
21. Chao A, Jost L. Coverage-based rarefaction and extrapolation: standardizing samples by completeness rather than size. *Ecology.* 2012;93:2533–2547.
22. Colwell RK, Chao A, Gotelli NJ, et al. Models and estimators linking individual-based and sample-based rarefaction, extrapolation and comparison of assemblages. *J Plant Ecol.* 2012;5:3–21.
23. Chao A, Gotelli NJ, Hsieh TC, et al. Rarefaction and extrapolation with Hill numbers: a unified framework for sampling and estimation in biodiversity studies. *Ecol Monogr.* 2014;84:45–67.
24. King J, Porter S. Evaluation of sampling methods and species richness estimators for ants in upland ecosystems in Florida. *Environ Entomol.* 2005;34: 1566–1578.
25. Morrison L, Porter S. Positive association between densities of the red imported fire ant, *Solenopsis invicta* (Hymenoptera: Formicidae), and generalized and arthropod diversity. *Environ Entomol.* 2003;32:548–554.
26. Borgelt A, New T. Pitfall trapping for ants (Hymenoptera, Formicidae) in mesic Australia: the influence of trap diameter. *J Insect Conserv.* 2005;9:219–221.
27. Pendola A, New T. Depth of pitfall traps—does it affects interpretation of ant (Hymenoptera, Formicidae) assemblages? *J Insect Conserv.* 2007;11:199–201.
28. Beier P, Albuquerque F. Evaluating  $\beta$  diversity as a surrogate for species representation at fine scale. *PLoS ONE.* 2016;11:e0151048.
29. Showler AT, Knaus RM, Reagan TE. Foraging territoriality of the imported fire ant, *Solenopsis invicta* Buren, in sugarcane as determined by neutron activation analysis. *Insect Soc.* 1989;36:235–239.
30. Showler AT, Reagan TE. Effects of sugarcane borer, weed, and nematode control strategies in Louisiana sugarcane. *Environ Entomol.* 1991;20:358–370.
31. Belchior C, Sendoya SF, del-Claro K. Temporal variation in the abundance and richness of foliage-dwelling ants mediated by extrafloral nectar. *PLoS ONE* 2016;11:e0158283.
32. Lach L, Hobbs ER, Majer E. Herbivory-induced extrafloral nectar increases native and invasive ant worker survival. *Popul Ecol.* 2009;51:237–243.
33. Byk J, Del-Claro K. Ant-plant interaction in the Neotropical savanna: direct beneficial effects of extrafloral nectar on ant colony fitness. *Popul Ecol.* 2011;53:327–332.
34. Ness JH, Morris WF, Bronstein JL. For ant protected plants, the best defense is a hungry offense. *Ecol.* 2009;90:2823–2831.
35. Ruffner GA, Clark WD. Extrafloral nectar of *Ferocactus acanthodes* (Cactaceae): composition and its importance to ants. *Am J Bot.* 1986;73:185–189.
36. Ness JH, Morris WF, Bronstein JL. For ant protected plants, the best defense is a hungry offense. *Ecol.* 2009;90:2823–2831.