

Comparison of noninvasive genetics and camera trapping for estimating population density of ocelots (*Leopardus pardalis*) on Barro Colorado Island, Panama

Authors: Rodgers, Torrey W., Giacalone, Jacalyn, Heske, Edward J., Janečka, Jan E., Phillips, Christopher A., et al.

Source: Tropical Conservation Science, 7(4) : 690-705

Published By: SAGE Publishing

URL: <https://doi.org/10.1177/194008291400700408>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Research Article

Comparison of noninvasive genetics and camera trapping for estimating population density of ocelots (*Leopardus pardalis*) on Barro Colorado Island, Panama

Torrey W. Rodgers^{1,3}, Jacalyn Giacalone^{2,3}, Edward J. Heske^{1,4}, Jan E. Janečka⁵, Christopher A. Phillips⁴ and Robert L. Schooley⁶

¹ Department of Animal Biology, University of Illinois, Urbana, IL, USA

² Montclair State University, Montclair, NJ, USA

³ Smithsonian Tropical Research Institute, Ancon, Panama

⁴ Illinois Natural History Survey, Prairie Research Institute, University of Illinois, Champaign, IL, USA

⁵ Department of Biological Sciences, Duquesne University, Pittsburgh, PA, USA

⁶ Department of Natural Resources and Environmental Sciences, University of Illinois, Urbana, IL, USA

Corresponding author: Torrey W. Rodgers, Department of Animal Biology, University of Illinois, 505 S. Goodwin Ave., Urbana, IL, USA. Email: Torrey.w.rodgers@gmail.com

Abstract

Estimates of population density are essential for the effective conservation and management of any threatened species. Accurately estimating density of elusive carnivores can be a challenge, however. One approach to this challenge is integration of DNA collected noninvasively from feces with capture-recapture modeling. To date, the bias and precision of this technique have seldom been evaluated in the field. We compared density estimates of ocelots (*Leopardus pardalis*) derived from fecal noninvasive genetic techniques to density estimates from camera trapping in the same population, during the same study period. Density estimates from the two techniques were comparable, especially when using spatially explicit capture-recapture models. Population density estimated using the program *DENSITY* was 1.74/km² (SE = 0.584) from noninvasive genetics and 1.59/km² (SE = 0.464) from camera trapping. These estimates also represent the highest reported ocelot population density within the species range.

Keywords: Elusive species conservation, ocelot, spatially-explicit capture-recapture.

Received: 19 August 2014; Accepted 23 October 2014; Published: 15 December 2014

Copyright: © Torrey W. Rodgers, Jacalyn Giacalone, Edward J. Heske, Jan E. Janečka, Christopher A. Phillips and Robert L. Schooley. This is an open access paper. We use the Creative Commons Attribution 4.0 license <http://creativecommons.org/licenses/by/3.0/us/>. The license permits any user to download, print out, extract, archive, and distribute the article, so long as appropriate credit is given to the authors and source of the work. The license ensures that the published article will be as widely available as possible and that your article can be included in any scientific archive. Open Access authors retain the copyrights of their papers. Open access is a property of individual works, not necessarily journals or publishers.

Cite this paper as: Rodgers, T. W., Giacalone, J., Heske, E. J., Janečka, J. E., Phillips, C. A. and Schooley, R. L. 2014. Comparison of noninvasive genetics and camera trapping for estimating population density of ocelots (*Leopardus pardalis*) on Barro Colorado Island, Panama. *Tropical Conservation Science* Vol.7 (4): 690-705. Available online: www.tropicalconservationscience.org

Introduction

Population density is a parameter of vital importance for species conservation. For rare or elusive species, however, acquiring reliable density estimates based on visual observation or capture can be difficult or impossible [1]. As a result, two noninvasive approaches are now commonly used to estimate abundance or density of elusive carnivores in conjunction with capture-recapture (CR) models: camera trapping [2, 3] and noninvasive genetics [4, 5]. To obtain CR density estimates from camera trapping, individual animals are identified from photographs based on morphological differences such as unique pelage patterns [3, 6], allowing researchers to construct capture histories for each individual. For the many elusive species without unique individual markings, however, one realistic option for estimating population density using CR methods is noninvasive genetics, in which individuals are identified from remotely collected DNA using genetic markers [7].

Since its first application by Karanth in 1995 [2], camera trapping has been widely used, reviewed, and evaluated. To date >100 published studies have used camera trapping in conjunction with CR techniques to estimate abundance or density, including >50 since 2010 [8]. Additionally, the accuracy of abundance and density estimates from camera trapping has been evaluated through comparisons with populations of known size [9], comparisons with estimates from telemetry [9-13], and via simulation [14, 15]. As a result camera trapping is considered a reliable method for estimating abundance and density in rare or elusive species [16, 17].

Use of noninvasive genetics to estimate abundance has recently become more common, but few studies have compared estimates derived from noninvasive genetics with estimates from other sources to evaluate the accuracy or potential bias of the technique [but see 18-21]. In addition, most noninvasive genetics studies have only estimated abundance and not population density [1, 19, 20]. Abundance by itself is a less valuable parameter for conservation, as it makes comparisons with other studies and other populations difficult [1]. Estimating abundance with CR models is relatively straightforward. Estimating density can be more difficult, however, because it requires accurate estimation of the effective survey area (ESA) [22], which is a challenge in geographically open populations and in areas with patchy habitat [23]. In the past, most camera trap and noninvasive genetics studies that estimated density have used derivations of the mean maximum distance moved (MMDM) between captures to estimate ESA, but the validity of this method has been debated [10, 11, 15, 24]. More recently, sophisticated maximum-likelihood and Bayesian spatially-explicit capture-recapture (SECR) models have been developed that estimate density directly by modeling ESA explicitly, using the geographic locations of captures [25-27].

To our knowledge, only one study [28] has directly compared estimates of population density from noninvasive genetics with estimates from camera trapping in the same population during the same time. In the previous study, however, the number of recaptures from noninvasive genetics was far below the minimum number required for the SECR analysis used [29], and thus was invalid. Our primary objective was to directly compare density estimates from noninvasive genetics with estimates from camera trapping, using sample sizes large enough for valid SECR analyses. As a secondary objective, we compared traditional CR models with a newer SECR model. We did this in a population of ocelots (*Leopardus pardalis*), a medium-sized felid, on Barro Colorado Island (BCI) in Panama. Ocelots are considered of least conservation concern in the core of their range, but many populations are threatened by habitat loss, and some northern populations are critically endangered [30]. Ocelots are a suitable organism for such a study because individuals can be identified for CR analyses based both on unique spot patterns [6] and unique DNA genotypes collected noninvasively from fecal samples. Our island study site also is a suitable setting

for such a study because it is relatively closed geographically, yet is large enough to maintain a stable breeding population. For our density estimates from camera trapping, we did not face the same problems with estimating ESA inherent in other studies, as we were simply able to use island area as our ESA. We think our camera-trapping results reflect a relatively unbiased estimate of population density, making them an excellent standard for comparing estimates from fecal noninvasive genetic techniques.

Methods

Study Area

Field work was conducted on BCI, a 15.43-km² island in the Panama Canal waterway, at a research station operated by the Smithsonian Tropical Research Institute (Fig. 1). BCI (9°10'N, 79°51'W) sits within Gatun Lake, an artificial body of water created in 1912 by the damming of the Chagres River to create the Panama Canal, and is part of the protected 54-km² Barro Colorado Nature Monument. Vegetation is tropical moist forest, and topography is dominated by hills that reach a maximum elevation of 165 m above sea level. Mean temperature is 27 °C, and average annual precipitation is 2,600 mm, with 90% of rainfall occurring from May through November [31]. Forest cover on the island is a mix of old growth forest and mature secondary forest >80 years old, all of which is suitable habitat for ocelots.




Fig. 1: Map of Barro Colorado Island in Panama, our study area for estimating population density of ocelots (*Leopardus pardalis*) from camera trapping and noninvasive genetics.

Scat surveys

Ocelots commonly defecate at latrine sites that are used by multiple individuals of both sexes [32, 33]. These latrines are typically located in large cavities or overhanging buttresses of large trees, underneath buttresses of fallen trees, or underneath human structures (Fig. 2). The primary objective of our surveys was to locate as many ocelot latrines as possible. All scats used in this study were found at such latrines. Seven latrines were located during a pilot study in 2011, five of which were still in use by ocelots in 2012. To find additional latrines in 2012, we walked all 39.5 km of trails on BCI (Fig. 1) a minimum of 3 times (once every 33 days). In addition, we walked >390 km of random, off-trail transects throughout the island in search of latrines. Once a latrine was located, all scats were removed, and it was revisited every 4-7 days for the remainder of the study to collect additional scats. Scat collection was conducted over 99 days from 29-January to 6-May 2012. The study length was chosen as a balance between being long enough to obtain sufficient data to estimate density, while still being short enough to satisfy assumptions of demographic closure [15]. Scats decompose quickly in the warm and tropical environment of Panama, so all collected scats were deposited by animals during the study period or shortly beforehand (within several days).



Fig. 2. An ocelot (*Leopardus pardalis*) defecating at a latrine under the buttress of a large tree on Barro Colorado Island, Panama.

Camera trapping

We conducted camera trapping using an array of 21 Reconyx PC900 and RC55 trail cameras (Reconyx Inc., Holmen, Wisconsin). Cameras were distributed evenly throughout the entire island so that gaps between cameras that could contain an ocelot home range were unlikely (Fig. 1), ensuring all individuals on the island had a capture probability > 0 [2]. The total grid size of camera traps (10.18 km^2) was several times larger than an ocelot home range (3.5 km^2 for males, 1.5 km^2 for females [34]) as suggested for unbiased density estimates [15, 35]. We placed all cameras along trails to maximize capture probability. Eighteen cameras were part of an ongoing, multi-year camera trap survey of the BCI mammal community; three additional cameras were placed to fill gaps. Each camera station consisted of one camera and could therefore photograph only one side of an animal at a time. For all individuals, however, right and left sides were observed in the same sequence of photographs at least once either before or during our study period, allowing profiles to be paired together for individual identification. Cameras were active for the same 99-day period that scat surveys were conducted. Photo sequences were considered independent if they were >30 minutes apart.

DNA extraction and species identification

We extracted DNA from scats using the Qiagen QIAamp DNA stool mini kit (Qiagen, Valencia, California) following the manufacturer's recommendations. For species identification, we amplified a 126-bp fragment of the mitochondrial gene ATP6 by polymerase chain reaction (PCR) using primers ATP6-DF3 and ATP6-DR1 following conditions from Chaves et al. [36]. PCR products were sequenced on an Applied Biosystems 3730xl DNA analyzer, and resulting sequences were compared to reference sequences using the online tool DNA Surveillance Carnivora [36].

Individual identification

Scat samples were genotyped at 4 microsatellite loci (FCA075, FCA077, FCA088, and FCA132) originally developed for the domestic cat (*Felis catus*) [37]. We initially screened 22 loci previously found to be variable in ocelots [38]. The 4 loci used for individual identification were selected based on degree of variability, success of amplification, and ease and clarity of allele scoring. As DNA from noninvasively collected fecal samples is often low quality and prone to genotyping errors such as allelic dropout and false alleles [39, 40], we used a multiple tubes approach [41] whereby each sample was genotyped three to nine times until reliable consensus genotypes were obtained. Genotypes were only accepted as reliable if a minimum of three identical heterozygote profiles, or five identical homozygote profiles, were observed (see Appendix 1 for PCR conditions). Checks for departure from Hardy-Weinberg equilibrium and probability of individual identity [42] were calculated using the program GENALEX [43].

Sex identification

For sex identification, we used felid-specific primers that amplify a 200-bp segment of the AMELY gene [44], which is only present on the Y-chromosome of males (See Appendix 1 for PCR conditions). PCR amplifications were performed in triplicate along with male and female positive controls and a negative control, and PCR products were visualized on agarose gel. Samples were identified as male if they showed amplification of the Y-linked marker for all three replicates, and were identified as female if no amplification was observed.

Density estimation

We used two CR models to estimate density from both our noninvasive genetic and camera trap data to provide a comparison of common approaches. First, we ran traditional closed CR analyses in the program CAPTURE [45] to estimate abundance. CAPTURE was implemented within the program MARK [46]. To

convert abundance estimates to density from our genetic data, we buffered sample locations with MMDM as well as $\frac{1}{2}$ -MMDM [3, 47], which can be conceptualized as a proxy for home range radius. We then used the sum of the resulting buffers to estimate ESA [3, 19]. For our camera trapping data, both the MMDM and the $\frac{1}{2}$ -MMDM buffer areas were larger than the area of the entire island, so we instead used island area (15.432 km²) as our ESA for estimating density. MMDM and ESA estimates were calculated in the program DENSITY [29]. Density estimates were then derived by dividing abundance estimates by ESA.

We also estimated density directly using the SECR model in the program DENSITY [29], which estimates density using a maximum-likelihood approach [25, 26, 48]. For our genetic data, we considered ocelot latrines as one might consider camera traps or hair snares, as discrete geographic locations where animals could be passively detected. DENSITY allows detectors to be considered as active or inactive during each sampling occasion. Thus, latrines known before commencement of the study were considered as active during the entire study period, and latrines discovered in the interim of the study were considered as inactive prior to discovery and active thereafter. Sampling occasions were three days, giving a total of 33 occasions during the study. All models were run under full likelihood with a Poisson distribution and the half-normal detection function. We ran multiple model combinations including various levels of individual heterogeneity, and we ran all model combinations both with and without sex as a covariate. The best-fit model was chosen using the Akaike Information Criterion adjusted for small sample size (AIC_c) [26, 48].

Results

Density from noninvasive genetic sampling

We collected 63 scats from 19 ocelot latrines on BCI, and we genetically confirmed 55 of these (87%) as being from ocelots. From these 55 scats, we obtained complete 4-locus genotypes from 43 samples (78%), which consisted of 12 unique genotypes and 31 recaptures from 16 latrines. Six individuals were identified as male, and six as female. All four loci were in Hardy-Weinberg equilibrium, and the number of alleles per locus was 3-6 (Appendix 2). Based on allele frequencies within the sampled population, probabilities of individual identity among unrelated individuals $P_{(ID)}$ was 0.00031, and among siblings $P_{(ID)sibs}$ was 0.038 [42].

The discriminant function model selection algorithm in CAPTURE selected model M_0 (the null model) as the most appropriate model (selection score of 1), with model $M_{h-jackknife}$ a close second (selection score of 0.94). As model M_0 is highly sensitive to violations of the assumption of homogeneous capture probability [45], and as some heterogeneity in capture probability at ocelot latrines is probable, we report results from both model M_0 and M_h (Table 1). Model M_0 produced an abundance estimate of 12 (SE = 0.918) and model M_h produced an abundance estimate of 15 (SE = 2.433). Capture probability was 0.189 for model M_0 and 0.152 for model M_h . MMDM from noninvasive genetic sampling was 632 m (SE = 117), resulting in ESA buffers of 9.760 km² (MMDM) and 3.114 km² ($\frac{1}{2}$ -MMDM). The program DENSITY selected the null model $g(0) \cdot \sigma(\cdot)$, without sex as a covariate, as the best-fitting model (Appendix 3), resulting in a density estimate of 1.742/km² (SE = 0.584).

Density from camera trapping

Twenty-one cameras were active during the 99-day sampling period resulting in 1,824 camera days. Seven cameras failed for short durations of the study, and this censoring was accounted for in the analyses [16]. This effort resulted in 398 independent photo sequences and 370 recaptures of 28 individual ocelots. Of these, four individuals were kittens photographed with their mother, which were excluded from further

density analyses. Mean number of captures per individual was 15.3 (SE = 2.71). Mean number of captures per camera was 17.5 (SE = 3.45). Of the 24 photographed adult ocelots, nine were males and 15 were females.

The discriminant function model selection algorithm in CAPTURE selected model $M_{h\text{-jackknife}}$ as the best-fitting model of those available. Model M_h is also the most commonly used model in camera trapping studies as it accounts for heterogeneity in capture probability [49]. Model M_h resulted in an abundance estimate of 28 (SE = 2.12) ocelots (Table 1), with a capture probability of 0.262. MMDM from camera trapping was 1,509 m (SE = 289). DENSITY selected the null model $g0(.)\sigma(.)$, with sex included as a covariate, as the best-fitting model (Appendix 4) resulting in a density estimate of 1.587/km² (SE = 0.464).

Table 1 Population density estimates of ocelots (*Leopardus pardalis*) from Barro Colorado Island, Panama. CAPTURE and DENSITY are software programs designed to estimate abundance or density using capture-recapture models. MMDM is a method of estimating effective survey area based on the mean of the maximum distance moved between captures.

Method and Model	Abundance (N)	Density		
		N/KM ²	SE	95% CI
Camera trapping				
CAPTURE $M_{h(jackknife)}$	28	1.815	0.163	1.686 – 2.464
DENSITY	-----	1.587	0.464	0.908 - 2.786
Noninvasive genetics				
CAPTURE M_0 - MMDM	12	1.229	0.094	1.229 - 1.844
CAPTURE M_0 - ½ MMDM	12	3.854	0.294	3.854 - 5.780
CAPTURE $M_{h(jackknife)}$ MMDM	15	1.537	0.249	1.332 - 2.459
CAPTURE $M_{h(jackknife)}$ ½ MMDM	15	4.817	0.781	4.174 - 7.707
DENSITY	-----	1.742	0.585	0.917 - 3.302

Discussion

Our estimates of ocelot population density derived from noninvasive genetics and camera trapping were highly comparable. Point estimates were similar, and 95% confidence intervals from all models overlapped, except those that used ½-MMDM to estimate ESA. Estimates from camera trapping had greater precision than those from noninvasive genetics, but these differences were not substantial (Table 1). Because we obtained density estimates from noninvasive genetics that were comparable, despite a relatively small sample size, to estimates from camera trapping, from which we had a large sample with many recaptures, our results support the validity of noninvasive genetic techniques for estimating density of elusive carnivores.

Density estimates from noninvasive genetics that used ½-MMDM to calculate ESA were substantially higher than other estimates, in some cases by more than a factor of three. Thus, we agree with other researchers that use of ½-MMDM can greatly overestimate density [10-12, 15, 19]. Our density estimates that used full MMDM to estimate ESA were more in line with our other estimates, but this result is likely

coincidental rather than biologically meaningful. Although MMDM methods have been used to estimate density in many studies, we agree with recent criticisms that use of MMDM methods has little theoretical or biological justification [15, 24]. This shortcoming is evident in our study, as MMDM distances calculated from camera trapping (1,509 m) and noninvasive genetics (632 m) were substantially different even though they were derived from the same population at the same time. The underestimation of ocelot movements from noninvasive genetics can likely be attributed to two factors. First, our camera array covered the entire island, whereas ocelot latrines had a more clumped distribution (Fig. 1). Second, our sample size from camera trapping was larger, and contained a far greater number of recaptures. It is logical that a large number of recaptures will be needed to effectively estimate movement distances [15], an issue widely acknowledged for estimation of home range size from telemetry data [50, 51]. Another problem with using MMDM to estimate survey area is that area buffers may contain both suitable and non-suitable habitat. For example, when we used MMDM and $\frac{1}{2}$ -MMDM to estimate ESA from camera trapping, both were larger than the area of the entire island and thus contained large portions of water, which is obviously not suitable ocelot habitat.

For our camera trapping dataset, DENSITY selected models that included sex as a covariate to be preferable to those that did not include sex (Appendix 4). This suggests, not surprisingly, that there is heterogeneity between the sexes in terms of movement and capture probability. Including sex as a covariate in the model, however, did not have a large effect on estimates of density. For our noninvasive genetic dataset, models that included sex as a covariate did not outperform those without sex (Appendix 3). This is likely due to our much smaller sample size from noninvasive genetics. Large datasets are typically necessary to benefit from inclusion of covariates, as covariates increase the number of model parameters [15].

For all of the reasons above, recently developed SECR models such as DENSITY [29] are clearly preferable for estimating population density in noninvasive studies. This is especially true for studies with small sample sizes, small survey areas [52], or in study locations with heterogeneous habitat. A key advantage is that SECR models do not require contentious post-hoc estimates of survey area to convert estimates of abundance to density [25, 53]. Furthermore, SECR models allow the user to specify suitable and non-suitable habitat for inclusion in the model, and they allow for inclusion of covariates such as sex [15]. Additionally, SECR models allow users to designate which detectors were operational during each sampling occasion, avoiding bias introduced by occasional camera trap failure common in traditional CR models [16].

Despite our efforts to obtain scat samples from throughout BCI, the spatial extent of our effective sampling area was unequal between camera trapping and noninvasive genetics, due to the difficulty of finding ocelot latrines. This difference reflects the reality that most researchers will face when conducting studies that rely on collection of scats. With camera trapping, researchers pre-define the sampling area by choosing the placement of camera traps. With fecal noninvasive genetic sampling, effective sampling area is instead defined by where study animals defecate, and by the ability of researchers to recover scat samples [1]. Fortunately, SECR models take differences in sampling area into account when estimating density, making it possible to compare density estimates from different spatial extents. We think that our ability to estimate similar densities from noninvasive genetics and camera trapping, despite differences in spatial extent, lends further support to noninvasive genetics as a viable method.

Our estimated ocelot densities were higher than estimates reported from anywhere else in the species range [54, 55]. Even our density estimates based on minimum number known alive ($1.55/\text{km}^2$; a

conservative lower bound estimate) are 1.6 times greater than the densest ocelot population previously reported from the northwestern Amazon of Peru ($0.947/\text{km}^2$) [55]. Our unusually high densities could be the result of numerous factors, including high prey availability [56], frustrated dispersal due to the high cost of dispersing from island to mainland across open water, or the vigilant protection BCI receives from poaching of both ocelots and their prey. High densities could also be the result of mesopredator release, as jaguars (*Panthera onca*) and pumas (*Puma concolor*) were resident in the BCI area in the early 20th century, but are now only infrequent visitors to the island [57, 58].

Implications for conservation

Our study provides evidence that noninvasive genetic techniques can generate accurate estimates of population density, especially when used in conjunction with SECR models. We do caution, however, that further studies should be conducted on a variety of species in diverse environments to further verify the accuracy of these techniques. Our study was conducted in a high-density population of a species that defecates at latrine sites, and thus we were able to find scats relatively easily. For low density populations, it may be necessary to use aids such as scat detection dogs, which although expensive can greatly increase detection rates [22]. In sum, we think that fecal noninvasive genetics provides a promising tool for the estimation of density in elusive animals, especially for species in which individuals cannot be identified from camera trapping. These techniques have great potential to aid in the conservation of the many elusive species worldwide for which population density and conservation status are poorly known.

Acknowledgements

We thank Patrick Jansen, Roland Kays, Ricardo Moreno and Joseph Wright for help with study design, George Smallwood for field assistance, and Natalie Pawlikowski and James Lyko for help with photo identification. Charles Kontos, Jr. helped to establish the ocelot photo identification procedure. Gregory Willis provided financial support and mammal census fieldwork. This work was funded by the Smithsonian Tropical Research Institute, American Society of Mammalogists, and University of Illinois Graduate College.

References

- [1] Rodgers, T. and Janečka, J. 2013. Applications and techniques for non-invasive faecal genetics research in felid conservation. *European Journal of Wildlife Research* 59:1-16.
- [2] Karanth, K. U. 1995. Estimating tiger *panthera-tigris* populations from camera-trap data using capture-recapture models. *Biological Conservation* 71:333-338.
- [3] Karanth, K. U. and Nichols, J. D. 1998. Estimation of tiger densities in India using photographic captures and recaptures. *Ecology* 79:2852-2862.
- [4] Kohn, M. H., York, E. C., Kamradt, D. A., Haugt, G., Sauvajot, R. M. and Wayne, R. K. 1999. Estimating population size by genotyping faeces. *Proceedings of the Royal Society of London Series B-Biological Sciences* 266:657-663.
- [5] Perez, I., Geffen, E. and Mokady, O. 2006. Critically Endangered Arabian leopards *Panthera pardus nimr* in Israel: estimating population parameters using molecular scatology. *Oryx* 40:295.
- [6] Trolle, M. and Kery, M. 2003. Estimation of ocelot density in the pantanal using capture-recapture analysis of camera-trapping data. *Journal of Mammalogy* 84:607-614.
- [7] Sollmann, R., Mohamed, A., Samejima, H. and Wilting, A. 2013. Risky business or simple solution - Relative abundance indices from camera-trapping. *Biological Conservation* 159:405-412.

- [8] Web of Science. Accessed 11-June-2013. <http://thomsonreuters.com/thomson-reuters-web-of-science/>.
- [9] Balme, G. A., Hunter, L. T. B. and Slotow, R. 2009. Evaluating methods for counting cryptic carnivores. *Journal of Wildlife Management* 73:433-441.
- [10] Soisalo, M. K. and Cavalcanti, S. M. C. 2006. Estimating the density of a jaguar population in the Brazilian Pantanal using camera-traps and capture-recapture sampling in combination with GPS radio-telemetry. *Biological Conservation* 129:487-496.
- [11] Dillon, A. and Kelly, M. J. 2008. Ocelot home range, overlap and density: comparing radio telemetry with camera trapping. *Journal of Zoology* 275:391-398.
- [12] Sharma, R. K., Jhala, Y., Qureshi, Q., Vattakaven, J., Gopal, R. and Nayak, K. 2010. Evaluating capture-recapture population and density estimation of tigers in a population with known parameters. *Animal Conservation* 13:94-103.
- [13] Nunez-Perez, R. 2011. Estimating jaguar population density using camera-traps: a comparison with radio-telemetry estimates. *Journal of Zoology* 285:39-45.
- [14] Blanc, L., Marboutin, E., Gatti, S. and Gimenez, O. 2013. Abundance of rare and elusive species: Empirical investigation of closed versus spatially explicit capture-recapture models with lynx as a case study. *Journal of Wildlife Management* 77:372-378.
- [15] Tobler, M. W. and Powell, G. V. N. 2013. Estimating jaguar densities with camera traps: Problems with current designs and recommendations for future studies. *Biological Conservation* 159:109-118.
- [16] Foster, R. J. and Harmsen, B. J. 2012. A critique of density estimation from camera-trap data. *Journal of Wildlife Management* 76:224-236.
- [17] O'Connell, A. F., Nichols, J. D. and Karanth, K. U., Eds., *Camera traps in animal ecology*, (Springer, Japan, 2011).
- [18] Solberg, K. H., Bellemain, E., Drageset, O.-M., Taberlet, P. and Swenson, J. E. 2006. An evaluation of field and non-invasive genetic methods to estimate brown bear (*Ursus arctos*) population size. *Biological Conservation* 128:158-168.
- [19] Janečka, J., Munkhtsog, B., Jackson, R., Naranbaatar, G., Mallon, D. and Murphy, W. 2011. Comparison of noninvasive genetic and camera-trapping techniques for surveying snow leopards. *Journal of Mammalogy* 92:771-783.
- [20] Mondol, S., Ullas Karanth, K., Samba Kumar, N., Gopaldaswamy, A. M., Andheria, A. and Ramakrishnan, U. 2009. Evaluation of non-invasive genetic sampling methods for estimating tiger population size. *Biological Conservation* 142:2350-2360.
- [21] Hedges, S., Arlyne Johnson, Marissa Ahlering, Martin Tyson and Eggert, L. S. 2013. Accuracy, precision, and cost-effectiveness of conventional dung density and fecal DNA based survey methods to estimate Asian elephant (*Elephas maximus*) population size and structure. *Biological Conservation* 159:101-108.
- [22] Kelly, M. J., Betsch, J., Wultsch, C., Mesa, B. and Mills, L. S. 2012. Noninvasive sampling for carnivores in *Carnivore ecology and conservation: a handbook of techniques*, L. P. R. A. Boitani, Ed. Oxford University Press, USA: 47-69.
- [23] Miotto, R. A., Cervini, M., Kajin, M., Begotti, R. A. and Galetti, P. M. 2014. Estimating puma Puma concolor population size in a human-disturbed landscape in Brazil, using DNA mark-recapture data. *Oryx* 48:250-257.
- [24] Pesenti, E. and Zimmermann, F. 2013. Density estimations of the Eurasian lynx (*Lynx lynx*) in the Swiss Alps. *Journal of Mammalogy* 94:73-81.
- [25] Efford, M. 2004. Density estimation in live-trapping studies. *Oikos* 106:598-610.
- [26] Borchers, D. L. and Efford, M. G. 2008. Spatially explicit maximum likelihood methods for capture-recapture studies. *Biometrics* 64:377-385.
- [27] Royle, J. A. and Young, K. V. 2008. A hierarchical model for spatial capture-recapture data. *Ecology* 89:2281-2289.

- [28] Anile, S., Ragni, B., Randi, E., Mattucci, F. and Rovero, F. 2014. Wildcat population density on the Etna volcano, Italy: a comparison of density estimation methods. *Journal of Zoology* 293:252-261.
- [29] Efford, M. G. 2012. Density 5.0: software for spatially explicit capture-recapture. Department of Mathematics and Statistics, University of Otago, Dunedin, New Zealand.
<http://www.otago.ac.nz/density>
- [30] IUCN. 2014. 2014 IUCN Red List of Threatened Species. www.iucnredlist.org
- [31] Leigh, E. G., Jr. 1999. *Tropical forest: A view from Barro Colorado Island* Oxford University Press, New York
- [32] Moreno, R. and Giacalone, J. 2008. Ecological data obtained from latrine use by ocelots (*Leopardus pardalis*) on Barro Colorado Island, Panama. *Tecnociencia* 8:7-21.
- [33] Rodgers, T. W., Giacalone, J., Heske, E. J., Janečka, J. E., Phillips, C. A., Jansen, P. A. and Schooley, R. L. *in press*. Socio-spatial organization and kin structure in ocelots from integration of camera trapping and noninvasive genetics. *Journal of Mammalogy*
- [34] Moreno, R., Kays, R., Giacalone-Willis, J., Aliaga-Rossel, E., Mares, R. and Bustamante, A. 2012. Home range and circadian activity of ocelots (*leopardus pardalis*) in Barro Colorado Island, Panama. *Mesoamericana* 16:30-39.
- [35] Noss, A. J., Gardner, B., Maffei, L., Cuellar, E., Montano, R., Romero-Munoz, A., Sollman, R. and O'Connell, A. F. 2012. Comparison of density estimation methods for mammal populations with camera traps in the Kaa-Iya del Gran Chaco landscape. *Animal Conservation* 15:527-535.
- [36] Chaves, P. B., Graeff, V. G., Lion, M. B., Oliveira, L. R. and Eizirik, E. 2012. DNA barcoding meets molecular scatology: short mtDNA sequences for standardized species assignment of carnivore noninvasive samples. *Molecular Ecology Resources* 12:18-35.
- [37] Menotti-Raymond, M., David, V. A., Lyons, L. A., Schaffer, A. A., Tomlin, J. F., Hutton, M. K. and O'Brien, S. J. 1999. A genetic linkage map of microsatellites in the domestic cat (*Felis catus*). *Genomics* 57:9-23.
- [38] Janečka, J. E., Tewes, M. E., Laack, L. L., Caso, A., Grassman, L. I., Haines, A. M., Shindle, D. B., Davis, B. W., Murphy, W. J. and Honeycutt, R. L. 2011. Reduced genetic diversity and isolation of remnant ocelot populations occupying a severely fragmented landscape in southern Texas. *Animal Conservation* 14:608-619.
- [39] Taberlet, P., Waits, L. P. and Luikart, G. 1999. Noninvasive genetic sampling: look before you leap. *Trends in Ecology and Evolution* 14:323-327.
- [40] Broquet, T., Menard, N. and Petit, E. 2007. Noninvasive population genetics: a review of sample source, diet, fragment length and microsatellite motif effects on amplification success and genotyping error rates. *Conservation Genetics* 8:249-260.
- [41] Taberlet, P. and Fumagalli, L. 1996. Owl pellets as a source of DNA for genetic studies of small mammals. *Molecular Ecology* 5:301-305.
- [42] Waits, L. P., Luikart, G. and Taberlet, P. 2001. Estimating the probability of identity among genotypes in natural populations: cautions and guidelines. *Molecular Ecology* 10:249-256.
- [43] Peakall, R. and Smouse, P. E. 2006. GENALEX 6: genetic analysis in Excel. Population genetic software for teaching and research. *Molecular Ecology Notes* 6:288-295.
- [44] Murphy, W. J., Sun, S., Chen, Z. Q., Pecon-Slattery, J. and O'Brien, S. J. 1999. Extensive conservation of sex chromosome organization between cat and human revealed by parallel radiation hybrid mapping. *Genome Research* 9:1223-1230.
- [45] Otis, D. L., Burnham, K. P., White, G. C. and Anderson, D. R. 1978. Statistical inference from capture data on closed animal populations. *Wildlife Monographs*:1-135.
- [46] White, G. C. and Burnham, K. P. 1999. Program MARK: survival estimation from populations of marked animals. *Bird Study* 46:120-139.

- [47] Wilson, K. R. and Anderson, D. R. 1985. Evaluation of 2 density estimators of small mammal population-size. *Journal of Mammalogy* 66:13-21.
- [48] Efford, M. G., Dawson, D. K. and Borchers, D. L. 2009. Population density estimated from locations of individuals on a passive detector array. *Ecology* 90:2676-2682.
- [49] Harmsen, B. J., Foster, R. J. and Doncaster, C. P. 2011. Heterogeneous capture rates in low density populations and consequences for capture-recapture analysis of camera-trap data. *Population Ecology* 53:253-259.
- [50] Harris, S., Cresswell, W. J., Forde, P. G., Trehwella, W. J., Woollard, T. and Wray, S. 1990. Home-range analysis using radio-tracking data - a review of problems and techniques particularly as applied to the study of mammals. *Mammal Review* 20:97-123.
- [51] Seaman, D. E., Millspaugh, J. J., Kernohan, B. J., Brundige, G. C., Raedeke, K. J. and Gitzen, R. A. 1999. Effects of sample size on kernel home range estimates. *Journal of Wildlife Management* 63:739-747.
- [52] Sollmann, R., Gardner, B. and Belant, J. L. 2012. How does spatial study design influence density estimates from spatial capture-recapture models? *PLoS One* 7
- [53] Borah, J., Sharma, T., Das, D., Rabha, N., Kakati, N., Basumatary, A., Ahmed, M. F. and Vattakaven, J. 2014. Abundance and density estimates for common leopard *Panthera pardus* and clouded leopard *Neofelis nebulosa* in Manas National Park, Assam, India. *Oryx* 48:149-155.
- [54] Di Bitetti, M. S., Paviolo, A., De Angelo, C. D. and Di Blanco, Y. E. 2008. Local and continental correlates of the abundance of a neotropical cat, the ocelot (*Leopardus pardalis*). *Journal of Tropical Ecology* 24:189-200.
- [55] Kolowski, J. M. and Alonso, A. 2010. Density and activity patterns of ocelots (*Leopardus pardalis*) in northern Peru and the impact of oil exploration activities. *Biological Conservation* 143:917-925.
- [56] Terborgh, J. 1992. Maintenance of diversity in tropical forests. *Biotropica* 24:283-292.
- [57] Wright, S. J., Gompper, M. E. and Deleon, B. 1994. Are large predators keystone species in neotropical forests - the evidence from Barro-Colorado Island. *Oikos* 71:279-294.
- [58] Moreno, R. S., Kays, R. W. and Samudio, R. 2006. Competitive release in diets of ocelot (*Leopardus pardalis*) and puma (*Puma concolor*) after jaguar (*Panthera onca*) decline. *Journal of Mammalogy* 87:808-816.

Appendix 1. PCR Conditions

PCR conditions for species identification can be found in Chaves et al. (2012). PCR conditions for microsatellite amplification and sex ID were as follows. Reactions included 1 µl 10X Buffer, 0.8 µl of MgCl, 0.2 µl of 10 mM DNTs, 0.2 µl of 20 mM fluorescently-labeled forward primer, 0.2 µl of 20 mM reverse primer, 0.04 µl of AmpliTaq Gold® 360 DNA Polymerase (Life Technologies; Grand Island, NY, USA), 0.1µl of BSA, 0.2 µl of GC-enhancer, 5.76 µl of water, and 1.5 µl of template DNA for a total reaction volume of 10 µl. Thermocycling conditions were as follows: initial denaturation at 95°C/10', followed by 50 cycles of 95°C/5", 55°C/15", 72°C/45", and a final extension of 72°C/10'. Microsatellite forward primers were fluorescently labeled with one of four dyes (FAM, NED, PET, or VIC). Each microsatellite was amplified individually and then the loci were pooled together and genotyped on an Applied Biosystems 3730xl DNA analyzer.

Appendix 2. Summary of 4 microsatellite loci used for individual identification. H_o is observed heterozygosity. H_e is expected heterozygosity under Hardy-Weinberg equilibrium. All loci were in Hardy-Weinberg equilibrium (HWE probability > 0.05).

Locus	Size	# alleles	H_o	H_e	HWE probability
FCA075	110-128	5	0.769	0.782	0.826
FCA077	133-137	3	0.615	0.492	0.463
FCA088	92-110	5	0.846	0.809	0.564
FCA132	167-179	6	0.769	0.788	0.559
Mean	na	4.75	0.762	0.7226	0.5872

Appendix 3. AIC table for density estimates from noninvasive genetic sampling from the program DENSITY.

Model	Sex as covariate	# Parameters	ML Log likelihood	AICc	Δ AICc	Density	SE	95% CI
g0[.]s[.]	No	3	-186.49	381.97	--	1.7423	0.5846	0.919-3.305
g0[h2]s[.]	No	5	-186.00	392.00	10.03	1.7417	0.5851	0.918-3.306
g0[.]s[h2]	No	5	-186.45	392.90	10.93	1.7410	0.5828	0.919-3.298
g0[h2]s[h2]	No	6	-185.86	400.52	18.55	1.7456	0.5874	0.919-3.317
g0[.]s[.]	Yes	6	-191.58	411.95	29.98	1.8074	0.9057	0.717-4.577
g0[h2]s[.]	Yes	8	-191.11	446.21	64.24	1.8186	0.9176	0.717-4.629
g0[.]s[h2]	Yes	8	-191.57	447.15	65.18	1.8057	0.9046	0.716-4.570
g0[h2]s[h2]	Yes	9	-191.58	491.15	109.1	1.8053	0.9085	0.713-4.587

Appendix 4. AIC table for density estimates from camera trapping from the program **DENSITY**.

Model	Sex as covariate	# Parameters	ML Log likelihood	AICc	ΔAIC_c	Density	SE	95% CI
g0[.]s[.]	Yes	6	-1072.52	2161.98	--	1.5867	0.463	0.908-2.786
g0[.]s[h2]	Yes	8	-1071.33	2168.26	6.28	1.5835	0.463	0.906-2.782
g0[h2]s[.]	Yes	8	-1071.43	2168.46	6.48	1.5842	0.463	0.906-2.783
g0[h2]s[h2]	Yes	9	-1071.33	2173.51	11.53	1.5826	0.463	0.905-2.781
g0[.]s[.]	No	3	-1115.23	2237.65	75.67	1.5617	0.322	1.105-2.330
g0[.]s[h2]	No	5	-1113.69	2240.71	78.73	1.5575	0.321	1.104-2.325
g0[h2]s[.]	No	5	-1114.12	2241.57	79.59	1.5613	0.322	1.105-2.330
g0[h2]s[h2]	No	6	-1113.92	2244.77	82.79	1.5551	0.321	1.104-2.321