

Comparison of Soil Microbial Composition and Diversity Between Mixed and Monoculture Rubber Plantations in Hainan Province, China

Authors: Jatoi, Muhammad Tahir, Lan, Guoyu, Wu, Zhixiang, Sun, Rui, Yang, Chuan, et al.

Source: Tropical Conservation Science, 12(1)

Published By: SAGE Publishing

URL: <https://doi.org/10.1177/1940082919876072>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Comparison of Soil Microbial Composition and Diversity Between Mixed and Monoculture Rubber Plantations in Hainan Province, China

Tropical Conservation Science
Volume 12: 1–9
© The Author(s) 2019
Article reuse guidelines:
sagepub.com/journals-permissions
DOI: 10.1177/1940082919876072
journals.sagepub.com/home/trc



Muhammad Tahir Jatoi^{1,2} , Guoyu Lan^{2,3,4,5}, Zhixiang Wu^{2,3,4,5} ,
Rui Sun^{2,3}, Chuan Yang^{2,3}, and Zhenghong Tan¹

Abstract

This study aimed to compare monoculture and mixed rubber plantations in terms of their soil bacterial and fungal composition. An Illumina MiSeq sequencing analysis was performed to investigate the composition and diversity of the soil bacterial and fungal communities among three different rubber (*Hevea brasiliensis*) plantations: monoculture, Mixed 1 (*Hevea brasiliensis* and *Mytilaria laosensis*), and Mixed 2 (*Hevea brasiliensis* and *Michelia macclurei*) in Hainan. The results showed that the bacterial composition of the three rubber plantations was basically similar. However, there was a significant difference in fungal communities among the three rubber plantations at both the phylum and operational taxonomic unit level. The species richness, Chao, and Shannon diversity of bacterial communities of monoculture rubber plantations were higher than the Mixed 1 and Mixed 2 rubber plantations, whereas all diversity indexes of fungal communities were relatively equal for the monoculture and mixed rubber plantations. Soil nutrition (such as total nitrogen and total potassium) and soil pH are the main drivers of the bacterial composition ($p < .001$). However, soil pH and water content are the main drivers of the fungal composition ($p < .001$), and to some extent, soil pH can increase soil bacteria diversity. We suggest that alkaline fertilizers should be applied in mixed rubber plantations to improve the soil pH and, consequently, to increase the total diversity of the rubber plantation.

Keywords

soil microbial composition, bacterial diversity, monoculture rubber plantation, mixed rubber plantation, soil chemical properties

Introduction

Rubber (*Hevea brasiliensis* Muell. Arg) is an important economic crop in the world, and its cultivation is one of the main sources of income in tropical region of Asia. During the last two decades, rubber plantations (RPs) seem to have extended quickly throughout Asia (Li et al., 2015), and a large area of natural forests have been changed into RPs (Chen et al., 2017; Yan et al., 2015). Deforestation influences local temperature, light, dampness, and litter conditions, bringing about changes of microhabitats and loss of forests (Chaudhary, Burivalova, Koh, & Hellweg, 2016). The conservation of tropical forests has been found to have global importance (Cole, Bhagwat, & Willis, 2014; Naeem, Duffy, & Zavaleta, 2012). Currently, southeast Asian croplands are under pressure due to increasing population and demands of crop products in the world (Koh, Miettinen, Liew, & Ghazoul, 2011; Sodhi, Koh, Brook, & Ng, 2004; Wilcove, Giam, Edwards, Fisher, &

¹College of Ecology and Environment, Hainan University, Haikou City, P. R. China

²Rubber Research Institute, Chinese Academy of Tropical Agricultural Sciences, Danzhou City, P. R. China

³Danzhou Investigation & Experiment Station of Tropical Crops, Ministry of Agriculture and Rural Affairs, Danzhou City, P. R. China

⁴Key Laboratory of Biology and Genetic Resources of Rubber Tree, Ministry of Agriculture and Rural Affairs, Danzhou City, P. R. China

⁵State Key Laboratory Incubation Base for Cultivation & Physiology of Tropical Crops, Danzhou City, P. R. China

Received 2 April 2019; Accepted 21 August 2019

Corresponding Author:

Guoyu Lan, Rubber Research Institute, Chinese Academy of Tropical Agricultural Sciences, Danzhou City, Hainan Province, 571737, P. R. China
Email: langyrrri@163.com

Zhixiang Wu, Rubber Research Institute, Chinese Academy of Tropical Agricultural Sciences, Danzhou City, Hainan Province, 571737, P. R. China
Email: wzxrrri@163.com



Creative Commons Non Commercial CC BY-NC: This article is distributed under the terms of the Creative Commons Attribution-NonCommercial 4.0 License (<http://www.creativecommons.org/licenses/by-nc/4.0/>) which permits non-commercial use, reproduction and distribution of the work without further permission provided the original work is attributed as specified on the SAGE and Open Access pages (<https://us.sagepub.com/en-us/nam/open-access-at-sage>).

Koh, 2013). This has aroused the interest of biodiversity scholars around the world. Hainan is considered to be the largest island in the Indo-Burma biodiversity hotspot and harbors many endangered species (Myers, Mittermeier, Mittermeier, Fonseca, & Kent, 2000). Now, RPs compose one fourth of the vegetation on Hainan Island (Wang, Lan, Wu, & Xie, 2012), and they are one of the most important vegetation compositions in Hainan Province (Lan, Wang, Wu, & Xie, 2013). Studies have shown that cropping systems involving the removal of an agricultural byproduct can reduce soil organic carbon (Singh, Rai, & Singh, 2016). Moreover, forest conversion also affects the composition of soil microorganisms (Cai, Zhang, Yang, & Wang, 2018; Rodrigues et al., 2013).

Microbial activity plays a key role in nutrient cycling and amelioration of plant stresses and is responsible for a broad range of the ecological functions of soil (Singh et al., 2016). The soil microbial diversity is a critical marker of soil quality (He, Zheng, Chen, He, & Zhang, 2008; Kirk et al., 2004). It was reported that soil microbial diversity is affected by a variety of environmental factors (Balser, Kinzig, & Firestone, 2002; Schimel, 1995). Previous findings revealed that abiotic soil nutrients drive the soil bacterial communities on RPs in Hainan (Lan, Li, Wu, & Xie, 2017a). Soil pH is another important factor that can affect the soil bacterial community structure (Krashevskaya, Klarner, Widyastuti, Maraun, & Scheu, 2015). Seasonal changes are also main drivers of the bacterial composition of RPs and other tropical forests (Lan, Li, Lesueur, Wu, & Xie, 2018). In addition, other factors controlling the distribution of soil microorganisms are being studied (Rousk et al., 2010). For instance, it is important to know whether different planting patterns of RP (monoculture and mixed plantations) affect the composition and diversity of soil microbes.

Monoculture has often been criticized for reducing biodiversity, causing widespread outbreaks of pests and diseases (Erskine, Lamb, & Bristow, 2006; Lamb, Erskine, & Parrotta, 2005) and not supporting numerous conventional ecosystem services (Singh, Rai, Banyal, Chauhan, & Singh, 2018). Thus, there is an emphasis on growing mixed plantations rather than monoculture. This includes an expanded yield, better natural assurance, biodiversity protection, and remodeling (Montagnini, Gonzales, & Porras, 1995). We aimed to test if there were any differences in the composition and diversity of soil microorganisms between monoculture and mixed RPs. In this study, we examined the bacterial and fungal composition and diversity of monoculture and mixed RPs by utilizing an extensive throughput Illumina MiSeq sequencing analysis. The objectives of this study were (a) to identify the soil microbial composition and changes in community structure among

monoculture, Mixed 1 and Mixed 2 plantations and (b) to understand the main factors that affect the microbial community. We aimed to test whether soil microbial diversity in mixed RPs was higher than in a pure RP.

Methods

Study Site

Hainan Province (18°10' –20°10'N and 108°37' –111°03'E), with a territory of 33,920 km², is located in the south of China. Hainan Island is one of the most important hot spots of biodiversity conservation in the Indo-Burma region. Hainan Island's tropical rainforests are situated at the northern edge of tropical Asia (Zhu & Zhou, 2002) and are recognized for their high biodiversity (Zheng, Cour, & Zou, 2002). This island has a tropical rainstorm environment with a wet period from May to October and a dry period from November to April. The annual precipitation was recorded more than 1,600 mm and the precipitation was irregularly distributed between wet and dry seasons (Francisco-Ortega et al., 2010). The lowest mean monthly temperature was recorded in January (16°C) and the highest in July 27.5°C. Precipitation frequently occurs on the eastern part of Hainan Island because of storms in the Pacific Ocean, while the western part of the island is dry (Xu, Tan, Wang, & Gai, 2016).

Sample Collection

The study site was located in Wenchang area, northeast of Hainan Island. We selected three plots (20 m × 20 m) on three RPs, that is, a monoculture RP, a Mixed 1 RP (*Hevea brasiliensis* and *Mytilaria laosensis*), and a Mixed 2 RP (*Michelia macclurei*). Soil sampling design in three rubber plantations were shown in Figure A1. Monoculture RPs have understory herbaceous species such as *Parabaena sagittata*, *Cyrtococcum patens*, *Eupatorium odoratum*, and *Mimosa pudica*. Because of the high canopy density, the coverage of the herbaceous plants in Mixed 1 is relatively lower than in the monoculture. Mixed 1 comprised *Cyrtococcum patens* and *Ageratum conyzoides*. The coverage of Mixed 2 comprised *Cyrtococcum patens* and *Eupatorium odoratum*.

Soil samples were randomly collected from each RP. Six soil samples were randomly gathered within each plot by using a 5 cm diameter stainless steel cylinder. The soil samples were then separated into three sections; the first was sieved throughout a 2 mm net instantly and stored at 4°C until analysis of MBC; the second one was air-dried and passed through a 0.25 mm sieve for SOM, soil pH, and TN; and the third one remained stored at –80°C for DNA extraction. The soil pH was estimated in a 1:1 soil:water mixture. The soil moisture was

determined gravimetrically. Total phosphorus (TP) was digested by molybdenum-antimony, and measurement was performed by spectrophotometric method (Soil Science Society of China, 2000). Total potassium (TK) content was measured by flame photometry. MBC was examined by chloroform fumigation and an extraction technique utilizing the correction factor 0.35 (Shen et al., 2014). The forests and soil properties are shown in Table 1.

DNA Extraction, PCR Amplification, and Illumina MiSeq Sequencing

Microbial DNA was extracted from 5.0 g of soil using the E.Z.N.A.[®] Soil DNA Kit (Omega Biotek, Norcross, GA, USA) according to the manufacturer's protocols. For bacteria, the V3–V4 hypervariable regions of the bacterial 16S rRNA gene were amplified with primers 338 F (5'-ACTCCTACGGGAGGCAGCAG-3') and 806 R (5'-GGACTACHVGGGTWTCTAAT-3') by a thermocycler polymerase chain reaction system (Xu et al., 2016; GeneAmp 9700, ABI, USA). Fungal Internal Transcribed Spacer (ITS) Region 1 was amplified using the ITS1F (5'-CTTGGTCATTTAGAGG AAGTAA-3') and ITS2 (5'-GCTGCGTTCTTCATCG ATGC-3') primer pair (Kerfahi, Tripathi, Dong, Go, & Adams, 2016). Polymerase chain reactions (PCRs) were conducted in triplicate and the tests were conducted using a 20 µL blend consisting of 4 µL of 5 × FastPfu Buffer, 2 µL of 2.5 mM dNTPs, 0.8 µL of the sample each primer (5 µM), 0.4 µL of FastPfu Polymerase, and 10 ng of template DNA. Amplicons were separated on 2% agarose gels and purified utilizing the AxyPrep DNA gel extraction kit (Axygen Biosciences, Union City, CA, USA) as indicated by the manufacturer's instructions and measured utilizing QuantiFluorTM—ST (Promega, USA). Purged amplicons were mutually equimolar and combined end-sequenced (2 × 250) on an Illumina MiSeq stage was conducted as indicated by standard conventions.

Bioinformatics and Statistical Analyses

Basic FASTQ records were demultiplexed and separated utilizing QIIME (version 1.17). The adjusted groupings were collected into operational taxonomic units (OTUs)

characterized through 97% comparability (Stackebrandt & Goebel, 1994) utilizing the CD-HIT-OTU driver (Wu, Zhu, Fu, Niu, & Li, 2011). The phylogenetic attachments of all 16S and 18S rRNA gene arrangements were examined by the Ribosomal Database Project classifier of the SILVA (SSU117/119) 16S and 18S rRNA catalog utilizing a confidence threshold of 70%. We calculated the exposure proportion using Good's technique. The Shannon diversity index and nonparametric measures of genus richness (Chao1) were designed for each illustration using Mothur (Schloss et al., 2009). Principal coordinate analysis (PCoA) based on the Bray-Curtis distance matrix of family composition was achieved. To check the microbial community's similarity index, the similarities were statistically analyzed to test whether there was any significant variance among the RPs.

The seven ecological variables included in the analysis were the type of RP (1: monoculture RP; 2: Mixed 1 RP; 3: Mixed 2 RP), soil organic matter (SOM), TP, total nitrogen (TN), TK, water content (WC), and soil pH. To identify relationship between microbial communities and ecological factors, a redundancy analysis was conducted. Using the *Vegan* package within R, seven variables and taxonomic composition (family level) data in 18 communities were analyzed. The numerical consequences were evaluated utilizing Monte Carlo permutation's technique on 999 permutations. The statistical analyses were conducted using SPSS 16.0. Duncan's technique was utilized to test the significant differences in compositions and diversity among the mixed and monoculture RPs.

Results

Taxonomic Compositions

There were a total of 1,896 OTUs belonging to 170 families and 7 phyla in the soil fungal communities from 18 soil samples (6 monoculture RPs, 6 Mixed 1 RPs, and 6 Mixed 2 RPs), among which there were a total of 1,758 OTUs belonging to 196 families and 23 phyla in the soil bacterial communities. At the phylum level, there was no difference in soil bacterial compositions between the monoculture and mixed RPs (Figure 1(a)). For fungal structure, the most abundant phylum in soils of monoculture, Mixed 1, and Mixed 2 RPs was Ascomycota.

Table 1. Soils Characteristics of the Three Rubber Plantations in Wenchang, Hainan China.

Forest types	pH	WC (%)	MBC (mg/kg)	SOM (g/kg)	TN (g/kg)	TP (g/kg)	TK (g/kg)
Monoculture RP	4.62 ± 0.03a	32.42 ± 0.79a	96.32 ± 11.01a	4.36 ± 0.43a	0.19 ± 0.01a	0.17 ± 0.00a	0.03 ± 0.00b
Mixed 1 RP	4.45 ± 0.01c	28.60 ± 0.35b	87.21 ± 9.32b	4.01 ± 0.18a	0.17 ± 0.00a	0.16 ± 0.01a	0.04 ± 0.00a
Mixed 2 RP	4.52 ± 0.01b	28.36 ± 0.59b	86.71 ± 8.36b	4.19 ± 0.28a	0.18 ± 0.00a	0.17 ± 0.01a	0.03 ± 0.00b

Note. Values shown are means and standard error ($n = 6$). Statistically significant difference at $p < .05$. MBC = microbial biomass carbon; SOM = soil organic matter; TN = total nitrogen; TP = total phosphorus; TK = total potassium; WC = water content; pH = soil pH.




Figure 1. Phylum compositions of soil bacterial (a) and fungal (b) communities in the three rubber plantations (monoculture, Mixed 1, Mixed 2 RPs) * $p < .05$.




Figure 2. Observed OTU richness and Chao I and Shannon diversity of soil bacteria (a–c) and fungi species level communities in the three rubber plantations (monoculture, Mixed 1, Mixed 2 RPs). a,b,c, $p < 0.05$

The relative abundance of Ascomycota was higher in monoculture soil than in the mixed RPs; however, the relative abundance of Basidiomycota was lower in the monoculture (Figure 1(b)). Mixed 1 and Mixed 2 RPs were fundamentally similar.

Bacterial Diversity

Diversity estimations at the OTU level of the 16S and 18S rRNA gene banks of three communities from Illumina MiSeq sequencing analysis are shown in Figure 2. In terms of soil bacterial community, the number of OTUs

in the monoculture RPs was significantly higher than in the Mixed 1 and Mixed 2 RPs, whereas Mixed 2 was higher than Mixed 1. The Chao1 diversity and Shannon diversity indexes showed that monoculture was significantly higher than the Mixed 1 RP. The number of species, Chao diversity and Shannon diversity indexes of fungal communities had no significant differences among the monoculture, Mixed 1, and Mixed 2 RPs.

PCoA of the bacterial and fungal communities at the OTU level of the three RPs showed some spatial heterogeneity of bacterial and fungal taxonomic compositions between the soils of the monoculture and mixed RPs




Figure 3. PCoA of bacterial (a) and fungal (b) communities of the three rubber plantations. Points those are closer together on the ordination show communities that are more similar (monoculture, Mixed I, Mixed 2 RPs). PCoA = principal coordinates analysis.




Figure 4. Redundancy analysis ordination of the soil samples and family compositions of bacterial (a) and fungal (b) communities across the three rubber plantations (monoculture, Mixed I, Mixed 2 RPs). SOM = soil organic matter; TN = total nitrogen; TP = total phosphorus; TK = total potassium; WC = water content; pH = soil pH. Different values were assigned to the three rubber plantations (1: monoculture; 2: Mixed I; 3: Mixed 2 RPs), thus the direction of the arrow of the rubber forest points to the mixed forest.

(Figure 3). For bacteria, the analysis of similarity (ANOSIM) results showed that there was no significant difference in taxonomic composition ($p = .0569$, at the OTU level) among the monoculture, Mixed 1, and Mixed 2 RPs. However, the ANOSIM results showed that monoculture, Mixed 1, and Mixed 2 RPs were different from each other in fungal taxonomic composition at the OTU level ($p < .001$).

Affecting Factors

Compared to Mixed 1 and Mixed 2, the WC was higher in the monoculture. Furthermore, the soil pH in the

monoculture was significantly higher than in the Mixed 1 and Mixed 2 (Table 1). In addition, SOM, TN, and TP did not differ between the monoculture and mixed plantations. The highest TK was measured in Mixed 1 RPs, followed by monoculture and Mixed 2 RPs. Seven variables and the abundance data of the bacterial and fungal dominant family were used for redundancy analysis. For bacteria, this variable combination explains 44.95% of the total variance of family abundance (Figure 4(a)). TN, TK, pH, and WC described 22.47%, 18.94%, 15.23%, and 15.18% of the total variance, respectively. Additional factors including Rubber

Forest, TP, and SOM on the bacterial community composition had relative little effect, accounting for 6.53%, 9.74%, and 11.19% of the variance, respectively. For fungi, this variables combination explained 27.77% of the total variance of the family abundance (Figure 4(b)). The pH and WC explained 15.14% and 8.96% of the total variance, respectively. Additional factors, that is, TN, Rubber Forest, TP, TK, and SOM for fungal community composition had little effect, describing only 5.50%, 5.49%, 3.61%, 3.15%, and 2.96% of the variance, respectively.

Discussion

Our results showed that the bacterial composition of the soil from the three RPs was basically similar. However, there was a significant difference in fungal communities among the three RPs at both the phylum and OTU level. The mixed RPs, that is, Mixed 1 and Mixed 2 RPs, were very similar in terms of phylum composition (ANOSIM results, $p = .65$) and OTU composition (ANOSIM results, $p = .07$). This indicated that intercropping other tree species into a RP has only a limited effect on soil bacterial composition. A previous study conducted in Hainan and Xishuangbanna showed that rubber forests and other tropical forests have different soil bacterial (Lan et al., 2017a; Lan, Li, Wu, & Xie, 2017b) and fungal compositions (Lan et al., 2017b). A new viewpoint has been drawn from our research, that is, monoculture and mixed RPs are similar in soil bacterial composition but different in fungal composition. Previous studies also showed mature rubber-based agroforestry systems can alleviate the adverse effects of rubber monoculture on bacterial biodiversity (Liu et al., 2019). Therefore, the effect of mixed RPs on the soil microbial community still needs long-term investigation.

In our investigation, the species richness, Chao, and Shannon diversity of the bacterial communities of the monoculture were higher than the Mixed 1 and Mixed 2 RPs, whereas all diversity indexes of the fungal communities were comparable between the monoculture and mixed RPs. These results showed that mixed RPs did not result in a higher diversity of bacteria. A study conducted in the Amazon showed that the local taxonomic and phylogenetic diversity of the soil bacteria was higher in agroecosystems compared to natural ecosystems (Rodrigues et al., 2013). A previous study observed that soil bacteria in a bamboo forest and a fir forest showed a prominent diversity, but with a lower total sequencing number than from native forest and mixed forest. A high variety of bacterial groups in monoculture rubber forests of agricultural practices would result in a considerable variety of soil microscopic organisms (Kerfahi et al., 2016; Lan et al., 2017, 2017a, 2017b). Previous studies have shown that the percentage of Gram-negative bacterial phospholipid fatty acids in jungle RPs was lower than in monoculture RPs

(Krashevskaya et al., 2015). Our results further confirmed that monoculture RPs have a higher bacterial diversity than mixed forests. In addition, our results also revealed that there were no significant differences in fungal diversity of soils between monoculture and mixed RPs. Our findings are similar to previous study conducted in the Jambi Province of southwest Sumatra, Indonesia (Krashevskaya et al., 2015), which showed that the total fungal phospholipid fatty acids of jungle RPs and monoculture RPs were comparable.

Our results showed that soil nutrition (such as TN and TK) and soil pH are the main drivers of the bacterial composition. However, soil pH and WC are the main drivers of the fungal composition. Our results are in line with a previous study in Danzhou, Hainan Island, which revealed that soil nutrients explained 43.05% of the total variance of soil bacterial composition in RPs (Lan et al., 2017a). In fact, numerous studies have shown that soil chemical properties, including soil pH, nitrogen, and carbon proportions together correspond with the bacterial community's composition and diversity (Lee, Barbier, Bottos, McDonald, & Cary, 2012; Smith, Barrett, Tusnady, Rejto, & Cary, 2010; Van Horn et al., 2013; Zeglin et al., 2011). Among the soil properties, pH is an imperative variable that impacts both soil bacterial communities and their alpha diversity (Fierer & Jackson, 2006). In our study, soil pH in the monoculture was relatively higher than in the mixed RP, which resulted in higher diversity among bacterial community in the monoculture than mixed RP. Very similar results were found in Hainan (Lan et al., 2018; Lan et al., 2017a) and Xishuangbanna (Lan et al., 2017b). Increased pH presumably is a key factor favoring nutrient capture in rubber production systems (Krashevskaya, Klarner, Widyastuti, Maraun, & Scheu, 2015). A previous study also confirmed that soil pH was the main driver of soil microbial composition in RPs (Krashevskaya et al., 2015).

Implications for Conservation

Our study revealed that there was a significant difference in fungal composition between monoculture and mixed plantations. The diversity of soil bacterial communities was higher in monoculture than in mixed RPs due to the higher soil pH in the monoculture. Our results also confirmed that soil pH is the main driver for both bacterial and fungal soil structure. From the point of view of soil microbial diversity, it seems that monoculture RPs are better than mixed RPs. However, from the perspective of plant diversity above ground, the mixed RP is better than the monoculture RP. Here, we suggest that more alkaline fertilizers should be applied in mixed RPs to improve the soil pH and soil microbial diversity. In this way, the total diversity would be enhanced to

some extent. At the same time, this treatment can also




Figure A1. Soil sampling design in three rubber plantations in Wenchang, Hainan Island.

increase the rubber latex yield.

Appendix

Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) disclosed receipt of the following financial support for the research, authorship, and/or publication of this article: This research work was financially sponsored by the National Natural Science Foundation of China (Grant No. 31770661), the Finance Science and Technology Project of Hainan Province (ZDYF2019145), the Lancang-Mekong River International Cooperation Project (081720203994192003), and the Earmarked Fund for China Agriculture Research System (CARS-34- ZP3).

ORCID iD

Muhammad Tahir Jatoi  <https://orcid.org/0000-0001-6191-3653>

Zhixiang Wu  <https://orcid.org/0000-0002-7344-8717>

References

- Balser, T., Kinzig, A., & Firestone, M. (2002). Linking soil microbial communities and ecosystem functioning. In A. Kinzig, S. Pacala, & D. Tilman (Eds.), *The functional consequences of biodiversity: Empirical progress and logical extensions* (pp. 265–293). Princeton, NJ: Princeton University Press.
- Cai, Z. Q., Zhang, Y. H., Yang, C., & Wang, S. (2018). Land-use type strongly shapes community composition, but not always diversity of soil microbes in tropical china. *Catena*, 165, 369–380.
- Chaudhary, A., Burivalova, Z., Koh, L. P., & Hellweg, S. (2016). Impact of forest management on species richness: Global meta-analysis and economic trade-offs. *Scientific Reports*, 6, 23954.
- Chen, L. C., Wang, H., Yu, X., Zhang, W. D., Lü, X. T., & Wang, S. L. (2017). The recovery time of soil carbon pools of conventional Chinese fir plantations from broadleaved forests in subtropical regions, China. *Science of the Total Environment*, 587, 296–304.
- Cole, L. E. S., Bhagwat, S. A., & Willis, K. J. (2014). Recovery and resilience of tropical forests after disturbance. *Nature Community*, 5, 3906.
- Ersine, P. D., Lamb, D., & Bristow, M. (2006). Tree species diversity and ecosystem function: Can tropical multi-species plantations generate greater productivity? *Forest Ecology Management*, 233, 205–210.
- Fierer, N., & Jackson, R. B. (2006). The diversity and biogeography of soil bacterial communities. *Proceedings of the National Academy of Sciences (USA)*, 103, 626–631.

- Francisco-Ortega, J., Wang, Z. S., Wang, F. G., Xing, F. W., Liu, H., Xu, H., . . . An, S. Q. (2010). Seed plant endemism on Hainan Island: A framework for conservation actions. *Botanical Review*, 76, 346–376.
- He, J. Z., Zheng, Y., Chen, C. R., He, Y. Q., & Zhang, L. M. (2008). Microbial composition and diversity of an upland red soil under long-term fertilization treatments as revealed by culture-dependent and culture-independent approaches. *Journal of Soils Sediments*, 8, 349–358.
- Kerfahi, D., Tripathi, B. M., Dong, K., Go, R., & Adams, J. M. (2016). Rainforest conversion to rubber plantation may not result in lower soil diversity of bacteria, fungi, and nematodes. *Microbial Ecology*, 72, 359–371.
- Kirk, J. L., Beaudette, L. A., Hart, M., Moutoglis, P., Klironomos, J. N., Lee, H., & Trevors, J. T. (2004). Methods of studying soil microbial diversity. *Journal of Microbiological Methods*, 58, 169–188.
- Koh, L. P., Miettinen, J., Liew, S. C., & Ghazoul, J. (2011). Remotely sensed evidence of tropical peatland conversion to oil palm. *Proceedings of the National Academy of Sciences (USA)*, 108, 5127–51232.
- Krashevskaya, V., Klarner, B., Widayastuti, R., Maraun, M., & Scheu, S. (2015). Impact of tropical lowland rainforest conversion into rubber and oil palm plantations on soil microbial communities. *Biology and Fertility of Soils*, 51(6), 697–705.
- Lamb, D., Erskine, P. D., & Parrotta, J. A. (2005). Restoration of degraded tropical forest landscapes. *Science*, 310, 1628–1632.
- Lan, G. Y., Li, Y. W., Jatoi, M. T., Tan, Z. H., Wu, Z. X., & Xie, G. S. (2017). Change in soil microbial community compositions and diversity following the conversion of tropical forest to rubber plantations in Xishuangbanna, Southwest China. *Tropical Conservation Science*, 10, 1–14.
- Lan, G. Y., Li, Y. W., Lesueur, D., Wu, Z. X., & Xie, G. S. (2018). Seasonal changes impact soil bacterial communities in a rubber plantation on Hainan Island, China. *Science of Total Environment*, 626, 826–834.
- Lan, G. Y., Li, Y. W., Wu, Z. X., & Xie, G. S. (2017a). Soil bacterial diversity impacted by conversion of secondary forest to rubber or eucalyptus plantations—A case study of Hainan Island, south China. *Forest Science*, 63, 87–93.
- Lan, G. Y., Li, Y. W., Wu, Z. X., & Xie, G. S. (2017b). Impact of tropical forest conversion on soil bacterial diversity in the tropical region of China. *European Journal of Soil Biology*, 83, 91–97.
- Lan, G. Y., Wang, J. K., Wu, Z. X., & Xie, G. S. (2013). Flora composition of seed plants in rubber forests in Hainan Island. *Journal of Northwest Forestry University*, 28, 37–41. (in Chinese)
- Langenheder, S., Bulling, M. T., Solan, M., & Prosser, J. I. (2010). Bacterial biodiversity-ecosystem functioning relations are modified by environmental complexity. *PLoS One*, 5, e10834.
- Lee, C. K., Barbier, B. A., Bottos, E. M., McDonald, I. R., & Cary, S. C. (2012). The intervalley soil comparative survey: The ecology of dry valley edaphic microbial communities. *The ISME Journal*, 6, 1046–1057.
- Li, Y. W., Xia, Y. J., Lei, Y. B., Deng, Y., Chen, H., Sha, L. Q., . . . Deng, X. B. (2015). Estimating changes in soil organic carbon storage due to land use changes using a modified calculation method. *iForest—Biogeosciences and Forestry*, 8, 45–52.
- Liu, C. G., Jin, Y. Q., Hu, Y. N., Tang, J. W., Xiong, Q. L., Xu, M. X., . . . Beng, K. C. (2019). Drivers of soil bacterial community structure and diversity in tropical agroforestry systems. *Agriculture Ecosystems & Environment*, 278, 24–34.
- Montagnini, F., Gonzales, E., & Porras, C. (1995). Mixed and pure forest plantations in the humid neotropics: A comparison of early growth, pest damage, and establishment cost. *Commonwealth Forestry Review*, 74, 306–314.
- Myers, N., Mittermeier, R. A., Mittermeier, C. G., Fonseca, G. A., & Kent, J. (2000). Biodiversity hotspots for conservation priorities. *Nature*, 403, 854–858.
- Naeem, S., Duffy, J. E., & Zavaleta, E. (2012). The functions of biological diversity in an age of extinction. *Science*, 336, 1401–1406.
- Rodrigues, J. L. M., Pellizari, V. H., Mueller, R., Baek, K., Jesus, E. D., Paula, F. S., . . . Nüsslein, K. (2013). Conversion of the Amazon rainforest to agriculture results in biotic homogenization of soil bacterial communities. *Proceedings of the National Academy of Sciences (USA)*, 110, 988–993.
- Rousk, J., Bååth, E., Brookes, P. C., Lauber, C. L., Lozupone, C., Caporaso, J. G., & Fierer, N. (2010). Soil bacterial and fungal communities across a pH gradient in an arable soil. *The ISME Journal*, 4, 1340–1351.
- Schimel, J. (1995). Ecosystem consequences of microbial diversity and community structure. In F. Staurt Chapin III & C. Körner (Eds.), *The Arctic and Alpine biodiversity: Patterns, causes and ecosystem consequences* (vol. 113, pp. 239–254). New York, NY: Springer.
- Schloss, P. D., Westcott, S. L., Ryabin, T., Hall, J. R., Hartmann, M., Hollister, E. B., . . . Weber, C. F. (2009). Introducing mothur: Open-source, platform-independent, community-supported software for describing and comparing microbial communities. *Applied and Environmental Microbiology*, 75, 7537–7541.
- Shen, C., Liang, W. J., Shi, Y., Lin, X. G., Zhang, H. Y., Wu, X., . . . Chu, H. (2014). Contrasting elevational diversity patterns between eukaryotic soil microbes and plants. *Ecology*, 95, 3190–3202.
- Singh, A. K., Rai, A., & Singh, N. (2016). Effect of long term land use systems on fractions of glomalin and soil organic carbon in the Indo-Gangetic plain. *Geoderma*, 277, 41–50.
- Singh, A. K., Rai, A., Banyal, R., Chauhan, S., & Singh, N. (2018). Plant community regulates soil multifunctionality in a tropical dry forest. *Ecological Indicators*, 95, 953–963.
- Smith, J. L., Barrett, J. E., Tusnady, G., Rejto, L., & Cary, S. C. (2010). Resolving environmental drivers of microbial community structure in Antarctic soils. *Antarctic Science*, 22, 673–680.
- Sodhi, N. S., Koh, L. P., Brook, B. W., & Ng, P. K. L. (2004). Southeast Asian biodiversity: An impending disaster. *Trends in Ecology and Evolution*, 19, 654–660.
- Soil Science Society of China. (2000). *Methods of soil agricultural chemical analysis*. Beijing: China Agricultural Science and Technology Press.
- Stackebrandt, E., & Goebel, B. M. (1994). Taxonomic note: A place for DNA-DNA reassociation and 16S rRNA sequence analysis in the present species definition in

- bacteriology. *International Journal of Systematic and Evolutionary Microbiology*, 44, 846–849.
- Van Horn, D. J., Van Horn, M. L., Barrett, J. E., Lee, M., John, V. H., . . . Takacs-Vesbach, C. D. (2013). Factors controlling soil microbial biomass and bacterial diversity and community composition in a cold desert ecosystem: The role of geographic scale. *PLoS One*, 8, e66103.
- Wang, J. K., Lan, G. Y., Wu, Z. X., & Xie, G. S. (2012). Investigation and analysis of plant resources under rubber forests in Hainan Island. *Chinese Journal of Tropical Agriculture*, 31, 31–36. (in Chinese)
- Wilcove, D. S., Giam, X., Edwards, D. P., Fisher, B., & Koh, L. P. (2013). Navjot's nightmare revisited: Logging, agriculture, and biodiversity in Southeast Asia. *Trends in Ecology and Evolution*, 28, 531–540.
- Wu, S., Zhu, Z., Fu, L., Niu, B., & Li, W. (2011). Web MGA: A customizable web server for fast metagenomic sequence analysis. *BMC Genomics*, 12, 444.
- Xu, N., Tan, G. C., Wang, H. Y., & Gai, X. P. (2016). Effect of biochar additions to soil on nitrogen leaching, microbial biomass, and bacterial community structure. *European Journal of Soil Biology*, 74, 1–8.
- Yan, W. B., Mahmood, Q., Peng, D. L., Fu, W. J., Chen, T., Wang, Y., . . . Liu, D. (2015). The spatial distribution pattern of heavy metals and risk assessment of most bamboo forest soil around lead-zinc mine in Southeastern China. *Soil Tillage Research*, 153, 120–130.
- Zeglin, L., Dahm, C., Barrett, J., Gooseff, M., Fitzpatrick, S., & Takacs-Vesbach, C. D. (2011). Bacterial community structure along moisture gradients in the para fluvial sediments of two ephemeral desert streams. *Microbial Ecology*, 61, 543–556.
- Zheng, Z., Cour, P., & Zhou, H. P. (2002). Modern pollen rain in Hainan Island, Southern China: Altitudinal pollen distribution in the tropical rain forest. *Acta Palaeontologica Sinica*, 41(4), 487–496.
- Zhu, H., & Zhou, H. X. (2002). A comparative study of the tropical rainforests of Xishuangbanna and Hainan. *Acta Botanica Yunnanica*, 24(1), 1–13.