

## **Climate and Emission Changes Contributing to Changes in Near-surface Ozone in Europe over the Coming Decades: Results from Model Studies**

Authors: Engardt, Magnuz, Bergström, Robert, and Andersson, Camilla

Source: AMBIO: A Journal of the Human Environment, 38(8) : 452-458

Published By: Royal Swedish Academy of Sciences

URL: <https://doi.org/10.1579/0044-7447-38.8.452>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

# Climate and Emission Changes Contributing to Changes in Near-surface Ozone in Europe over the Coming Decades: Results from Model Studies

We used an off-line, regional, model of atmospheric transport and chemistry to investigate current and future levels of near-surface ozone and accumulated ozone exposure over a threshold of 40 ppb(v) (AOT40) in Europe. To describe the current situation and enable an evaluation of the model's performance we simulated a number of years around 2000. To assess changes in ozone concentrations due to possible emission changes in Europe, the model was run with the meteorology of the early 2000s and precursor emissions from a set of Clean Air for Europe (CAFE) emissions scenarios. By extrapolation of the observed increase in near-surface O<sub>3</sub> at coastal locations in northwest Europe we constructed model boundaries that were used to simulate the impact of increasing hemispheric background in 2020. To assess changes in ozone concentrations due to climate change, the model was run with recent (2000) emissions but using meteorology from a regional climate model simulating a control (1961–1990) and a future (2021–2050) climate. The results indicate that climate change will have a small impact on ozone concentrations and AOT40 in the Nordic countries. Changes in hemispheric background concentrations and changes in precursor emissions in Europe will have a larger effect on ozone in Northern Europe. The situation is quite different in southern Europe, where climate change is expected to result in a very large increase in near-surface ozone concentrations.

## INTRODUCTION

Ozone (O<sub>3</sub>) constitutes a threat to human health (1) and is causing damage to vegetation (2) in regions with elevated concentrations. The present, near-surface, O<sub>3</sub> concentration in Europe is estimated to be at least double that in preindustrial times (3). In order to mitigate the consequences of elevated O<sub>3</sub> across Europe, international agreements (e.g., the Gothenburg Protocol) based on scientific evaluations of air pollutants' atmospheric cycle and their damage to humans and ecosystems are now in place aiming at reducing the emissions of O<sub>3</sub>-generating species.


O<sub>3</sub> is formed through photochemical reactions in environments rich in ozone precursors, i.e., nitrogen oxides (NO<sub>x</sub>), carbon monoxide (CO), and volatile organic compounds (VOCs). High temperature, intense solar radiation, stable atmosphere, and low wind speed—frequently occurring in southern Europe during summer—promote the production and accumulation of O<sub>3</sub> in the planetary boundary layer. Near-surface O<sub>3</sub> concentrations in northern Europe (the Nordic and Baltic countries) are generally much lower than in central and southern Europe as a result of the lower precursor concentrations here, and as a result of weather conditions less favorable for O<sub>3</sub> production at these latitudes. Episodes of enhanced near-surface O<sub>3</sub> in northern Europe typically occur

during spring and summer following the advection of O<sub>3</sub>-laden air from continental Europe after periods of O<sub>3</sub> buildup there.

The lifetime of boundary-layer O<sub>3</sub> is on the order of hours to days; changed precursor concentrations or weather conditions translate into O<sub>3</sub> changes within a few hours. This is evident when examining the diurnal variation of near-surface O<sub>3</sub>, which typically features maximum concentrations in the afternoon and low concentrations during night and early morning. O<sub>3</sub> is taken up at the surface and subsequently destroyed when it reacts with organic or other materials. At night, the photochemical production is turned off, while the vertical mixing—responsible for replenishing the O<sub>3</sub> continuously destroyed at the surface—is operating less efficiently than during daytime.

In this work we used the Multiple-scale Atmospheric Transport and Chemistry Model (MATCH)—a regional off-line, chemistry transport model (CTM)—to investigate current and future concentrations of O<sub>3</sub> in Europe. We investigated changes in near-surface O<sub>3</sub> brought about by altered precursor emissions in Europe, as well as increasing hemispheric O<sub>3</sub> and changes in climate following anthropogenic emissions of greenhouse gases and aerosol particles. We will restrict our discussion to accumulated ozone exposure over a threshold of 40 ppb (AOT40), which is one of the measures that is used in Europe to quantify damage to crops and natural vegetation brought about by near-surface O<sub>3</sub>. AOT40 is defined as the sum of the differences between the hourly mean O<sub>3</sub> concentration (in ppb[v]) and the threshold 40 ppb(v), when the concentration exceeds 40 ppb(v) during daytime. AOT40 crops (AOT40c) has traditionally been used as a measure of crop impacts in Europe and should be deduced at canopy height and accumulated over a 90-d growing season, which varies between different climatic regions. In this work, we set the growing season for crops to May–July and define daytime as between 06.00 and 18.00 coordinated universal time, across all of Europe. We aim at reproducing background measurements, which are taken at 2–10 m above ground and not specifically at canopy height. AOT40 forests (AOT40f) has been used as a measure of O<sub>3</sub> impacts on natural forests and is accumulated over the 6-mo period April through September. AOT40 will in this presentation be expressed in kppb(v)·h, i.e., =10<sup>3</sup> ppb(v)·h, sometimes denoted as ppm(v)·h.

The aim of this work is to study how O<sub>3</sub>, and AOT40 in particular, will develop in the Nordic and Baltic countries compared to the rest of Europe and to investigate the relative importance of the different factors controlling future O<sub>3</sub> in Europe. AOT40 is selected to illustrate the current and future impact of O<sub>3</sub> on vegetation. AOT40 is a concentration-based measure that we deduce from the modeled hourly O<sub>3</sub> concentrations (Fig. 1). The work can be regarded as a continuation and refinement of Laurila et al. (4), which used coarser-resolution models and older emission inventories to estimate changes in O<sub>3</sub> concentrations in the Nordic countries following global emission changes and regional climate change. The work also complements Simpson et al. (5), who use the



**Figure 1.** Layout of model system.

European Monitoring and Evaluation Program (EMEP) Meteorological Synthesizing Center—West (MSC-W) model to investigate the O<sub>3</sub>-induced impacts on vegetation in Europe in 2000 and in 2020 following one CAFE emission scenario as well as changed background concentrations of O<sub>3</sub> and methane.

## METHODS

For the mapping of current and future O<sub>3</sub> in Europe we use MATCH (6, 7)—an off-line CTM developed at the Swedish Meteorological and Hydrological Institute (SMHI). The model is flexible in that it accepts input from different sources and, depending on application, operates on different scales and generates different types of output, see Figure 1. Running the model over Europe is a standard task for MATCH, and the system's ability to calculate near-surface O<sub>3</sub> in the region is described and evaluated in a number of previous publications (8–10). The interested reader is directed to these papers for general descriptions of model setup and performance.

Year-to-year variation in meteorological conditions has a profound impact on near-surface O<sub>3</sub> in Europe (11, 12). Annual mean, near-surface, O<sub>3</sub> concentration for an arbitrary year typically deviates 5–10% from the long-term average, mainly as a consequence of the variability in the driving meteorology. In order to discriminate trends in O<sub>3</sub> brought about by changes in precursor emissions, background concentrations, or climate it is therefore essential to construct robust estimates of present and future concentrations by appropriate averaging of O<sub>3</sub> concentrations and selected indices (e.g., AOT40) over a number of years.

The present work is a compilation and extension of two previous (partly unpublished) studies where we examined the influence of decreasing emissions as well as increasing hemispheric background concentrations and the impact of climate change on European near-surface O<sub>3</sub> concentrations. The two studies used similar configurations of MATCH but operated with different input to produce hourly, three-dimensional fields of tropospheric O<sub>3</sub> across Europe. The first study dealt with the change in O<sub>3</sub> concentrations from the early 2000s to 2020 following changes in European precursor emissions and hemispheric background concentrations of O<sub>3</sub>; the model resolution was 0.4° × 0.4°, and the model atmosphere was divided into 22 layers up to 5 km. The second study (13) investigated the impact of one possible climate change

realization from 1961–1990 to 2020–2050 by driving MATCH with meteorological data representing recent and future climate; here the model resolution was 0.44° × 0.44° and the atmosphere was divided into 15 layers up to 6 km where the top boundary of the CTM was set. The geometries of the two model setups differ slightly and can be deduced from the maps below. Both studies utilized recent compilations of European emissions. The anthropogenic emissions were released with different temporal and spatial variation depending on sector. Biogenic isoprene emissions are calculated on-line in MATCH (7, 14). The experiments are summarized in Table 1 and further explained below.

## Model Evaluation

To evaluate the model's ability to simulate O<sub>3</sub> and AOT40 we ran MATCH with meteorology representing 4 y around the turn of the century (1999, 2000, 2001, 2003). The weather data were generated with the operational weather forecast model in use at SMHI (the high-resolution limited area model [HIRLAM]). Emissions of anthropogenic NO<sub>x</sub>, SO<sub>x</sub>, NH<sub>x</sub>, NMVOC, and CO were taken from EMEP's expert emissions (15), which reproduce the anthropogenic emissions occurring in Europe during these years. Monthly varying model boundary concentrations of all species of concern in the CTM were estimated from a combination of ground-based measurements and vertical soundings and should describe the conditions during each respective year. The results from this experiment constitute our best estimate of the situation during the early 2000s, which we will compare with measurements collected during the same years.

## Emission Reductions in Europe

To isolate the impact of changing European anthropogenic precursor emissions, we first ran MATCH using meteorology and boundary concentrations representing four different years (1999, 2000, 2001, 2003). In these calculations the model was fed with EMEP's expert emissions valid for one close-to-present year, 2004 (*BASE\_2004*). These simulations were contrasted with simulations where we ran MATCH with the same meteorological input and boundary values, but utilizing emission data from two different scenarios constructed by the International Institute for Applied Systems Analysis (IIASA) in the CAFE project.

The emissions in *D23A* are described in Amann et al. (16), where the scenario is denoted Case A, joint optimization. These gridded emissions constitute an optimized emission scenario where the aim was to reach a number of EU environmental targets (covering years of life lost due to particulate material smaller than 2.5 μm (PM<sub>2.5</sub>), acidification, eutrophication, and ozone) at the lowest possible cost. *D23A* is similar to the thematic strategy scenario in Amann et al. (17). MFR-DEEP stands for maximum technically feasible reduction plus a deep-cutting EU climate-taxation of 90 € per tonne CO<sub>2</sub> (18). This is currently believed to be close to what is possible to reach in

**Table 1.** Model experiments.

| Experiment      | Meteorology                | Emissions              | O <sub>3</sub> boundaries |
|-----------------|----------------------------|------------------------|---------------------------|
| Early 2000s     | 1999–2001, 2003 (HIRLAM)   | 1999–2001, 2003 (EMEP) | 1999–2001, 2003           |
| BASE_2004       | 1999–2001, 2003 (HIRLAM)   | 2004 (EMEP)            | 1999–2001, 2003           |
| D23A            | 1999–2001, 2003 (HIRLAM)   | D23A (IIASA/CAFE)      | 1999–2001, 2003           |
| MFR-DEEP        | 1999–2001, 2003 (HIRLAM)   | MFR-DEEP (IIASA/CAFE)  | 1999–2001, 2003           |
| 2020 Background | 1999–2001, 2003 (HIRLAM)   | D23A (IIASA/CAFE)      | 1999–2001, 2003 +4 ppb(v) |
| Control Climate | 1961–1990 (RCA3/ECHAM4 A2) | 2000 (EMEP)            | 2000                      |
| Future Climate  | 2021–2050 (RCA3/ECHAM4 A2) | 2000 (EMEP)            | 2000                      |

**Table 2. MATCH's ability to reproduce the early 2000s and control climate. Observations refers to data from EMEP monitoring sites (25). Model results are extracted from the grid cells encompassing the respective monitoring site.**

|                  | Observations<br>(kppb[v]-h) (period) | Model results<br>(kppb[v]-h) (period) | Correlation | Bias (%) | RMSE (kppb[v]-h) | No. samples |
|------------------|--------------------------------------|---------------------------------------|-------------|----------|------------------|-------------|
| Early 2000s*     |                                      |                                       |             |          |                  |             |
| AOT40c           | 7.0 (1999–2001, 2003)                | 7.8 (1999–2001, 2003)                 | 0.78        | +10.1    | 2.2              | 363         |
| AOT40f           | 12.4 (1999–2001, 2003)               | 13.4 (1999–2001, 2003)                | 0.83        | +8.0     | 3.5              | 355         |
| Control climate† |                                      |                                       |             |          |                  |             |
| AOT40c           | 7.7 (1997–2003)                      | 8.2 (1961–1990)                       | 0.86        | +6.3     | 2.8              | 60          |
| AOT40f           | 13.8 (1997–2003)                     | 13.3 (1961–1990)                      | 0.90        | –2.8     | 4.1              | 61          |

\* Model data reduced to 3 m aboveground. † Model data from lowest model layer (ca. 30 m).

Europe in 2020 and can therefore be regarded as the cleanest possible option.

### Increasing Hemispheric Background of O<sub>3</sub>

A number of authors (19, 20) have reported that background concentrations of O<sub>3</sub>, represented by data collected at remote locations around Europe, seem to have been increasing during the last few decades. The increase occurs in spite of recent emission reductions in North America and Europe and is thought to be caused by increasing emissions of O<sub>3</sub> precursors in the emerging economies of Asia, Africa, and Latin America, or increasing emissions from international shipping. Other factors, such as increased frequency of boreal forest fires, increased stratospheric-tropospheric exchange, or climate change, are also suggested. The increase is not uniform across all background stations, and the trend is not always clear. Measurements collected on western Ireland (21) indicate an average annual increase of  $\sim 0.25$  ppb(v) y<sup>–1</sup> for baseline air for the period 1990–2001. Derwent et al. (22) report a similar value ( $0.31 \pm 0.12$ ) ppb(v) y<sup>–1</sup> for the 20-y period 1987–2007. At background stations in southern and southwestern Sweden, annual mean ozone levels show upward trends of about  $\sim 0.24$  ppb(v) y<sup>–1</sup> for the period 1990–2006 (23).

To investigate the consequences of increasing hemispheric concentrations of O<sub>3</sub> on AOT40 in Europe during 2020 we executed the model with increased O<sub>3</sub> concentrations on the boundaries of the model domain (2020 background)—everything else (including the boundary concentration of other species) as the D23A simulation. We consider D23A to be a reasonably realistic emission scenario for Europe in 2020 and therefore use this experiment as the base for exploring the consequence of increasing hemispheric O<sub>3</sub> concentrations. The increased O<sub>3</sub> boundaries were deduced from an extrapolation of the  $0.25$  ppb(v) y<sup>–1</sup> trend of near-surface O<sub>3</sub>, which results in a 4 ppb(v) increase of O<sub>3</sub> from 2004 to 2020. The +4 ppb(v) was applied uniformly to all five boundaries (which all lie in the troposphere), during all 12 months.

### Climate Change

In the final analysis, we used long time series of meteorological data generated by a regional climate model, to drive MATCH. Control climate corresponds to the climatological average and variability during a recent, close to present, period. Future climate is constructed by running MATCH with an identical setup as the control climate simulation but using meteorology from a regional climate model simulating one possible climate change realization.

The driving meteorological data for the climate change experiments were generated by the Rossby Centre regional atmospheric climate model (RCA3) driven by boundary data from the global model ECHAM4/OPYC3 forced by the SRES

A2 emission scenario (24). The control climate and future climate both form 30-y averages representing roughly 1961–1990 and 2021–2050 in terms of climate but ca. 2000 in terms of emissions and boundary values. In this work we are interested in the change in AOT40 caused by climate change and will therefore consider the difference between these two 30-y averages.

The ability of the regional climate model to realistically simulate recent climate is discussed in Kjellström et al. (24), while the performance of MATCH forced by RCA3 data is evaluated in Andersson and Engardt (13). A separate evaluation of MATCH's ability to reproduce AOT40 measured at EMEP stations across Europe is discussed below along with the evaluation of the model's ability to reproduce the situation during the early 2000s.


## RESULTS

### Model Performance

Table 2 presents a summary of MATCH's capability of reproducing measured AOT40 during the early 2000s and the control climate. Since the data used for generating the statistical measures are not from the same locations and time periods, the two sets of results are not expected to be identical. The scores are also generated somewhat differently. For the early 2000s, the correlation coefficient (*r*), the bias (average relative model deviation), and the root mean-square error (RMSE) are calculated from the annual values at each station for each individual year. In the control climate these measures are formed by comparing the long-term averages (7 and 30 y, respectively) of observations and model results at each station. The latter method thus only takes the spatial variation into account, while the former also considers temporal variation between the years 1999, 2000, 2001, and 2003. For the statistics we only include those sites where the model and station elevation are within 250 m, and we use only stations that have at least 90% data coverage during the 3 and 6 months needed for the AOT40c and AOT40f calculations. The first criterion excludes a number of high-elevation sites, which MATCH cannot locate vertically because the model's limited horizontal resolution averages out the complex topography of the real world.

The observational data are typically collected at ca. 3 m above ground. In the climate change simulations, the statistical scores were marginally better if we used the average concentration in the lowest model layer instead of model values downscaled to 3 m, which is the normal quantity we use for “near-surface” O<sub>3</sub> in MATCH. The reason, we believe, is a consequence of the coarser vertical resolution in the meteorological data from the climate model (RCA3) compared to the weather forecast model (HIRLAM). Consequently, O<sub>3</sub> from the lowest model layer will be used to represent near-surface O<sub>3</sub> and





**Figure 2. Average modeled AOT40c over Europe, using 2004 emissions during the meteorological years 1999–2001, 2003. Units: kppb(v)-hours.**

for the calculations of AOT40c in the climate change simulations.


The correlation coefficient between observations and model result is close to or above 0.8, which means that MATCH can reproduce the spatial variability of AOT40 across Europe. The bias is within  $\pm 10\%$ , indicating that, on average, the model

realistically describes AOT40 values over Europe. The absolute error may, of course, be much larger at individual stations, which is also indicated by the relatively high RMSE. For the early 2000s the model slightly overestimates the values observed across Europe. For the climate simulations the 3-mo value is overestimated while the 6-mo value is underestimated. At first glance, it appears that the climate simulations perform better than the model's realization of the early 2000s. It is, however, more difficult to reproduce individual years than averages over a number of years, which the climate simulation scores are calculated for.


### Current Situation

Figure 2 shows average AOT40c over Europe using meteorology for 1999–2001 and 2003 and emissions valid for 2004 (*BASE\_2004*). According to the analysis above, the model, on average, overestimated measurements during the early 2000s by  $\sim 10\%$ . The spatial distribution, with high AOT40c in southern Europe and low values in the Nordic countries, the Baltic countries, and the British Isles was, however, well reproduced.

A recent compilation of dose-response relationships (26) established a linear relationship between growing season AOT40 and relative yield loss of several crops. For wheat, AOT40c of 3 kppb(v)-h corresponds to 5% yield loss, 6 kppb(v)-h cause 10% yield loss, etc. This means that substantial parts of Europe are currently at risk for severe ozone-related crop damages. New research has, however, pointed out that the AOT40 index is not optimal to assess crop losses and that stomatal uptake calculations are a far more realistic measure to assess the ozone-induced damages to vegetation (27). For example, during periods with reduced water availability (which is typical in southern Europe during summer), the plants will close their stomata to save water and, as a consequence, they will also be less affected by high  $O_3$  concentrations. Simpson et



**Figure 3. Change in AOT40c from 2004 to 2020 after changing the emission in Europe. (a) The average of the four D23A simulations minus the average of the four BASE\_2004 simulations; (b) The average of the four MFR-DEEP simulations minus the average of the four BASE\_2004 simulations. Units: kppb(v)-hours.**



**Figure 4.** Change in AOT40c after adjusting the background concentrations from the 2004 values to an extrapolated value representative of 2020. The figure depicts the average value of the meteorological years 1999–2001, 2003. Units: kppb(v)-hours.

al. (5) discuss the spatial differences in areas at risk using the AOT40 index compared to the accumulated stomatal flux. The former measure showing a much higher exceedence in southern Europe compared to the flux-based measure, which displays more uniform exceedences across Europe. In this setup of MATCH we did not explicitly calculate and store the stomatal uptake of  $O_3$  in different crops; our results are therefore depicting the estimated risk based on the accumulated dose of the concentrations of near-surface  $O_3$  only.

### Emission Reductions in Europe

Figure 3 shows the change in AOT40c from 2004 to 2020 when European precursor emissions evolve according to the emission scenarios D23A and MFR-DEEP, respectively. AOT40c will decrease substantially, all over Europe, in both scenarios. As expected from the definition and generation of the emission scenarios, the absolute decrease is largest where the highest AOT40c occur today (i.e., in southern and central Europe). The Nordic and Baltic countries will also experience significant decrease in AOT40c if Europe follows either of the two suggested emission paths. In the D23A scenario AOT40c is calculated to decrease 1–4 kppb(v)-h in southern Scandinavia, southern Finland, and the Baltic countries.

The difference between the two emission scenarios is substantial, meaning that strong commitments within Europe will substantially improve the air quality within Europe itself. Stronger commitments will clearly also have effects on AOT40c in Northern Europe, as evident from the difference between Figure 3a and Figure 3b.

### Increased Hemispheric $O_3$ Background

Increased  $O_3$  on the model boundaries, representing an elevated hemispheric burden of tropospheric  $O_3$  in 2020 compared to 2004, causes increased AOT40c throughout the model domain (Fig. 4). The increase is not uniform due to nonlinear effects in the chemistry and deposition of  $O_3$  and because AOT40c is a

cumulative index based on exceedence of a threshold (which is not always reached). Depending on the ambient concentrations of other species ( $NO_x$  in particular) different parts of the domain will react differently to increased  $O_3$ . A general feature (not shown) is that the additional  $O_3$  on the boundaries last longer over the oceans than over land due to the slow dry deposition of  $O_3$  to water surfaces.

Southern Sweden, Denmark, and parts of Norway, as well as southwestern Europe (Spain and Portugal) and most grid cells along the Atlantic coast will clearly be affected by the increasing hemispheric background. These land areas will experience an increase of AOT40c of 1–2 kppb(v)-h. While this amount in southern Europe is small compared to the expected decrease in AOT40c following emission changes up to 2020, 1–2 kppb(v)-h is typically what is expected to be gained in the Nordic and Baltic countries following emission changes in Europe over the same period (Fig. 3a,b).

### Climate Change

While most parts of southern and central Europe will experience a considerable increase in AOT40c due to climate change, the expected change in AOT40c from the late 20th century to the mid 21st century is small in the Nordic countries (Fig. 5a).


Andersson and Engardt (13) showed that  $O_3$  is likely to increase in southwest Europe due to decreased dry deposition of  $O_3$  in dryer climates and increased emissions of isoprene (which is an important biogenic  $O_3$  precursor) in a future climate. The decreased AOT40c over the Alps mountainous region and in the far north in Figure 5 is connected to the decreasing snow cover, which is radically increasing the dry deposition of  $O_3$ .

The climate change calculations cover a period of 60 y, while the emission change and boundary values calculations discussed above span ca. 16 y. In order to compare the climate effect with the other effects, we assume the climate change signal to be linearly increasing from 1961–1990 to 2021–2050 and present in Figure 5b the climate change signal from 2004 to 2020. Over this time period the effect of climate change is similar to the effect of increasing hemispheric background (+0.2–1 kppb(v)-h) over most parts of southern and western Europe. The values are smaller than the anticipated effect of the changing emissions but, together with the increasing hemispheric concentrations of  $O_3$ , they act to lessen the benefits gained by reducing  $O_3$  precursor emissions in Europe. Northern Europe is not expected to experience any significant changes in AOT40c following this climate change realization.

### DISCUSSION

In this work, we investigated the relative importance of emission changes in Europe, increased hemispheric background concentrations of  $O_3$ , and climate change on AOT40 in Europe.

To study emission changes in Europe, we used two different  $O_3$  precursor emission scenarios. One scenario describes a likely future situation; the second has been regarded as close to what is technically feasible. However, following the ambitious climate-mitigation commitments currently discussed in Europe, measures aimed at reducing European  $CO_2$  emissions, may, in fact, simultaneously reduce  $O_3$  precursor emissions below the MFR-DEEP scenario. Although there are many reports of increasing hemispheric background concentration of  $O_3$ , the magnitude of the change varies from location to location, which makes statements of future hemispheric background concentrations even more uncertain. The detailed change in climate in Europe over the coming years and its impact on near-surface  $O_3$  is also very uncertain. This is both because of the imperfection of the models and tools we use to predict future climate and because climate is inherently dependent on actions and decisions by



**Figure 5.** Change in AOT40c from control climate to future climate. (a) 1961–1990 to 2021–2050; (b) 2004 to 2020. Only grid cells with changes significant at the 99% level are shown. Units: kppb(v)-hours.

mankind during the coming decades. In our study we used one particular CTM operating on data from one regional climate model simulating one CO<sub>2</sub> emission scenario and forced by one particular global model on the boundaries. Using another CTM or climate model would give different details, but the overall effect of regional climate change on European O<sub>3</sub> in this study is similar to other recent studies (28–30).

## CONCLUSIONS

Although the results presented above are not strictly additive due to nonlinearities in O<sub>3</sub> chemistry, the following conclusions can be drawn:

Anticipated reductions in European O<sub>3</sub> precursor emissions will have a large beneficial impact on near-surface O<sub>3</sub> across Europe. The resulting reduction in AOT40c will be most pronounced in southern and central Europe—where AOT40c is currently at its highest, but the effect will also be clearly evident in the Nordic countries. Increasing hemispheric background concentrations of O<sub>3</sub> will, however, to a large extent offset the improvements in AOT40c in the Nordic countries over the coming 16 y.

Climate change has the potential to greatly increase O<sub>3</sub> concentrations in most of continental Europe. Given that the committed climate change is already significant and irrevocable, we regard the long-term effect of climate change a very serious impediment in reaching and retaining acceptable O<sub>3</sub> levels in continental Europe during the 21st century. Over the 16-y period studied in the present work, climate change, together with the increasing hemispheric background of O<sub>3</sub>, will to some extent counteract the gains in AOT40c achieved through reducing O<sub>3</sub> precursor emissions in Europe. In the Nordic countries climate change results in a small decrease of AOT40c, both over a 16-y perspective and in a 60-y perspective.

## References and Notes

- World Health Organization. 2003. *Health Aspects of Air Pollution with Particulate Matter, Ozone, and Nitrogen Dioxide*. EUR/03/5042688. Bonn, Germany, 94 pp.
- Emberson, L., Ashmore, M. and Murray, F. 2003. *Air Pollution Impacts on Crops and Forests. A Global Assessment*. Air Pollution Reviews, Vol. 4. Imperial College Press, London, 372 pp.
- Volz, A. and Kley, D. 1988. Evaluation of the Montsouris series of ozone measurements made in the nineteenth century. *Nature* 332, 240–242.
- Laurila, T., Jonson, J.E., Langner, J., Sundet, J., Tuovinen, J.-P., Bergström, R., Foltescu, V., Tarvainen, V., et al. 2004. *Ozone Exposure Scenarios in the Nordic Countries during the 21st Century*. EMEP/MSC-W Technical Report 2/2004. Norwegian Meteorological Institute, Oslo, 41 pp.
- Simpson, D., Ashmore, M.R., Emberson, L. and Tuovinen, J.-P. 2007. A comparison of two different approaches for mapping potential ozone damage to vegetation. A model study. *Environ. Poll.* 39, 715–725.
- Robertson, L., Langner, J. and Engardt, M. 1999. An Eulerian limited-area atmospheric transport model. *J. Appl. Meteor.* 38, 190–210.
- Langner, J., Bergström, R. and Pleijel, K. 1998. *European Scale Modeling of Sulfur, Oxidised Nitrogen and Photochemical Oxidants. Model Development and Evaluation for the 1994 Growing Season*. Swedish Meteorological and Hydrological Institute, RMK No. 82, 71 pp. (With errata).
- Timmes, S., Brandt, J., Flatøy, F., Bergström, R., Flemming, J., Langner, J., Christensen, J.H., Frohn, L.M., et al. 2002. Comparison of five Eulerian air pollution forecasting systems for the summer of 1999 using the German ozone monitoring data. *J. Atmos. Chem.* 42, 91–121.
- Vautard, R., van Loon, M., Schaap, M., Bergström, R., Bessagnet, B., Brandt, J., Buitjes, P.J.H., Christensen, J., et al. 2006. Is regional air quality model diversity representative of uncertainty for ozone simulation? *Geophys. Res. Lett.* 33, L24818. (doi: 10.1029/2006GL027610)
- van Loon, M., Vautard, R., Schaap, M., Bergström, R., Bessagnet, B., Brandt, J., Buitjes, P.J.H., Christensen, J., et al. 2007. Evaluation of long-term ozone simulations from seven regional air quality models and their ensemble. *Atmos. Environ.* 41, 2083–2097.
- Andersson, C. and Langner, J. 2007. Inter-annual variations of ozone and nitrogen dioxide over Europe during 1958–2003 simulated with a regional CTM. *Water Air Soil Pollut.: Focus* 7, 15–23. (doi: 10.1007/s11267-006-9088-4)
- Andersson, C., Langner, J. and Bergström, R. 2007. Interannual variation and trends in air pollution over Europe due to climate variability during 1958–2001 simulated with a regional CTM coupled to the ERA40 reanalysis. *Tellus* 59B, 77–98. (doi: 10.1111/j.1600-0889.2006.00196.x)
- Andersson, C. and Engardt, M. 2009. European ozone in a future climate—the importance of changes in dry deposition and isoprene emissions. *J. Geophys. Res.* (doi: 10.1029/2008JD011690) (In press).
- Simpson, D., Guenther, A., Hewitt, C.N. and Steinbrecher, R. 1995. Biogenic emissions in Europe I. Estimates and uncertainties. *J. Geophys. Res.* 100, 22875–22890.
- Vestreng, V. 2003. *Review and Revision. Emission Data Reported to CLRTAP*. MSC-W Status Report 2003, EMEP/MSC-W Note 1/2003. Norwegian Meteorological Institute, Oslo.
- Amann, M., Bertok, I., Cabala, R., Cofala, J., Heyes, C., Gyarmas, F., Klimont, Z., Schöpp, W., et al. 2005. *A Final Set of Scenarios for the Clean Air for Europe (CAFE) Programme—CAFE Scenario Analysis Report Nr. 6*. IIASA.
- Amann, M., Bertok, I., Cabala, R., Cofala, J., Heyes, C., Gyarmas, F., Klimont, Z., Schöpp, W., et al. 2005. *A Further Emission Control Scenario for the Clean Air for Europe (CAFE) Programme—CAFE Scenario Analysis Report Nr. 7*. IIASA.
- Amann, M., Cabala, R., Cofala, J., Heyes, C., Klimont, Z., Schöpp, W., Tarrason, L., Simpson, D., et al. 2004. *The “Current Legislation” and the “Maximum Technically Feasible Reduction” Cases for the CAFE Baseline Emission Projections, Version 2, November 2004—CAFE Scenario Analysis Report Nr. 2*. IIASA.
- Lelieveld, J., van Aardenne, J., Fischer, H., de Reus, M., Williams, J. and Winkler, P. 2004. Increasing ozone over the Atlantic Ocean. *Science* 304, 1483–1487.
- Simmonds, P., Derwent, R., Manning, A.L. and Spain, G. 2004. Significant growth in surface ozone at Mace Head, Ireland 1987–2003. *Atmos. Environ.* 38, 4769–4778.



21. Carslaw, D.C. 2005. On the changing seasonal cycles and trends of ozone at Mace Head, Ireland. *Atmos. Chem. Phys.* 5, 3441–3450.
22. Derwent, R.G., Simmonds, P.G., Manning, A.J. and Spain, T.G. 2007. Trends over a 20-year period from 1987 to 2007 in surface ozone at the atmospheric research station, Mace Head, Ireland. *Atmos. Environ.* 41, 9091–9098.
23. Lin Tang 2009. *Regional and Local Surface Ozone Variations in Relation to Meteorological Conditions in Sweden*. PhD thesis, Department of Earth Sciences, University of Gothenburg, Sweden.
24. Kjellström, E., Barring, L., Gollvik, S., Hansson, U., Jones, C., Samuelsson, P., Rummukainen, M., Ullerstig, A., et al. 2005. *A 140-year Simulation of the European Climate with the New Version of the Rossby Centre Regional Atmospheric Climate Model (RC43)*. SMHI report, RMK No. 108. Swedish Meteorological and Hydrological Institute, Norrköping, Sweden, 54 pp.
25. EMEP measurement data, extracted from <http://www.emep.int>, accessed 29 April 2008.
26. Mills, G., Buse, A., Gimeno, B., Bermejo, V., Holland, M., Emberson, L. and Pleijel, H. 2007. A synthesis of AOT40-based response functions and critical levels of ozone for agricultural and horticultural crops. *Atmos. Environ.* 41, 2630–2643. 2643. (doi: 10.1016/j.atmosenv.2006.11.016)
27. Pleijel, H., Danielsson, H., Emberson, L., Ashmore, M.R. and Mills, G. 2007. Ozone risk assessment for agricultural crops in Europe: further development of stomatal flux-response relationships for European wheat and potato. *Atmos. Environ.* 41, 3022–3040.
28. Langner, J., Bergström, R. and Foltescu, V. 2005. Impact of climate change on surface ozone and deposition of sulphur and nitrogen in Europe. *Atmos. Environ.* 39, 1129–1141.
29. Meleux, F., Solmon, F. and Giorgi, F. 2007. Increase in summer European ozone amounts due to climate change. *Atmos. Environ.* 41, 7577–7587.
30. Hedegaard, G.B., Brandt, J., Christensen, J.H., Frohn, L.M., Geels, C., Hansen, K.M. and Stendel, M. 2008. Impacts of climate change on air pollution levels in the Northern

Hemisphere with special focus on Europe and the Arctic. *Atmos. Chem. Phys.* 8, 3337–3367.

Magnuz Engardt is a researcher at the Swedish Meteorological and Hydrological Institute (SMHI). His address: SMHI, SE-601 76 Norrköping, Sweden.  
E-mail: [Magnuz.Engardt@smhi.se](mailto:Magnuz.Engardt@smhi.se)

Robert Bergström is a researcher at SMHI. He is also affiliated with the Department of Chemistry, University of Gothenburg. His address: SMHI, SE-601 76 Norrköping, Sweden.  
E-mail: [Robert.Bergstrom@smhi.se](mailto:Robert.Bergstrom@smhi.se)

Camilla Andersson is a researcher at the Swedish Meteorological and Hydrological Institute. Her address: SMHI, SE-601 76 Norrköping, Sweden.  
E-mail: [Camilla.Andersson@smhi.se](mailto:Camilla.Andersson@smhi.se)

# **UNITED STATES POSTAL SERVICE® (All Periodicals Publications Except Requester Publications)**

|  |   |  |
|--|---|--|
| 1. Publication Title<br>Ambio - A Journal of the Human Environment   | 2. Publication Number<br>0 0 8 - 1 0 2      | 3. Filing Date<br>Aug. 31, 2009  |
| 4. Issue Frequency<br>Feb., April, May, July, Aug., Oct., Nov. & Dec.  | 5. Number of Issues Published Annually<br>8 | 6. Annual Subscription Price<br>\$105.00   |
| 7. Complete Mailing Address of Known Office of Publication (Not printer) (Street, city, county, state, and ZIP+4®)<br>Ambio, 810 E. 10th Street, Lawrence, KS 66044-8897 |   | Contact Person<br>Daniela Bone<br>Telephone (include area code)<br>785-843-1234 ext. 242 |

8. Complete Mailing Address of Headquarters or General Business Office of Publisher (Not printer)  
Ambio, 810 E. 10th Street, Lawrence, KS 66044-8897

9. Full Names and Complete Mailing Addresses of Publisher, Editor, and Managing Editor (Do not leave blank)  
Publisher (Name and complete mailing address)  
Angle Fine, Allen Press, 810 E. 10th Street, Lawrence, KS 66044-8897

Editor (Name and complete mailing address)  
Elisabeth Kessler, Ambio, Royal Swedish Academy of Sciences, Box 50005, SE-10405 Stockholm, Sweden

Managing Editor (Name and complete mailing address)  
Beverly Lindeen, Allen Press, 810 E. 10th Street, Lawrence, KS 66044-8897

10. Owner (Do not leave blank. If the publication is owned by a corporation, give the name and address of the corporation immediately followed by the names and addresses of all stockholders owning or holding 1 percent or more of the total amount of stock. If not owned by a corporation, give the names and addresses of the individual owners. If owned by a partnership or other unincorporated firm, give its name and address as well as those of each individual owner. If the publication is published by a nonprofit organization, give its name and address.)

| Full Name                         | Complete Mailing Address              |
|-----------------------------------|---------------------------------------|
| Royal Swedish Academy of Sciences | Box 50005, SE-10405 Stockholm, Sweden |
|                                   |                                       |
|                                   |                                       |
|                                   |                                       |

11. Known Bondholders, Mortgagees, and Other Security Holders Owning or Holding 1 Percent or More of Total Amount of Bonds, Mortgages, or Other Securities. If none, check box ☒ None

| Full Name | Complete Mailing Address |
|-----------|--------------------------|
|           |                          |
|           |                          |
|           |                          |
|           |                          |

12. Tax Status (For completion by nonprofit organizations authorized to mail at nonprofit rates) (Check one)  
☒ The purpose, function, and nonprofit status of this organization and the exempt status for federal income tax purposes:  
☐ Has Not Changed During Preceding 12 Months  
☐ Has Changed During Preceding 12 Months (Publisher must submit explanation of change with this statement)

PS Form 3526, September 2007 (Page 1 of 3 (Instructions Page 3)) PSN 7530-01-000-9931 PRIVACY NOTICE: See our privacy policy on [www.usps.com](http://www.usps.com)

|   |  |  |   |
|---|--|--|---|
| 13. Publication Title<br>Ambio - A Journal of the Human Environment   |  | 14. Issue Date for Circulation Data Below<br>July 2009   |   |
| 15. Extent and Nature of Circulation  |  | Average No. Copies Each Issue During Preceding 12 Months | No. Copies of Single Issue Published Nearest to Filing Date |
| a. Total Number of Copies (Net press run)   |  | 2036   | 1800  |
| b. Paid Circulation (By Mail and Outside the Mail)  | (1) Mailed Outside-County Paid Subscriptions Stated on PS Form 3541 (include paid distribution above nominal rate, advertiser's proof copies, and exchange copies) | 247  | 187   |
|   | (2) Mailed In-County Paid Subscriptions Stated on PS Form 3541 (include paid distribution above nominal rate, advertiser's proof copies, and exchange copies)      | 0  | 0   |
|   | (3) Paid Distribution Outside the Mail: Including Sales Through Dealers and Carriers, Street Vendors, Counter Sales, and Other Paid Distribution Outside USPS®     | 153  | 158   |
|   | (4) Paid Distribution by Other Classes of Mail Through the USPS (e.g. First-Class Mail®)   | 1367   | 1266  |
| c. Total Paid Distribution (Sum of 15b (1), (2), (3), and (4))  |  | 1767   | 1631  |
| d. Free or Nominal Rate Distribution (By Mail and Outside the Mail)   | (1) Free or Nominal Rate Outside-County Copies Included on PS Form 3541  | 32   | 31  |
|   | (2) Free or Nominal Rate In-County Copies Included on PS Form 3541   | 0  | 0   |
|   | (3) Free or Nominal Rate Copies Mailed at Other Classes Through the USPS (e.g. First-Class Mail)   | 0  | 0   |
|   | (4) Free or Nominal Rate Distribution Outside the Mail (Carriers or other means)   | 3  | 0   |
| e. Total Free or Nominal Rate Distribution (Sum of 15d (1), (2), (3) and (4))   |  | 35   | 31  |
| f. Total Distribution (Sum of 15c and 15e)  |  | 1802   | 1662  |
| g. Copies not Distributed (See Instructions to Publishers #4 (page 83))   |  | 234  | 138   |
| h. Total (Sum of 15f and g)   |  | 2036   | 1800  |
| i. Percent Paid (15c divided by 15f times 100)  |  | 98.1   | 98.1  |
| 16. Publication of Statement of Ownership<br><input checked="" type="checkbox"/> If the publication is a general publication, publication of this statement is required. Will be printed in the November 2009 issue of this publication. <input type="checkbox"/> Publication not required. |  |  |   |
| 17. Signature and Title of Editor, Publisher, Business Manager, or Owner<br>Daniela Bone, Publishing Coordinator  |  |  | Date<br>8/21/09   |

I certify that all information furnished on this form is true and complete. I understand that anyone who furnishes false or misleading information on this form or who omits material or information requested on the form may be subject to criminal sanctions (including fines and imprisonment) and/or civil sanctions (including civil penalties).

PS Form 3526, September 2007 (Page 2 of 3)