

On Brazilian Ethanol and the Ecological Footprint

Authors: AZAR, CHRISTIAN, BERNDES, GÖRAN, HANSSON, JULIA,
and GRAHN, MARIA

Source: BioScience, 56(1) : 5

Published By: American Institute of Biological Sciences

URL: [https://doi.org/10.1641/0006-3568\(2006\)056\[0005:OBEATE\]2.0.CO;2](https://doi.org/10.1641/0006-3568(2006)056[0005:OBEATE]2.0.CO;2)

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

On Brazilian Ethanol and the Ecological Footprint

Marcelo Dias de Oliveira and colleagues assess the environmental impacts of ethanol production in Brazil and in the United States (*BioScience* 55: 593–602), and conclude that “the use of ethanol as a substitute for gasoline proved to be neither a sustainable nor an environmentally friendly option.”

Their paper offers interesting insights into ethanol production, but we are nevertheless in disagreement with their conclusion for Brazilian sugarcane-based ethanol. The reason for this is that their conclusion follows from a common but problematic use of the ecological footprint concept.

Dias de Oliveira and colleagues estimate the ecological footprint of burning gasoline as the forest area required for capturing the associated carbon dioxide (CO₂) emissions. For ethanol, the footprint is estimated as the amount of land used for the sugarcane production plus the forest area required for capturing the CO₂ emissions from energy input in the production (which is assumed to be fossil fuel-based). In total, the footprint for the ethanol car is approximately 0.56 hectare and that of a gasohol car (76 percent gasoline, 24 percent ethanol) about 0.63 hectare. Thus, according to their calculation of the ecological footprint, an ethanol car seems to be essentially equivalent to a gasohol car.

The major problem with their approach is that it fails to consider that the sequestration of CO₂ in a forest cannot continue forever. This means that when the established forest matures, the carbon sink ceases, a new area has to be planted, and the old area has to be managed so that the carbon stock remains intact.

Taking this dynamic feature into account yields a different picture. The footprint for ethanol remains constant, whereas the footprint for gasohol grows over time. This is illustrated in figure 1. Here we have assumed that bioenergy (instead of fossil fuels) is used in ethanol production, which gives a somewhat lower area requirement in this case.

Finally, although our analysis suggests that ethanol is preferable to gasoline from a dynamic footprint perspective, this does




Figure 1. The footprint of Brazilian ethanol and gasoline in a dynamic perspective.

not necessarily mean that it would be unproblematic if sugarcane ethanol became a dominant fuel in the transportation sector. If the 200 million cars in Europe ran on ethanol, they would require some 40 to 100 Mha of land (7 to 15 times the area used today for sugarcane in Brazil). Clearly, the scale of the potential demand is a cause for concern about potential negative environmental impacts. Other, complementary ways to solve the energy problem for transport are thus warranted.

CHRISTIAN AZAR
GÖRAN BERNDES
JULIA HANSSON
MARIA GRAHN

Christian Azar (e-mail: christian.azar@fy.chalmers.se) is a professor, Göran Berndes is an assistant professor, and Julia Hansson and Maria Grahn are graduate students at Chalmers University of Technology, 412 96 Göteborg, Sweden.

Response from Dias de Oliveira and Vaughan

Professor Azar and colleagues raise a number of interesting points. We agree that forest sequestration cannot continue indefinitely—we addressed the issue of uncertainty about forests’ potential to sequester CO₂ in our article—and we do not advocate it as a solution to the problem. Azar and colleagues note that the land area required to grow sugarcane or corn for 200 million cars fueled with ethanol is unrealistic. Indeed, that was one of the main points we tried to make, and it is true despite the imprecision involved in deter-

mining ecological footprint values, which we pointed out was problematic in some aspects. Nonetheless, the ecological footprint approach is still a valuable screening tool.

Regarding Azar and colleagues’ argument that the footprint of gasohol-fueled (but not ethanol-fueled) cars will grow over the years, it seems to us that the assimilation area needed per car would increase as well for ethanol. Consider the production of sugarcane: If only labor-intensive agricultural production (as in Brazil) is taken into account, the assimilation component with regard to ethanol would remain smaller. However, with any major scale-up of production, most countries would turn to industrial agriculture (e.g., supplementary nitrogen, pesticide, and fuel inputs) for biomass production, which would mean that the ecological footprint would increase.

Another important observation: After 30 years of the ethanol program, Brazilian production is still far from deserving the label of “clean energy.”

MARCELO E. DIAS DE OLIVEIRA
BURTON E. VAUGHAN

Marcelo E. Dias de Oliveira (e-mail: dias_oliveira@msn.com) was a graduate student at Washington State University (WSU) Tri-Cities, in Richland, Washington, when he and his coauthors wrote “Ethanol as Fuel: Energy, Carbon Dioxide Balances, and Ecological Footprint.” Burton E. Vaughan, a co-author of the article, is an adjunct professor of biological sciences at WSU Tri-Cities.

Letters to the Editor

BioScience
1444 I Street, NW, Suite 200
Washington, DC 20005
E-mail: bioscience@aibs.org

The staff of *BioScience* reserves the right to edit letters for clarity without notifying the author. Letters are published as space becomes available.