



## **Discerning the Environmental Drivers of Annual Migrations in an Endangered Amphibian**

Authors: Brooks, George C., Smith, Jennifer A., Gorman, Thomas A., and Haas, Carola A.

Source: Copeia, 107(2) : 270-276

Published By: The American Society of Ichthyologists and Herpetologists

URL: <https://doi.org/10.1643/CH-18-068>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

# Discerning the Environmental Drivers of Annual Migrations in an Endangered Amphibian

George C. Brooks<sup>1</sup>, Jennifer A. Smith<sup>1,2</sup>, Thomas A. Gorman<sup>1,3</sup>, and Carola A. Haas<sup>1</sup>

**The prevalence of studies that discern the drivers of animal migrations is increasing exponentially, in line with raised concerns over climate-driven shifts in phenological patterns. Amphibians that breed in ephemeral wetlands are particularly vulnerable owing to the stochastic nature of their breeding habitat and high survival and reproductive costs associated with untimely migrations. Knowledge of the effects of climatic cues on the timing of amphibian migration and breeding phenologies is required to understand the potential consequences of climate change and to inform conservation plans for at-risk species. Here we use six years of count data from Eglin Air Force Base, Florida, U.S., collected between 2010 and 2016, to model the phenology of migrations of *Ambystoma bishopi*. Models were fit with a negative binomial error distribution, with daily values of temperature and precipitation as covariates. Models were compared in an AIC framework. We found that salamander movement was strongly tied to environmental cues. A proclivity to move over a medial range of temperatures and precipitation resulted in a ‘Goldilocks’ set of conditions that facilitate migration. We also discovered a role for endogenous drivers of movement, with day of year predicting above-ground activity regardless of environmental conditions. These results contribute to our basic knowledge of the behavior of *A. bishopi* and provide managers with useful metrics to maximize the efficacy of conservation actions. More broadly, the methods employed within have utility to amphibian researchers to help predict the effects of climate change and tailor management strategies accordingly.**

ENVIRONMENTAL factors that determine phenological patterns in nature have received increasing attention, partly driven by increasing concerns of climate change and concomitant phenological mismatches (Roy and Sparks, 2000; Menzel et al., 2006; Kelly et al., 2016; Green, 2017). Climate-driven shifts in the timing of key life-history events are predicted to have consequences at the species and community level (Edwards and Richardson, 2004; Thackeray et al., 2010; Valtonen et al., 2011). For example, asynchronous shifts in phenologies across taxa may disrupt plant–pollinator mutualisms (Hegland et al., 2009), decouple latitudinal migrations from resource availability (Both and Visser, 2001; Post and Forchhammer, 2008), and lead to mismatches between peak food availability and reproduction (Visser et al., 1998, 2006; Plard et al., 2014). Developing a better understanding of the environmental cues that underlie phenologies is fundamental for predicting the responses of species to a changing environment, and thus development of steps toward mitigating the effects of climate change.

Many amphibians in temperate regions undertake annual migrations from terrestrial upland habitat to aquatic breeding habitats (Semlitsch, 2008). In the southeastern United States (U.S.), many amphibians rely on ephemeral (i.e., temporary) wetlands for reproduction and larval development (Russell et al., 2002; Gibbons et al., 2006). Adults must use environmental cues to time movements to coincide with seasonal precipitation events that fill wetlands in order to allow maximum development time for larvae before water levels recede. In addition, adults must time above-ground movements with environmental conditions that mitigate the risk of desiccation. Accordingly, amphibian migrations to and from breeding wetlands have been linked to climatic variables such as daily rainfall and temperature (Jakob et al.,

2003; Cook et al., 2006; Todd and Winne, 2006; Timm et al., 2007; Canavero et al., 2008).

Climate projections for the southeastern U.S. suggest warmer mean temperatures, shifts in the frequency and timing of rainfall events, and longer periods of drought (Strzepek et al., 2010; Bucklin et al., 2013; Ingram et al., 2013; Chandler et al., 2016). Clearly such environmental change is pertinent to amphibian ecology and future management of threatened and endangered species (Beebee, 1995; Walls et al., 2013; Chandler et al., 2016; Green, 2017). To direct conservation efforts, additional long-term studies of the environmental cues underlying amphibian movements are needed, especially given the observed declines of many species (Stuart et al., 2004; Alford, 2010).

Here, we evaluate linkages between environmental variables and the timing of breeding movements between upland habitat and ephemeral wetlands of Reticulated Flatwoods Salamanders (*Ambystoma bishopi*; hereafter flatwoods salamanders). Flatwoods salamanders are considered federally endangered under the U.S. Endangered Species Act (USFWS, 2009) and vulnerable by the IUCN (Palis and Hammerson, 2008). Loss and alteration of preferred habitat are often implicated in the decline of populations (Means et al., 1996; USFWS, 1999, 2009). Climate change may also have ramifications for the persistence of flatwoods salamander populations if it leads to shifts in the timing of above-ground movements that underlie breeding phenologies or shortened hydroperiods of breeding wetlands. Understanding the potential consequences of climate change requires knowledge of the effects of climatic cues on phenological patterns.

Flatwoods salamanders inhabit longleaf pine (*Pinus palustris*) flatwoods and savannas in the southeastern Coastal Plain in the U.S. Adults are fossorial and occupy mesic upland habitats, but migrate to ephemeral wetlands with

<sup>1</sup> Department of Fish and Wildlife Conservation, Virginia Tech, 310 West Campus Drive, Blacksburg, Virginia 24061; Email: (GCB) boa10gb@vt.edu; and (CAH) cahaas@vt.edu. Send reprint requests to GCB.

<sup>2</sup> Present address: Department of Environmental Science and Ecology, The University of Texas at San Antonio, One UTSA Circle, San Antonio, Texas 78249; Email: jennifer.smith@utsa.edu.

<sup>3</sup> Present address: Aquatic Resources Division, Washington State Department of Natural Resources, PO BOX 280, Castle Rock, Washington 98611; Email: gormant@vt.edu.

Submitted: 30 May 2018. Accepted: 25 March 2019. Associate Editor: J. Kerby.

© 2019 by the American Society of Ichthyologists and Herpetologists DOI: 10.1643/CH-18-068 Published online: 16 May 2019

well-developed herbaceous ground cover and relatively open canopy to breed (Palis, 1997; Gorman et al., 2009, 2014). Unlike most other North American amphibians, flatwoods salamanders lay their eggs in dry wetland basins before they fill, allowing embryos to develop so that eggs can hatch immediately when wetlands are inundated a few weeks, or months, after laying (Anderson and Williamson, 1976; Palis, 1997; Gorman et al., 2014). Although anecdotally flatwoods salamanders migrate following the first cool rains of fall, direct quantification of the drivers of movement are lacking. Previous studies of flatwoods salamander movements have been purely descriptive (Erwin et al., 2016), conducted over relatively short time periods (Palis, 1997), or have only considered crude measures of climatic variables (e.g., seasonal rainfall; Palis et al., 2006). Empirical assessment of the cues driving flatwoods salamander migrations is needed in order to inform conservation plans for this imperiled species. Being able to predict amphibian movement has additional utility, for scheduling research activities, protection efforts at road-crossings, or potentially disruptive activities in the uplands surrounding breeding sites.

## MATERIALS AND METHODS

**Study site.**—Our study was conducted on Eglin Air Force Base (hereafter Eglin), Okaloosa County, Florida, U.S. from 2010 to 2016. Eglin is a large military installation (188,459 ha), primarily consisting of actively managed Longleaf Pine dominated sandhills (approximately 145,000 ha) interspersed with treeless open test ranges, pine plantations, and mesic flatwoods. The landscape is punctuated by ephemeral wetlands consisting of shallow depressions of low topography, which fill in the fall or early winter and typically remain inundated throughout the spring (Chandler et al., 2016). We focused on two wetlands, approximately 500 m apart, which have been monitored intensively since 2009 and that are known to be occupied by flatwoods salamanders. Both wetlands are approximately 0.4 ha and are characterized by open overstories dominated by Longleaf Pine, Slash Pine (*P. elliotii*), and Pond Cypress (*Taxodium ascendens*), and mid-stories dominated by Myrtle Dahoon (*Ilex cassine* var. *myrtifolia*), Chapman's St. John's-wort (*Hypericum chapmanii*), and Swamp Titi (*Cyrilla racemiflora*). Wetland ecotones and portions of basins used for breeding are dominated by grasses, including *Aristida stricta* and bluestems (*Andropogon* spp.) and forbs, including Flattened Pipewort (*Eriocaulon compressum*).

**Sampling design.**—To monitor the movements of flatwood salamanders, both wetlands were completely encircled with drift fences and funnel traps (see Erwin et al., 2016 for details). The drift fences were constructed from 60 cm tall metal flashing buried in the sediment approximately 15–20 cm. Funnel traps measuring 85 cm x 20 cm were placed flush with the fence and ground at approximately 10 m intervals on both sides of the fence. We began drift fence operations on or before the first major rains were forecast in fall, typically late October or early November. Once we initiated drift fence operations, we checked traps multiple times per night through March if water levels receded and there were no larvae present, or May if the wetlands continued to hold water. In some years, we ceased operations for a week in late December and some weekends due to staffing shortages. Our data comprise 892 sampling nights spanning six breeding seasons from fall 2010 to spring 2016; specific dates of active

sampling are reported in Supplemental Table A (see Data Accessibility). Upon capture, we recorded the date and time of capture, and uniquely marked each individual using passive integrated transponder (PIT) tags or visual implant elastomer (VIE). Animals were sexed if possible, and released immediately after marking on the opposite side of the fence from their capture location. Between sampling events, we closed traps and removed sections of the fence panels to allow free movement of animals to and from wetlands.

**Environmental variables.**—Rain gauges were not installed for all but the last two years of our study, and so we used PRISM data at the highest resolution (800 m) in lieu of more precise information. For the two years in which rain gauge data were available, precipitation was highly correlated between the two datasets. We collected daily precipitation and temperature data from the PRISM Climate Group for the years 2010 to 2016 (Oregon State University, <http://prism.oregonstate.edu>, created 1 May 2017). The PRISM data set is derived from climatologically aided interpolation that averages across multiple stations and adjusts values for the grid reference of interest using output from a digital elevation model.

**Statistical analyses.**—To elucidate patterns in the timing of above-ground movements and the environmental factors that trigger migration, we fitted several competing generalized linear models (GLMs). Daily counts were modeled with daily precipitation, minimum daily temperature, and day of year as predictors. We considered both linear and quadratic functions of all predictors to retain flexibility in the relationships with the response. Year was included as a fixed effect, and an interaction between temperature and rainfall was explored. A high proportion of individuals of unknown sex prevented sex from being included as a random effect. Emerging metamorphs were excluded from the analysis. We evaluated collinearity between predictor variables using Spearman's rank correlation and omitted one of a pair of variables when  $|r| < 0.5$  to minimize challenges associated with multi-collinearity (Dormann et al., 2013). We used Durbin-Watson statistics on model residuals to check for serial autocorrelation.

Preliminary analyses revealed that the negative-binomial distribution (NB) outperformed Poisson, zero-inflated Poisson, and Poisson-log normal error distributions. The NB parameterization can account for overdispersion commonly seen in biological data (Bliss and Fisher, 1953; Ver Hoef and Boveng, 2007; Lindén and Mäntyniemi, 2011). Failure to account for overdispersion in count data could lead to erroneous conclusions regarding the effect of covariates and ultimately lead to misguided conservation actions (Terceiro, 2003; Ver Hoef and Boveng, 2007). All results reported assume a NB error distribution, and all models were fitted in R using the statistical package 'MASS' using the `glm.nb` function (R Core Team, 2015).

All possible combination of variables were fit to the data, and models were compared in an AIC framework. Model selection was based on AIC weights and minimizing uninformative parameters (Arnold, 2010). Cross correlation analysis was implemented to reveal any lag effects between predictor and response, and collinearity between predictors was tested through inspection of variance inflation factors. For individuals of known sex, comparison between dates of movement for males and females were made with Wilcoxon rank-sum tests.

**Table 1.** Model comparison of the top five performing models based on AIC values and Akaike weights ( $w_i$ ). The null model is included for reference. All models presented assume a negative binomial error distribution. Variables included are daily precipitation (ppt), minimum daily temperature ( $t_{\min}$ ), day of year, and year. Quadratic terms include all lower order polynomials (e.g.,  $\text{Day}^2 = \text{Day} + \text{Day}^2$ ).

Model	AIC	$\Delta\text{AIC}$	$w_i$
$\sim \text{ppt}^2 + t_{\min}^2 + \text{Day} + \text{Year}$	1634.3	0.0	0.33
$\sim \text{ppt}^2 + t_{\min}^2 + \text{Day} + \text{Year} + (\text{ppt} * t_{\min})$	1634.9	0.5	0.25
$\sim \text{ppt}^2 + t_{\min}^2 + \text{Day}^2 + \text{Year}$	1635.5	1.2	0.18
$\sim \text{ppt}^2 + t_{\min}^2 + \text{Day}^2 + \text{Year} + (\text{ppt} * t_{\min})$	1636.2	1.9	0.12
$\sim \text{ppt}^2 + t_{\min} + \text{Day} + \text{Year} + (\text{ppt} * t_{\min})$	1638.0	3.7	0.05
Null	1780.0	145.7	0.00


## RESULTS

Over the duration of the study, flatwoods salamanders were detected on 179 separate sampling nights ( $179/892 = 20\%$ ) yielding 1025 total captures (Year 1 = 164, 2 = 231, 3 = 128, 4 = 118, 5 = 79, 6 = 305). During the study period, minimum daily temperatures averaged  $9.1^\circ\text{C}$  (range:  $-8$ – $24.4^\circ\text{C}$ ), and precipitation averaged 5.5 mm (range: 0–139.9 mm). Of the nights that salamanders were captured, minimum temperatures averaged  $11.7^\circ\text{C}$  (range:  $-1.5$ – $22.5^\circ\text{C}$ ), and precipitation averaged 7.9 mm (range: 0–92.7 mm). Year-specific demographic and environmental information is summarized in Supplemental Table B (see Data Accessibility).

The top five performing models and corresponding AIC values are presented in Table 1. The top four models all have sufficient AIC weights to warrant consideration; however, the top model reduces the number of uninformative parameters and represents the most parsimonious description of the data. The top model includes quadratic terms for precipitation and temperature. Initially, we considered mean, minimum, and maximum daily temperatures, but subsequently excluded mean and maximum temperatures owing to the high degree of correlation between all three variables.

Both temperature and precipitation exhibited nonlinear relationships with salamander movement. The steepness of the contours, across both the precipitation and temperature axes, suggests distinct thresholds in environmental conditions that trigger salamander movement (Supplemental Figure A; see Data Accessibility). Intermediate values of temperature and precipitation were correlated with the highest movements, with a monotypic peak in movement occurring at  $14^\circ\text{C}$  and 43 mm of rainfall. The relationships between climatic variables and movement were qualitatively similar across all model parameterizations. There was further evidence for a lagged effect of minimum daily temperatures on salamander movement; overland migrations were more strongly correlated with temperature in previous time steps (results not shown). In contrast, the effect of precipitation on salamander movement showed no evidence of a time lag. The second best performing model also contained an interaction between temperature and precipitation. At low temperatures, increased precipitation was predicted to have a larger positive effect on salamander movement compared with an equivalent increase at high temperatures.

Irrespective of environmental conditions, day of year influenced salamander activity; movements were concentrated around October, representing peak arrival dates, with above-ground movement declining as the season progressed. Year was retained in the top model parameterization,



**Fig. 1.** Daily counts of individuals as a proportion of the total number of individuals available for capture for each year. Major tick marks are set at January 1, and minor tick marks are months.

indicative of annual variation in movement patterns (Fig. 1). In three of the six years, over 20% of the breeding population migrated to wetlands on a single night.

Although we found no overall difference between the timing of movements of males and females (Wilcoxon rank-sum test,  $P = 0.84$ ), there were noticeable differences within years (Fig. 2). In fall of 2011 and 2015, males arrived earlier than females, whilst in fall of 2013, females arrived earlier than males (Fig. 2). In all other breeding seasons, there was no difference in arrival times between the sexes.


Predicted frequencies from the top-performing model provided a good fit to the observed data (Chi-Sq = 13,  $P = 0.2$ ). Predictive accuracy is high for peak movement nights ( $>40$  individuals) and daily counts below 15 individuals; however, the model tends to over-predict the frequency of nights with 25–35 individuals and under-predict nights with no captures. Comparison of the observed frequency distribution of arrivals with those expected under the top model are reported in Supplemental Figure B (see Data Accessibility).

Serial autocorrelation is a common concern in studies of count data, particularly when movement occurs in clusters; however, we found no evidence of correlation between the model residuals and thus serial autocorrelation does not appear to be an issue in our data (Durbin-Watson statistic = 2.3,  $P = 0.9$ ).

## DISCUSSION

In this study, we evaluated the associations between temperature and rainfall, and the movement of flatwoods salamanders in and out of their breeding wetlands. We see movements concentrated in October on rainy nights over a specific range of ambient temperatures. Drawing from multiple, interacting environmental cues in this fashion may be an adaptive trait to avoid moving during heavy summer rains frequent in the southeastern US. Alternatively, salamanders may concentrate movements on nights with low temperatures when snake predators may be less active and/or when energetic stressors are lower. We were unable to disentangle the effects of absolute temperatures from relative changes in temperature. The importance of changing temperatures, as opposed to temperature per se, has recently been demonstrated in avian nesting behavior (Schaper et al.,





**Fig. 2.** Comparison of arrival dates for male and female flatwoods salamanders. Plots are standardized from 1 October and separated by year, i.e., 10–11 is the fall 2010–spring 2011 breeding season. Within year comparisons are made with Wilcoxon rank-sum tests.

2012); we predict similar mechanisms operate in many amphibian phenologies and deserve further study.

We also found a significant effect of date on salamander movements. Date can serve as a proxy for photoperiod or reflect any number of endogenous cues decoupled from environmental conditions (Both et al., 2008; Canavero et al., 2008). Whilst such endogenous cues to migration behavior are commonly acknowledged in mammals and birds (Knudsen et al., 2011; Williams et al., 2014), they are not as well documented in analogous amphibian research (Bradshaw and Holzapfel, 2007; but see Grant et al., 2009 and references therein). The negative relationship between date and daily counts stems from the synchronous arrivals of individuals at the start of the breeding season, contrasting with the staggered, protracted departure of individuals subsequent to breeding. Although not categorized as an explosive breeder (Petranka, 1998), pulse movement events comprising a large proportion of the total population are still evident; half of all captures occurred on just 24 sampling nights.

Protandry, that is, the arrival of males before females at breeding sites, has been documented in several species within Ambystomatidae (Douglas, 1979; Semlitsch, 1985; Sexton et al., 1990; Semlitsch et al., 1993; Trenham et al., 2000). It has been suggested that males arriving early maximize the number of potential mating opportunities, especially when female arrivals are unpredictable. In contrast, females that defer arrival at breeding sites maximize mate choice (Morbey and Ydenberg, 2001). Our results provide evidence of protandry in flatwoods salamanders. However, males only preceded females in two years of our study. For the remaining years, synchrony in arrival times of male and females occurred in the 2010–2011, 2012–2013, and 2014–2015 seasons, corroborating the observations of Palis (1997), and females arrived earlier than males in the 2013–2014 season.

It has been suggested that unpredictable resources, and the high probability of reproductive failure, selects for synchronous arrivals in ephemeral pool breeders (Dodd, 1993; Hartel

et al., 2007). This is certainly an apt description of flatwoods salamander ecology, with only two successful recruitment years over the course of this study. Alternatively, synchronization can be attributed to sperm competition or other sexual behaviors, as shown in *Triturus* spp. (Hartel et al., 2007). Neither of these mechanisms, however, can explain variability across years in the timing of movement when comparing males and females. To the authors' knowledge, this is the first documentation of such inter-annual variability in arrival times for any pond-breeding amphibian. It is possible that in dry summers, some adults remain in the breeding ponds so are not detected immigrating in the following season. Additionally, after multiple years of reproductive failure, the adult population declines, and differential mortality by sex or small sample size may contribute to the difficulty in discerning patterns. More research on the relationship between environmental stochasticity and synchrony is needed.

A better understanding of the environmental factors driving migration is important for conservation and management of species of amphibians that make seasonal movements. Such studies also contribute to our basic knowledge of amphibian behavior and phenotypic constraints. Amphibian populations can suffer heavy mortality when they cross roads to move between their terrestrial and breeding habitats, and several mitigation strategies (such as temporary road closures or citizen volunteer “bucket-brigade” transport) depend on an ability to predict timing of amphibian movement (Schmidt and Zumbach, 2008; Beebe, 2013). In addition, evaluating whether a particular wetland has a suitable hydroperiod for a species of concern, or determining the temporal niches of different species of amphibians breeding in the same wetland, requires information on the timing of arrival and departure (Paton and Crouch, 2002).

Amphibian conservation operates on a relative shoestring budget, and intensive long-term monitoring is often unsustainable (Beebe, 2014; Walls et al., 2017; Meredith et al.,

2018). Future sampling efforts should be tailored to maximize capture rates with the minimum cost. The negative binomial model provided useful threshold limits for precipitation and temperature beyond which salamander movement is likely, and identified a 'Goldilocks' set of environmental conditions that flatwoods salamanders favor. No captures were made on nights where temperature exceeded 22°C or fell below -1.5°C. Similarly, nights with more than ten captures did not occur before 25 October or after 25 February. This kind of information can be used to inform managers as to when surveys should be conducted and when monitoring can be curtailed with negligible loss of data. More broadly, the analytical tools described above can be employed in any study of pond-breeding amphibians to develop a more cost-effective monitoring schedule.

The strong association between climatic variables and salamander movement foreshadows future challenges in the face of climate change for this endangered species. Rising temperatures may delay arrival times at breeding wetlands (as shown for two species of salamander by Todd et al., 2011), thereby reducing the amount of time available for larval development. With climate projections forecasting more sporadic rainfall events over the species range (Mulholland et al., 1997; Chandler et al., 2016), the timing of salamander movements will likely become decoupled from environmental conditions conducive to successful reproduction; given enough change, those conditions may cease to exist entirely. The impacts of this phenological mismatch are predicted to be extremely detrimental to long-term population viability (Jakob et al., 2003; Chandler et al., 2016). Indeed, the annual variability in protandry seen in these populations may reflect maladaptive consequences of environmental perturbations that have already occurred.

Amphibians are often considered amongst vertebrates the most at risk to climate change (Alford, 2010). To date, the relative contributions of climatic cues and circannual rhythms to amphibian phenologies has received little attention, yet in order to confidently predict the response of amphibians to future climate scenarios, the degree of plasticity in phenological traits needs to be addressed. Experimental work on the proximate drivers of movement would help to this end. To combat the bleak prospects that amphibians face, the methods described herein allow researchers to exercise thrift, both in terms of sampling design and survey effort, and shrewdness in their mitigation strategies.

#### DATA ACCESSIBILITY

Supplemental information is available at <https://www.copeiajournal.org/ch-18-068>.

#### ACKNOWLEDGMENTS

We would like to thank the team of people who have made this research possible. Special mention should be given to K. Jones, B. Rincon, S. Goodman, V. Porter, and the myriad seasonal technicians involved in data collection. We thank E. Frimpong for comments that greatly improved the manuscript. We thank the Natural Resources Branch of Eglin Air Force Base (Jackson Guard), the U.S. Fish and Wildlife Service, the Department of Defense Legacy Resource Management Program, the Florida Fish and Wildlife Conservation Commission's Aquatic Habitat Restoration and Enhancement Program, and the Department of Fish and Wildlife Conser-

vation at Virginia Tech for funding and logistical support on this project. This work was supported by the USDA National Institute of Food and Agriculture, McIntire Stennis project 1006328.

#### LITERATURE CITED

- Alford, R. A. 2010. Declines and the global status of amphibians, p. 13–45. *In*: Ecotoxicology of Amphibians and Reptiles. D. Sparling, G. Linder, C. A. Bishop, and S. K. Krest (eds.). SETAC Press, Pensacola, Florida.
- Anderson, J. D., and G. K. Williamson. 1976. Terrestrial mode of reproduction in *Ambystoma cingulatum*. *Herpetologica* 32:214–221.
- Arnold, T. W. 2010. Uninformative parameters and model selection using Akaike's information criterion. *Journal of Wildlife Management* 74:1175–1178.
- Beebee, T. J. C. 1995. Amphibian breeding and climate. *Nature* 374:219–220.
- Beebee, T. J. C. 2013. Effects of road mortality and mitigation measures on amphibian populations. *Conservation Biology* 27:657–668.
- Beebee, T. J. C. 2014. Amphibian conservation in Britain: a 40-year history. *Journal of Herpetology* 48:2–12.
- Bliss, C. I., and R. A. Fisher. 1953. Fitting the negative binomial distribution to biological data. *Biometrics* 9:176–200.
- Both, C., Í. L. Kaefer, T. G. Santos, and S. T. Cechin. 2008. An austral anuran assemblage in the Neotropics: seasonal occurrence correlated with photoperiod. *Journal of Natural History* 42:205–222.
- Both, C., and M. E. Visser. 2001. Adjustment to climate change is constrained by arrival date in a long-distance migrant bird. *Nature* 411:296–298.
- Bradshaw, W. E., and C. M. Holzapfel. 2007. Evolution of animal photoperiodism. *Annual Review of Ecology, Evolution, and Systematics* 38:1–25.
- Bucklin, D. N., J. I. Watling, C. Speroterra, L. A. Brandt, F. J. Mazzotti, and S. S. Romanach. 2013. Climate downscaling effects on predictive ecological models: a case study for threatened and endangered vertebrates in the southeastern United States. *Regional Environmental Change* 13: 57–68.
- Canavero, A., M. Arim, D. E. Naya, A. Camargo, I. D. Rosa, and R. Maneyro. 2008. Calling activity patterns in an anuran assemblage: the role of seasonal trends and weather determinants. *North-West Journal of Zoology* 4:29–41.
- Chandler, H. C., A. L. Rypel, Y. Jiao, C. A. Haas, and T. A. Gorman. 2016. Hindcasting historical breeding conditions for an endangered salamander in ephemeral wetlands of the southeastern USA: implications of climate change. *PLoS ONE* 11:1–17.
- Cook, D. G., P. C. Trenham, and P. T. Northen. 2006. Demography and breeding phenology of the California Tiger Salamander (*Ambystoma californiense*) in an urban landscape. *Northwestern Naturalist* 87:215–224.
- Dodd, C. K. 1993. Cost of living in unpredictable environment: the ecology of striped newts *Notophthalmus perstriatus* during a prolonged drought. *Copeia* 1993:605–614.
- Dormann, C. F., J. Elith, S. Bacher, C. Buchmann, G. Carl, G. Carré, J. R. García Márquez, B. Gruber, B. Lafourcade, P. J. Leitão, T. Münkemüller, C. McClean, P. E. Osborne, B. Reineking...S. Lautenbach. 2013. Collinearity: a review of methods to deal with it and a simulation study evaluating their performance. *Ecography* 36:27–46.

- Douglas, M. E. 1979. Migration and sexual selection in *Ambystoma jeffersonianum*. *Canadian Journal of Zoology* 57:2303–2310.
- Edwards, M., and A. J. Richardson. 2004. Impact of climate change on marine pelagic phenology and trophic mismatch. *Nature* 430:881–884.
- Erwin, K. J., H. C. Chandler, J. G. Palis, T. A. Gorman, and C. A. Haas. 2016. Herpetofaunal communities in ephemeral wetlands embedded within longleaf pine flatwoods of the Gulf Coastal Plain. *Southeastern Naturalist* 15:431–447.
- Gibbons, J. W., C. T. Winne, D. E. Scott, J. D. Willson, X. Glaudas, K. M. Andrews, B. D. Todd, L. A. Fedewa, L. Wilkinson, R. N. Tsaliagos, S. J. Harper, J. L. Greene, T. D. Tuberville, B. S. Metts, B. B. Rothermel. 2006. Remarkable amphibian biomass and abundance in an isolated wetland: implications for wetland conservation. *Conservation Biology* 20:1457–1465.
- Gorman, T. A., C. A. Haas, and D. C. Bishop. 2009. Factors related to occupancy of breeding wetlands by flatwoods salamander larvae. *Wetlands* 29:323–329.
- Gorman, T. A., S. D. Powell, K. C. Jones, and C. A. Haas. 2014. Microhabitat characteristics of egg deposition sites used by reticulated flatwoods salamanders. *Herpetological Conservation and Biology* 9:543–550.
- Grant, R. A., E. A. Chadwick, and T. Halliday. 2009. The lunar cycle: a cue for amphibian reproductive phenology? *Animal Behavior* 78:449–357.
- Green, D. M. 2017. Amphibian breeding phenology trends under climate change: predicting the past to forecast the future. *Global Change Biology* 23:646–656.
- Hartel, T., I. Sas, A. P. Pernetta, and I. C. Geltsch. 2007. The reproductive dynamics of temperate amphibians: a review. *North-Western Journal of Zoology* 3:127–145.
- Hegland, S. J., A. Nielsen, A. Lázaro, A-L. Bjerknes, and Ø. Totland. 2009. How does climate warming affect plant-pollinator interactions? *Ecology Letters* 12:184–195.
- Ingram, K., K. Dow, L. Carter, and J. Anderson. 2013. *Climate of the Southeast United States: Variability, Change, Impacts, and Vulnerability*. Island Press, Washington, D.C.
- Jakob, C., G. Poizat, M. Veith, A. Seitz, and A. J. Crivelli. 2003. Breeding phenology and larval distribution of amphibians in a Mediterranean pond network with unpredictable hydrology. *Hydrobiologia* 499:51–61.
- Kelly, J. F., K. G. Horton, P. M. Stepanian, K. de Beurs, T. Fagin, E. S. Bridge, and P. Chilson. 2016. Novel measures of continental-scale avian migration phenology related to proximate environmental cues. *Ecosphere* 7:1–13.
- Knudsen, E., A. Lindén, C. Both, N. Jonzén, F. Pulido, N. Saino, W. J. Sutherland, L. A. Bach, T. Coppack, T. Ergon, P. Gienapp, J. A. Gill, O. Gordo, A. Hedenström, N. C. Stenseth. 2011. Challenging claims in the study of migratory birds and climate change. *Biological Reviews of the Cambridge Philosophical Society* 86:928–946.
- Lindén, A., and S. Mäntyniemi. 2011. Using the negative binomial distribution to model overdispersion in ecological count data. *Ecology* 92:1414–1421.
- Means, D. B., J. G. Palis, and M. Baggett. 1996. Effects of slash pine silviculture on a Florida salamander. *Conservation Biology* 10:426–437.
- Menzel, A., T. H. Sparks, N. Estrella, E. Koch, A. Aasa, R. Ahas, K. Alm-Kübler, P. Bissolli, O. Braslavská, A. Briede, F. M. Chmielewski, Z. Crepinsek, Y. Curnel, A. Dahl, A. Zust. 2006. European phenological response to climate change matches the warming pattern. *Global Change Biology* 12:1969–1976.
- Meredith, H. M., F. A. St. John, B. Collen, S. A. Black, and R. A. Griffiths. 2018. Practitioner and scientist perceptions of successful amphibian conservation. *Conservation Biology* 32:366–375.
- Morbey, Y. E., and R. C. Ydenberg. 2001. Protandrous arrival timing to breeding areas: a review. *Ecology Letters* 4: 663–673.
- Mulholland, P. J., G. R. Best, C. C. Coutant, G. M. Hornberger, J. L. Meyer, P. J. Robinson, J. R. Stenberg, R. E. Turner, F. Vera-Herrera, and R. G. Wetzel. 1997. Effects of climate change on freshwater ecosystems of the south-eastern United States and the Gulf Coast of Mexico. *Hydrological Processes* 11:949–970.
- Palis, J. G. 1997. Breeding migration of *Ambystoma cingulatum* in Florida. *Journal of Herpetology* 31:71–78.
- Palis, J. G., M. J. Aresco, and S. Kilpatrick. 2006. Breeding biology of a Florida population of *Ambystoma cingulatum* (Flatwoods Salamander) during a drought. *Southeastern Naturalist* 5:1–8.
- Palis, J. G., and G. Hammerson. 2008. *Ambystoma bishopi*. The IUCN Red List of threatened species. <https://www.iucnredlist.org/details/136128/0> (accessed 28 July 2017).
- Paton, P. W. C., and W. B. Crouch, III. 2002. Using the phenology of pond-breeding amphibians to develop conservation strategies. *Conservation Biology* 16:194–204.
- Petranka, J. 1998. *Salamanders of the United States and Canada*. Smithsonian Institution Press, Washington, D.C.
- Plard, F., J-M. Gaillard, T. Coulson, A. J. M. Hewison, D. Delorme, C. Warnant, and C. Bonenfant. 2014. Mismatch between birth date and vegetation phenology slows the demography of roe deer. *PLoS Biology* 12:e1001828.
- Post, E., and M. C. Forchhammer. 2008. Climate change reduces reproductive success of an Arctic herbivore through trophic mismatch. *Philosophical Transactions of the Royal Society B: Biological Sciences* 363:2367–2373.
- R Core Team. 2015. R: a language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. <https://www.R-project.org/>
- Roy, D. B., and T. H. Sparks. 2000. Phenology of British butterflies and climate change. *Global Change Biology* 6: 407–416.
- Russell, K. R., D. C. Guynn, Jr., and H. G. Hanlin. 2002. Importance of small isolated wetlands for herpetofaunal diversity in managed, young growth forests in the Coastal Plain of South Carolina. *Forest Ecology and Management* 163:43–59.
- Schaper, S. V., A. Dawson, P. J. Sharp, P. Gienapp, S. P. Caro, and M. E. Visser. 2012. Increasing temperature, not mean temperature, is a cue for avian timing of reproduction. *American Naturalist* 179:55–69.
- Schmidt, B. R., and S. Zumbach. 2008. Amphibian road mortality and how to prevent it: a review, p. 157–167. *In: Urban Herpetology*. J. C. Mitchell, R. E. Jung Brown, and B. Bartholomew (eds.). Society for the Study of Amphibians and Reptiles, Salt Lake City, Utah.
- Semlitsch, R. D. 1985. Analysis of climatic factors influencing migrations of the salamander *Ambystoma talpoideum*. *Copeia* 1985:477–489.
- Semlitsch, R. D. 2008. Differentiating migration and dispersal processes for pond-breeding amphibians. *Journal of Wildlife Management* 72:260–267.



- Semlitsch, R. D., D. E. Scott, J. H. K. Pechmann, and J. W. Gibbons. 1993. Phenotypic variation in the arrival time of breeding salamanders: individual repeatability and environmental influences. *Journal of Animal Ecology* 62:334–340.
- Sexton, O. J., C. Phillips, and J. E. Bramble. 1990. The effects of temperature and precipitation on the breeding migration of the spotted salamander (*Ambystoma maculatum*). *Copeia* 1990:781–787.
- Strzepek, K., G. Yohe, J. Neumann, and B. Boehlert. 2010. Characterizing changes in drought risk for the United States from climate change. *Environmental Research Letters* 5:044012.
- Stuart, S. N., J. S. Chanson, N. A. Cox, B. E. Young, A. S. L. Rodrigues, D. L. Fischman, and R. W. Waller. 2004. Status and trends of amphibian declines and extinctions worldwide. *Science* 306:1783–1786.
- Terceiro, M. 2003. The statistical properties of recreational catch rate data for some fish stocks off the northeast US coast. *Fishery Bulletin* 101:653–672.
- Thackeray, S. J., T. H. Sparks, M. Frederiksen, S. Burthe, P. J. Bacon, J. R. Bell, M. S. Botham, T. M. Brereton, P. W. Bright, L. Carvalho, T. Clutton-Brock, A. Dawson, M. Edwards, J. M. Elliott, S. Wanless. 2010. Trophic level asynchrony in rates of phenological change for marine, freshwater and terrestrial environments. *Global Change Biology* 16:3304–3313.
- Timm, B. C., K. McGarigal, and B. W. Compton. 2007. Timing of large movement events of pond-breeding amphibians in western Massachusetts, USA. *Biological Conservation* 136:442–454.
- Todd, B. D., D. E. Scott, J. H. Pechmann, and J. W. Gibbons. 2011. Climate change correlates with rapid delays and advancements in reproductive timing in an amphibian community. *Proceedings of the Royal Society of London B: Biological Sciences* 278:2191–2197.
- Todd, B. D., and C. T. Winne. 2006. Ontogenetic and interspecific variation in timing of movement and responses to climatic factors during migrations by pond-breeding amphibians. *Canadian Journal of Zoology* 84: 715–722.
- Trenham, P. C., H. B. Shaffer, W. D. Koenig, and M. R. Stromberg. 2000. Life history and demographic variation in the California tiger salamander (*Ambystoma californiense*). *Copeia* 2000:365–377.
- U.S. Fish and Wildlife Service (USFWS). 1999. Final rule to list the flatwoods salamander as a threatened species. *Federal Register* 64:15691–15704.
- U.S. Fish and Wildlife Service (USFWS). 2009. Determination of endangered status for reticulated flatwoods salamander. *Federal Register* 74:6700–6774.
- Valtonen, A., M. P. Ayres, H. Roininen, J. Pöyry, and R. Leinonen. 2011. Environmental controls on the phenology of moths: predicting plasticity and constraint under climate change. *Oecologia* 165:237–48.
- Ver Hoef, J. M., and P. L. Boveng. 2007. Quasi-Poisson vs. negative binomial regression: how should we model overdispersed count data? *Ecology* 88:2766–2772.
- Visser, M. E., L. J. M. Hollemann, and P. Gienapp. 2006. Shifts in caterpillar biomass phenology due to climate change and its impact on the breeding biology of an insectivorous bird. *Oecologia* 147:164–172.
- Visser, M. E., A. J. van Noordwijk, J. M. Tinbergen, and C. M. Lessels. 1998. Warmer springs lead to mistimed reproduction in Great Tits (*Parus major*). *Proceedings of the Royal Society: Biological Sciences B* 265:1867–1870.
- Walls, S. C., L. C. Ball, W. J. Barichivich, C. K. Dodd, Jr., K. M. Enge, T. A. Gorman, K. M. O'Donnell, J. G. Palis, and R. D. Semlitsch. 2017. Overcoming challenges to the recovery of declining amphibian populations in the United States. *Bioscience* 67:156–165.
- Walls, S. C., W. J. Barichivich, and M. E. Brown. 2013. Drought, deluge and declines: the impact of precipitation extremes on amphibians in a changing climate. *Biology* 2: 399–418.
- Williams, C. T., B. M. Barnes, G. J. Kenagy, and C. L. Buck. 2014. Phenology of hibernation and reproduction in ground squirrels: integration of environmental cues with endogenous programming. *Journal of Zoology* 292:112–124.