

Real-Time PCR Reveals Endosymbiont Titer Fluctuations in *Metaseiulus occidentalis* (Acari: Phytoseiidae) Colonies Held at Different Temperatures

Authors: Jeyaprakash, Ayyamperumal, and Hoy, Marjorie A.

Source: Florida Entomologist, 93(3) : 464-466

Published By: Florida Entomological Society

URL: <https://doi.org/10.1653/024.093.0326>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

REAL-TIME PCR REVEALS ENDOSYMBIONT TITER FLUCTUATIONS IN *METASEIULUS OCCIDENTALIS* (ACARI: PHYTOSEIIDAE) COLONIES HELD AT DIFFERENT TEMPERATURES

AYYAMPERUMAL JEYAPRAKASH¹ AND MARJORIE A. HOY
Department of Entomology and Nematology, University of Florida, Gainesville, FL 32611

¹Presently at Bureau of Entomology, Nematology and Plant Pathology, Florida Department of Agriculture and Consumer Services, Division of Plant Industry, Gainesville, FL 32608

Real-time PCR (Polymerase Chain Reaction) can estimate *Wolbachia* endosymbiont population density in mosquitoes (Wiwatanaratnabutr & Kittayapong 2009). The COS (Carbaryl-QP-Sulfur resistant) colony of the phytoseiid *Metaseiulus* (= *Typhlodromus* or *Galendromus*) *occidentalis* (Nesbitt) harbors several endosymbionts, including *Cardinium*, *Wolbachia*, *Enterobacter*, and *Bacteroidetes* in their gut and reproductive tissues (Johanowicz & Hoy 1996; Hoy & Jeyaprakash 2005). Johanowicz & Hoy (1998) observed mating incompatibility when females from a heat-treated (33°C) COS colony (a treatment presumed to eliminate *Wolbachia*) was crossed with males from a room-temperature (23°C) COS colony containing *Wolbachia*, resulting in the production of shriveled eggs that did not hatch. It was not clear whether the *Wolbachia* population was reduced or completely eliminated from the heat-treated mites because a standard PCR protocol was used, which is relatively insensitive compared to high-fidelity PCR (Jeyaprakash & Hoy 2000). In this paper, we estimate the endosymbiont population density in COS colonies of *M. occidentalis* held under different temperatures in the laboratory using real-time PCR.

One colony of *M. occidentalis* was maintained at room-temperature (23-25°C) and 2 colonies initiated from this line were maintained at 32 and 34°C for More than 1 year in growth chambers

prior to experiments. Female mites (100) were isolated from each colony and their genomic DNA extracted by Puregene and QIAGEN methods (Jeyaprakash & Hoy 2007), and resuspended in 50 µL of sterile water. Four different plasmids carrying endosymbiont 16S rRNA sequences; pAJ233 (*Cardinium*), pAJ234 (*Bacteroidetes*), pAJ235 (*Wolbachia*) and pAJ239 (*Enterobacter*), were extracted and their yield estimated with a BioPhotometer Plus (Eppendorf AG, Hamburg, Germany). Species-specific primers were designed with Primer 3 software (<http://frodo.wi.mit.edu/primer3/>) and a probe was designed with Primer X software (Applied Biosystems, Foster City, CA) from the variable regions for each species (Table 1). Real-time PCR was performed in a 20-µL reaction volume containing genomic DNA from 2 females (1 µL) or serially-diluted plasmid DNA (100 µg to 1 fg) or a no DNA control (1 µL), forward and reverse primers (400 pM), probe (100 pM) and TaqMan master mix (10 µL). Two linked profiles: (1) one cycle of 50°C for 2 min and 95°C for 10 min, and (2) 60 cycles consisting of denaturation at 95°C for 15 s, annealing at 59°C (*Cardinium*) or 57°C (*Wolbachia* or *Enterobacter* or *Bacteroidetes*) for 30 s and extension at 72°C for 30 s, were used. Each treatment was replicated 3 times. The starting copy number of all plasmid DNA dilutions used in the real-time PCR was obtained with an open-source software ([TABLE 1. LIST OF SPECIES-SPECIFIC PRIMERS AND PROBES DESIGNED FOR REAL-TIME PCR.](http://</p></div><div data-bbox=)

Species	Primer and probe sequence	Product size
<i>Cardinium</i>	MoCardRT-F2 5'-GCCGCGCACCGGCGAATG-3'	66 bp
	MoCardRT-R1 5'-CGGAGGCTATCCCCAGTGT-3'	
	MoCardProbe1 6FAM-TGCGTAATGCACATGC-MGBNFQ	
<i>Wolbachia</i>	MoWoIRT-F1 5'-GCAACGCGAAAAACCTTACCAC-3' MoWoIRT-R1 5'-CCGACCCTATCCCTTCGAAT-3' MoWoProbe 1 6FAM-TCTTGACATGAAAATC-MGBNFQ	65 bp
<i>Bacteroidetes</i>	MoBactRT-F1 5'-AGCGGCAGGCCAATACATG-3' MoBactRT-R1 5'-CTCACCATCTTCGAGCAAGCT-3' MoBactProbe 1 6FAM-AAGTCGGACGGGATC-MGBNFQ	65 bp
<i>Enterobacter</i>	MoEnterRT-F1 5'-TGCCAGCAGCCGCGGTAAT-3' MoEnterRT-R1 5'-TTTACGCCAGTAATTCCGATT-3' MoEnterProbe 1 6FAM-CGGAGGGTGCAAGC-MGBNFQ	59 bp

/molbiol.edu.ru/eng/scripts/h01_07.html). A standard regression curve was generated with the Ct-value (Number of PCR cycles required for sample fluorescence to reach the threshold level) obtained from all 9 serial plasmid DNA dilutions. The Ct-value obtained from the 2 female mites was then used in the regression analysis to estimate the copy number of bacterial cells present, which was divided by 2 to obtain the copy numbers in a single female.

The titer of *Cardinium* and *Wolbachia* was much higher than that of the gut endosymbionts *Enterobacter* and *Bacteroidetes* in the room-temperature *M. occidentalis* COS colony (Table 2). The *Cardinium*, *Wolbachia* and *Enterobacter* titer was reduced, but not eliminated, in both heat-treated colonies. By contrast, *Bacteroidetes* had a higher titer in the heat-treated colonies, but their density was below detectable levels (1 or 2) in the room-temperature colony, and could not be reliably amplified by real-time PCR (Table 2).

We think that the titer of *Wolbachia* or *Cardinium* could play a role in mating compatibility when making crosses between heat-treated (uninfected) females and room-temperature (infected) males. However, it is not clear which endosymbiont actually causes mating incompatibility in the COS colony because this requires generating colonies infected with only *Cardinium* or *Wolbachia*. Weeks & Stouthamer (2004) reported that *Wolbachia* infection in another colony of *M. occidentalis* is lost when fed a *Wolbachia*-free two-spotted spider mite diet. During the summer of 2009 we determined, by high-fidelity PCR, that the two-spotted spider mite culture in our greenhouse had inadvertently lost *Wolbachia* after being heated to 40-42°C for several months (3-6) due to a maintenance problem and, subsequently, the 23-25, 32 and 34°C COS colonies that were fed this *Wolbachia*-free spider mite diet were determined to have lost *Wolba-*

chia (data not shown), suggesting that the *Wolbachia* in these colonies of *M. occidentalis* was obtained from their spider mite prey.

To determine whether the *Wolbachia*-free COS colonies reared at 23-25 and 34°C exhibited incompatibility several inbred lines were generated. They displayed no fluctuation in *Cardinium* and *Enterobacter* titers (Table 2). A total of 7 or 8 reciprocal single-pair and control crosses were performed on 3 separate dates. Equal numbers of reciprocal and control crosses were successful and produced unshriveled eggs that subsequently hatched, suggesting that *Cardinium* is not involved in causing cytoplasmic incompatibility in the COS colony at these titers (Table 2) and that *Wolbachia* might be involved. It appears that this COS colony is obtaining *Wolbachia* by feeding on its *Wolbachia*-infected prey. These results are consistent with earlier data indicating that there are no differences in *16S*, *fts* and *wsp* sequences from *Wolbachia* in the COS and *T. urticae* colonies (Hoy & Jeyaparakash 2005).

The standard PCR protocol previously used by Johanowicz & Hoy (1996) we now know is not sufficiently sensitive to be certain that endosymbionts truly are lacking (Jeyaparakash & Hoy 2000) and should not be used when comparing heat-treated and room-temperature colonies. Real-time PCR is an efficient tool to quantify endosymbiont titer based on this study and that by Wiwatanaratanaabutr & Kittayapong (2009). This research was supported in part by the Davies, Fischer and Eckes Endowment in Biological Control to M. A. Hoy.

SUMMARY

Real-time PCR amplification of *16S rRNA* sequences from 4 *M. occidentalis* endosymbionts; *Cardinium*, *Wolbachia*, *Enterobacter*, and

TABLE 2. MEAN NUMBER (S.D.) OF ENDOSYMBIONTS ESTIMATED PER ADULT FEMALE *METASEIULUS OCCIDENTALIS* MAINTAINED AT 3 DIFFERENT TEMPERATURES¹.

Date	Colony	Rearing temperature	<i>Cardinium</i>	<i>Wolbachia</i>	<i>Enterobacter</i>	<i>Bacteroidetes</i>
8/10/09	COS	23-25°C	26,666 ± 23	13,585 ± 22	379 ± 12	BDL
	COS	32°C	127 ± 16	68 ± 15	10 ± 2	543 ± 12
	COS	34°C	138 ± 16	74 ± 15	15 ± 2	499 ± 12
9/25/09 COS inbred lines	C10-BaBA	23-25°C	18,516 ± 5	ND	264 ± 5	BDL
	C10-BA	23-25°C	19,199 ± 5	ND	274 ± 5	BDL
	C10-CB	23-25°C	20,892 ± 5	ND	295 ± 5	BDL
	F10-FAB	23-25°C	16,711 ± 5	ND	278 ± 5	BDL
	F10-BAA	23-25°C	19,907 ± 5	ND	322 ± 5	BDL
	F10-IOO	23-25°C	19,315 ± 5	ND	329 ± 5	BDL

¹Means were obtained based on 3 reactions per condition.

BDL = Below detectable level (1 or 2); COS = Carbaryl-OP-Sulfur resistant; ND = Not detected (0) and had been fed two-spotted spider mite prey lacking *Wolbachia*.

Bacteroidetes, provided an estimate of their titer in colonies reared under 3 temperatures. This work also revealed, for the first time, that the *Wolbachia* and *Cardinium* titer were reduced in heat-treated COS colonies, but not completely eliminated. However, feeding *M. occidentalis* colonies the *Wolbachia*-free two-spotted spider mite diet did eliminate *Wolbachia* from this predator. No cytoplasmic incompatibility was detected when crosses were made between males reared at room-temperature (23-25°C) containing *Cardinium* (and lacking *Wolbachia*) and females reared at 34°C containing a much lower titer of *Cardinium* (and lacking *Wolbachia*), indicating that *Wolbachia* may be involved in causing previously observed incompatibilities rather than *Cardinium*.

REFERENCES CITED

- HOY, M. A., AND JEYAPRAKASH, A. 2005. Microbial diversity in the predatory mite *Metaseiulus occidentalis* (Acari: Phytoseiidae) and its prey, *Tetranychus urticae* (Acari: Tetranychidae). Biol. Control 32: 427-441.
- JEYAPRAKASH, A., AND HOY, M. A. 2000. Long PCR improves *Wolbachia* DNA amplification: *wsp* sequences found in 76% of sixty-three arthropod species. Insect Mol. Biol. 9(4): 393-405.
- JEYAPRAKASH, A., AND HOY, M. A. 2007. The mitochondrial genome of the predatory mite *Metaseiulus occidentalis* (Arthropoda: Chelicerata: Acari: Phytoseiidae) is unexpectedly large and contains several novel features. Gene 391: 264-274.
- JOHANOWICZ, D. L., AND HOY, M. A. 1996. *Wolbachia* in a predator-prey system: 16S ribosomal analysis of two phytoseiids (Acari: Phytoseiidae) and their prey (Acari: Tetranychidae). Ann. Entomol. Soc. America 89: 435-441.
- JOHANOWICZ, D. L., AND HOY, M. A. 1998. Experimental induction and termination of non-reciprocal reproductive incompatibilities in a parahaploid mite. Entomol. Expt. Appl. 87: 51-58.
- WEEKS, A. R., AND STOUTHAMER, R. 2004. Increased fecundity associated with infection by a *Cytophaga*-like intracellular bacterium in the predatory mite, *Metaseiulus occidentalis*. Proc. R. Soc. London B (Suppl.) 271: S193-S195.
- WIWATANARATANABUTR, I., AND KITTAAPONG, P. 2009. Effects of crowding and temperature on *Wolbachia* infection density among life cycle stages of *Aedes albopictus*. J. Invert. Path. 102: 220-224.