

New Collection Records of *Perileptus areolatus* (Coleoptera: Carabidae) from the Amanos Mountains of Turkey

Author: Avgin, Sakine Serap

Source: Florida Entomologist, 97(4) : 1364-1368

Published By: Florida Entomological Society

URL: <https://doi.org/10.1653/024.097.0410>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

NEW COLLECTION RECORDS OF *PERILEPTUS AREOLATUS* (COLEOPTERA: CARABIDAE) FROM THE AMANOS MOUNTAINS OF TURKEY

SAKINE SERAP AVGIN

Kahramanmaraş Sütçü İmam University, Faculty of Education, Division of Science Education, Avşar Campus,
46100 Kahramanmaraş, Turkey
E-mail: ssavgin@ksu.edu.tr

Supplementary material for this article Florida Entomologist 97(4) (December 2014)
is online at <http://purl.fcla.edu/fcla/entomologist/browse>.

ABSTRACT

Specimens from *Perileptus* Schaum, 1860 (Coleoptera: Carabidae) were collected from the Amanos Mountains of Turkey, and this genus was redescribed. Descriptions of *Perileptus* species are presented herein with zoogeographic, biological, and ecological observations. This is the first detailed study of Turkish *Perileptus*, and it provides the first records and information on *Perileptus* from the Amanos Mountains.

Key Words: Amanos Mountains, Anatolian Diagonal *Bembidion*, syrio-exemial faunal elements, *Trechus*

RESUMEN

Se redescrive de *Perileptus* Schaum, 1860 (Coleoptera: Carabidae) con especímenes recolectados de las Montañas Amanos en Turquía. Se presentan descripciones de las especies *Perileptus* con observaciones zoogeográficas, biológicas y ecológicas. Este es el primer estudio detallado de los *Perileptus* de Turquía y se provee los primeros datos e información sobre *Perileptus* de las Montañas Amanos.

Palabras Clave: Montañas Amanos, *Bembidion* Anatolia Diagonal, elementos faunísticos sirio-exemial, *Trechus*

The genus *Perileptus* Schaum, 1860 (Coleoptera: Carabidae) has been placed in the subfamily Trechiniinae. Most species of this genus have small, flat, and narrow individuals. *Perileptus* is superficially intermediate between *Bembidion* and *Trechus* as evidenced by the moderately reduced terminal segment of the maxillary palp (Lindroth 1985). In addition, the sutural stria is not recurrent at the apex of the elytra, but the frontal furrows are strongly divergent behind the eyes, as in *Trechus* (Lindroth 1985). Beetles in this genus have full wings. Males have 2 dilated segments on both the prothoracic and mesothoracic tarsi (Lindroth, 1985). Adult beetles in this genus have a spine on the outer edge of the anterior margin of the front tibia. Their eyes are pubescent and large. The penultimate labial palpomere has at least 5 setae. The average size of *Perileptus* beetles are less than 2.5 mm, which differs from other genera in this subfamily (Hurka 1996).

The larvae of this genus can be identified by their tarsi, which have 2 equal claws and a pair of flattened apical setae, and by the toothed cutting edge of their mandibles (Luff 1993). Larvae of *Perileptus* species live in fine gravel along the edge of fast flowing streams (Trautner & Geigenmüller 1987; Hurka 1996).

Worldwide, the genus *Perileptus* comprises 57 species and subspecies (Lorenz 1998). According to Hurka (1996), *Perileptus* is distributed mostly in the tropics of the Old World with a single species in

Europe. There are 26 species and subspecies known from the Palearctic region (Löbl & Smetana 2003). Of these, *Perileptus areolatus areolatus* Creutzer is the only species that has been reported from Turkey.

There have been some studies of carabids from the Amanos Mountains (Korell 1988, 2001), but there are no published records of *Perileptus* from this region. The Amanos Mountains encompass a national park that is important not only in Turkey but also worldwide. The Amanos Mountains, which are one of the most tectonic ranges in the world, possess a unique importance because of their geographical position (Okur & Yalçın-Özdilek 2008). These mountains provide a bridge between the Taurus Mountains, Lebanon Mountains, and North Syrian Desert, and allow species belonging to this region to disperse among these areas (Boulos et al. 1994; Okur & Yalçın-Özdilek 2008). The Amanos Mountain range provides a barrier for the distribution of syrio-exemial faunal elements in the west and Mediterranean faunal element in the east (Çıplak et al. 1993; Demirsoy 1996; Uğurtaş et al. 2000). In addition, this mountain range is thought to be a part of the hypothetical Anatolian Diagonal, which possibly affected the faunal structure of Anatolia (Çıplak et al. 1993; Uğurtaş et al. 2000). The Amanos Mountains have rich and diverse fauna and flora because of its geographical position.

This study redescribes the genus *Perileptus* from Turkey, and it provides some zoogeographic, biologi-

cal, and ecological observations and information about this genus from the Amanos Mountains.

MATERIALS AND METHODS

Sampling Locations

Perileptus specimens were collected in 2008 and 2010 from 4 locations at different elevations in the Amanos Mountains. Samples were collected by hand from gravelly-stoney river banks:

1. Amanos mountains, Hatay Province, Sinanlı, Karaçay: N 36° 05' 42" E 36° 04' 50", and 33 m asl (Fig. 1).
2. Amanos mountains, Hatay Province, Hassa, Demrek: N 36° 40' 47" E 36° 25' 17", and 458 m asl (Fig. 1).
3. Amanos mountains, Hatay Province, Dörtöl-Kuzuculu-Deliçay: N 36° 53' 21" E 36° 15' 17", and 152 m asl.
4. Amanos mountains, Hatay Province, Dörtöl-Payas-Karbeyaz: N 36° 43' 27" E 36° 14' 07", and 249 m asl.

The specimens collected were brought to laboratory, mounted, and identified to species using keys, published descriptions, comparing them with pinned specimens. Measurements were made using a SMZ 1000 Nikon stereobinocular microscope. In addition, photographs of the main diagnostic characters were taken with a Kameram 3.2 camera through the stereomicroscope. Type specimens of the new species were deposited.

Descriptions include details of external features and information about their geographical distribution within the study area, Turkey, and worldwide.

The following abbreviations were used for the measurements, which are given in mm:

- HL: length of head from anterior margin of clypeus to posterior margin of temples;
 HL(m): length of head from apex of mandibles to posterior margin of temples;
 HW: width of head, not including eyes;
 HW(e): width of head, including eyes;
 PL: length of pronotum along median line;
 PW: maximum width of pronotum;
 EL: length of elytra from humeral tubercle to the apex;
 EW: maximum width of elytra;
 BL(a): body length (from head to elytra apex);
 BL(b): total body length, from apices of mandibles to those of elytra.

Figures 1 and 2 are displayed in color online in supplementary material for this article in Florida Entomologist 97(4) (December 2014) at <http://purl.fcla.edu/fcla/entomologist/browse>.

RESULTS AND DISCUSSION

Genus *Perileptus* Schaum, 1860. [Ganglb., Käf. v. Mitteleur. I., p.185; Bedel, Cat. rais. Col. du Nord de l'Afr. 1896, pp. 82 (Apfelbeck 1904)]. [*Perileptus* Schaum, 1860, pp. 663. Type-species: *Carabus areolatus* Creutzer, 1799. For diagnosis see Jeannel (1926, pp. 402) and Moore (1972, pp. 12) (Baehr 1987)].

Description of *Perileptus areolatus* Creutzer from Turkey (Fig. 2A-D)

Body length (from head to elytra apex) 2.47-2.59 mm, width 0.88-0.92 mm; labrum without an indistinct central tubercle, vertex without punctures, media line strongly conspicuous and reddish-brown, posterior angles slightly protrude, elytra with black spot narrowed from basal to medial at 1st quarter and along apical margin with black spot, elytra color red-brown.

Perileptus areolatus Creutzer, 1799. Wide-spread in the Balkan Peninsula: Bosnia-Herzegovina, frequently on the banks of flowing water streams; Dalmatia (Knin, Castelnuovo, Cattaro), Montenegro (Rjeka-Mustajb.), Serbia (Maidan Kutšajna, Negotin Dol Milanovac Golubac-Merkl.), Bulgaria (Sofia on Iskerflusse-Apflb), Greece (Attica, Thessaly-Krüper; Doris Oertzen-v, dir p 206), and Crete (Zebe). In Asia Minor (Smyrna Krüper, according to V. Bodemeyer anywhere frequently) (Apfelbeck 1904)]. Sometimes seen under pebbles of large streams. Captured at Smyrna, the Hermon River, the Meandron River, the island of Lesbos, and in the valley of the Alps Bulghar Kara Dag (Shalberg 1913).

Material Examined

Amanos Mountains, Hatay Province, Hassa, Demrek: 04.VI.2010, 1♀; Amanos Mountains, Hatay Province, Dörtöl-Kuzuculu-Deliçay: 03.VI.2010, 1♂; Hatay-Sinanlı-Karaçay (Orman Ürümeleri Deposu): 05.VI.2010, 1♀; Amanos Mountains, Hatay Province, Dörtöl-Payas-Karbeyaz: 06.VI.2009, 2♀♀; 03.VI.2010, 1♀ (4 specimens were damaged when collected).

Description

Measurements (mm): HL: 0.39-0.42; HL(m): 0.55-0.59; HW: 0.42-0.45; HW(e): 0.58-0.62; PL: 0.49-0.53; PW: 0.65-0.70; EL: 1.46-1.51; EW: 0.88-0.92; BL(a): 2.47-2.59; BL(b): 2.63-2.76 (6 specimens measured).

Coloration, from reddish to dark-brown, head reddish-dark brown, antennae red-brown, maxillary palp and labial palp red-brown, mandible red-brown, pronotum from reddish to dark-brown, elytra red-brown, legs red-brown, body ventral from black to dark-brown.



Fig. 1. Collection locations of *Perileptus*; (A) Amanos Mountains, Hatay Province, Sinanlı, Karaçay, and (B) Amanos Mountains, Hatay Province, Hassa, Demrek. This figure is displayed in color online in supplementary material for this article in Florida Entomologist 97(4) (December 2014) at <http://purl.fcla.edu/fcla/entomologist/browse>.




Fig. 2. *Perileptus areolatus* Creutzer, 1799; (A) labrum; (B) pronotum; (C) habitus and dorsal aspect of male, and (D) head and labrum. This figure is displayed in color online in supplementary material for this article in Florida Entomologist 97(4) (December 2014) at <http://purl.fcla.edu/fcla/entomologist/browse>.

Head narrower than pronotum, glabrous, with punctures fine, labrum without an indistinct central tubercle (Fig. 2B), basal like a half moon, with 6 setae in basal marginal, the last segment of maxillary palp and labial palp very thin, mandible strongly large and apex thick, vertex without punctures.

Pronotum (PW/PL: 1.32) (EW/PW: 1.35-1.31) punctures fine and sparsely laying hairs on the marginal sides, with superficial weak microsculpture, anterior margin concave, lateral sides slightly sinuate and margins narrow toward posterior and posterior angles slightly protrude, media line strongly conspicuous and reddish-brown,

posterior angles with 1 seta, basal depressions slightly marked (Fig. 2A).

Elytra (EL/EW: 1.66-.164) elytra with black spot narrowed from basal to medial at 1st quarter and along apical margin with black spot (Fig. 2C).

Distribution

Asia: Turkey (Anatolia) (Casale & Vigna Taglianti 1999; Löbl & Smetana 2003), Iran, Israel, Saudi Arabia, Syria, Turkey (Löbl & Smetana 2003). Europe (Trautner & Geigenmüller 1987): Bohemia, Moravia, Slovakia (Hurka 1996); Azerbaijan, Albania, Armenia, Austria, Bosnia-Herzegovina, Belarus, Croatia, Czech Republic, France, Great Britain, Germany, Georgia, Greece, Hungary, Ireland, Italy, Latvia, Macedonia, Moldova, Norway, Poland, Portugal, Romania, Russia, Slovakia, Slovenia, Spain, Sweden, Switzerland, Ukraine, Yugoslavia (Löbl & Smetana 2003). North Africa: Algeria, Morocco, Tunisia (Löbl & Smetana 2003).

Biology and Ecological Notes

The period of reproduction for *P. areolatus* is in late spring, and females with mature eggs have been found in Jun (Lindroth 1985). This species lives in fine sand and pebbles by rivers; and it is known to fly. *Perileptus areolatus* has a primarily southern European distribution, extending as far north as southern Scandinavia (Luff 1998). Although, *P. areolatus* has been described from several countries, including Turkey, this is the first report of specimens from the Amanos Mountains. The specimens were mostly collected in May and Jun at altitudes of 33-458 m. Collections were difficult because these beetles move very quickly.

ACKNOWLEDGMENTS

I am grateful to the Scientific and Technical Research Council of Turkey (TUBITAK) for their financial support (Project 109T559).

REFERENCES CITED

- APFELBECK, V. 1904. Die Käferfauna der Balkanhalbinsel, mit Berücksichtigung Klein-Asiens und der Insel Kreta, Familienreihe Caraboidea [The beetle fauna of the Balkan Peninsula, with consideration of Asia Minor and the island of Crete, family series Caraboidea]. R. Frieländer & Son, Berlin, Germany, 422 pp.
- BAEHR, M. 1987. The Australian species of the Carabid genus *Perileptus* (Coleoptera: Carabidae: Trechinae). Invert. Taxon. 1: 1-16.
- BOULOS, L., A. G. MILLER, AND R. R. MILL. 1994. Regional overview: South West Asia and the Middle East, pp. 293-349 In S. D. Davis, V. H. Heywood and A. C. Hamilton [eds.], Centres of Plant Diversity, A guide and strategy for their conservation. Volume 1: Europe, Africa, South West Asia and the Middle East. Inform. Press, Oxford, UK.
- CASALE, A., AND VIGNA TAGLIANTI, A. 1999. Caraboid beetles (excl. Cicindelidae) of Anatolia, and their biogeographical significance (Coleoptera, Caraboidea). Biogeographia 20: 277-406.
- ÇIPLAK, B., DEMİRİSOY, A., AND BOZCUK, N. 1993. Distribution of Orthoptera in relation to the Anatolian diagonal in Turkey. Articulata 8(1): 1-20.
- DEMİRİSOY, A. 1996. Genel ve Türkiye Zoocoğrafyası "Hayvan Coğrafyası". Metaksan A.Ş. Ankara.
- HURKA, K. 1996. Carabidae of the Czech and Slovak Republics. V. Kabourek Publ., Zlin, Czech Republic, 565 pp.
- KORELL, A. 1988. Ein bemerkenswerter Fund von *Procerus syriacus* Kollar 1843 in der Provinz Hatay, Türkei (Coleoptera: Carabidae). Entomol. Z. 98(7): 92-95.
- KORELL, A. 2001. DIE Arten der Gattungen *Carabus* Linnaeus, 1758 und *Procerus* Dejean, 1828 der Provinz Hatay, Südtürkei (Coleoptera: Carabidae, Carabini) [The species of the genus *Carabus* Linnaeus, 1758 and *Procerus* Dejean, 1828 in the province of Hatay, south Turkey (Coleoptera: Carabidae, Carabini)]. Nachr. Entomol. Verein Apollo, Neue Folge 22(2): 66.
- LINDROTH, C. H. 1985. The Carabidae (Coleoptera) of Fennoscandia and Denmark. Fauna Entomol. Scandinavica, vol 15, part 1. Scandinavian Science Press, Ltd., Copenhagen, 225 pp.
- LORENZ, W. 1998. Systematic list of extant ground beetles of the world: Insecta, Coleoptera "Geadephaga": Trachypachidae and Carabidae incl. Paussinae, Cicindelinae, Rhysodinae. Published by the author, Tutzing, Germany, 503 pp.
- LÖBL, I., AND SMETANA, A. 2003. Catalogue of Palaearctic Coleoptera. Volume I. Archostemata-Myxophaga-Adephaga. Apollo Books, Stenstrup, Denmark, 819 pp.
- LUFF, M. L. 1993. The Carabidae (Coleoptera) larvae of Fennoscandia and Denmark. Fauna Entomol. Scandinavica, vol 27. E. J. Brill Publ., Leiden, The Netherlands, 186 pp.
- LUFF, M. L. 1998. Provisional Atlas of the Ground Beetles (Coleoptera, Carabidae) of Britain. Biological Records Centre, Huntingdon, England, 194 pp.
- OKUR, E., AND YALÇIN-ÖZDİLEK, Ş. 2008. Preliminary study of the fish community structure in Amanos mountain streams (Hatay-Turkey). Biologia 63: 427-438.
- SAHLBERG, J. 1913. Coleoptera Mediterranea Orientalia, quae in Aegypto, Palaestina, Syria, Carmania atque Anatolia Occidentalia Anno 1904 collegerunt John Sahlberg et Unio Saalas [Beetles of the eastern Mediterranean, Egypt, Palestine, Syria, west Anatolia, and west Carmania, and a collection of John Sahlberg and Unio Saalas in 1904]. Öfversikt af Finska Vetenskaps-Societetens Förhandlingar, A 55 (19): 281 pp.
- TRAUTNER J., AND GEIGENMÜLLER, K. 1987. Tiger Beetles, Ground Beetles. Illustrated key to the Cicindelidae and Carabidae of Europe. Josef Margraf Publ., Aichtal, Germany, 488 pp.
- UĞURTAŞ, İ. H., YILDIRIMHAN, H. S., AND ÖZ, M. 2000. Herpetofauna of the eastern region of the Amanos Mountains (Nur). Turkish J. Zool. 24: 257-261.