

Identification of Megachilid Species (Hymenoptera: Megachilidae) and Other Pollinators in Apple Orchards in Chihuahua, México

Authors: Rios-Velasco, Claudio, Berlanga-Reyes, David I., Ayala-Barajas, Ricardo, Salas-Marina, Miguel Ángel, Ibarra-Rendón, Jorge E., et al.

Source: Florida Entomologist, 97(4) : 1829-1834

Published By: Florida Entomological Society

URL: <https://doi.org/10.1653/024.097.0460>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

IDENTIFICATION OF MEGACHILID SPECIES (HYMENOPTERA: MEGACHILIDAE) AND OTHER POLLINATORS IN APPLE ORCHARDS IN CHIHUAHUA, MÉXICO

CLAUDIO RIOS-VELASCO¹, DAVID I. BERLANGA-REYES¹, RICARDO AYALA-BARAJAS², MIGUEL ÁNGEL SALAS-MARINA³, JORGE E. IBARRA-RENDÓN³, PAUL B. ZAMUDIO FLORES¹, ALEJANDRO ROMO-CHACÓN¹, AND CARLOS H. ACOSTA-MUÑOZ¹

¹Centro de Investigación en Alimentación y Desarrollo, A.C. Unidad Cuauhtémoc, Chihuahua, Av. Río Conchos S/N Parque Industrial. Z.C. 31570, Cuauhtémoc, Chihuahua, México

²Estación de Biología Chamela, Instituto de Biología, Universidad Nacional Autónoma de México, Z.C. 48980, San Patricio, Jalisco, México

³Departamento de Biotecnología y Bioquímica, Centro de Investigación y de Estudios Avanzados del Instituto Politécnico Nacional, Unidad Irapuato, Km. 9.6 Libramiento Norte, Z.C. 36821, Irapuato, Guanajuato, México

Corresponding author; E-mail: claudio.rios@ciad.mx

Animal pollination is a mutually beneficial to both plants and pollinators. Plants benefit by being pollinated, which assures seed formation and sexual reproduction; and in return, the plant provides the pollinator with nectar, pollen and sweet fruits. Pollination has contributed to the evolution of floral diversity, mainly of angiosperms (Cane 2008; Sheffield et al. 2012). Thirty five percent of the world's food production (vegetables, fruits, edible oil crops, stimulants, nuts and spices) depends on crop pollination by honey bees, bumblebees and solitary bees (Kraemer & Favi 2005; Klein et al. 2007; Gallai et al. 2009; Das et al. 2011). Recently an appreciation of the fragility of many plant-pollinator relationships has emerged, along with the realization that it is necessary to understand the complex relationships and factors that influence the richness and abundance of species in the ecosystem in order to prevent its decline in a wide range of habitats (Biesmeijer et al. 2006; Winfree 2010; Williams et al. 2010). Pollination management in apple orchards is a critically important activity to assure fruit set and quality (Kraemer & Favi 2005; Das et al. 2011). Solitary bees of the genus *Osmia* (Hymenoptera: Megachilidae), known as effective pollinators, have excellent anatomical and behavioral adaptations and lifestyles dependent entirely of floral phenology (Drummond & Stubbs 1997; Kraemer & Favi 2005; Felicioli & Pinzauti 2008). Pollinators bees including *Osmia* species are favored by warm climates; however, they are still active in cool and cloudy days, which are frequent during the flowering period in apple orchards of Cuauhtémoc and Guerrero, Chihuahua, México. *Osmia* species are non-aggressive to humans and can easily be induced to nest in artificial nests, which are the basis for increasing the use of *Osmia* species in the management of apple orchards (Felicioli & Pinzauti 2008). Given the importance of organisms locally adapted to pollinate apple trees (*Malus × domestica* Borkh.) (Rosaceae),

and since knowledge of the natural pollinators in México is poor, the main objective of this study was to observe and document the native pollinators in unmanaged apple orchards using artificial nests in Cuauhtémoc and Guerrero Chihuahua, México.

The experiment was established in 2 apple orchards 1) "La Concepción" (28° 23.9' N, 107° 01.58' W, 2194 m asl), and 2) "San Martín" (28° 24.51' N, 107° 03.88' W, 2256 m asl) located in Cuauhtémoc and Guerrero, Chihuahua, México respectively. Habitats surrounding apple orchards were a forest-complex species predominated by hardwood/coniferous woodland species such as pine tree *Pinus* spp. (Pinaceae), juniper *Juniperus* spp. (Cupressaceae), oak *Quercus* spp. (Fagaceae) and a mix of noncrop vegetation. Undergrowth in both orchards was composed primarily of sage brush (*Artemisia* sp. L.; Asteraceae) and grasses (Poaceae), typical plants in the Midwest plains of Chihuahua. Both apple orchards had been without chemical pesticide application for at least 10 years, and sited near water-bodies, a requirement for the establishment of Megachilid pollinators. Distance between orchards was approximately 4 km, and each had an area of 3 to 5 ha with mature 30-35-year-old 'Red Delicious' and 'Golden Delicious' trees.

Nesting wood traps designed to capture *Osmia* spp. were used. These were fabricated by beekeeper Eng. Ruben Rivera-Landeros based on a model developed by U.S. researchers (Bosch & Kemp 2001). Traps were installed 2 m high in the canopy around orchards, and were sampled periodically from 2010 to 2012 in both orchards (Tonietto et al. 2011). Traps were installed in mid-Mar and removed in Oct or Nov each year, and carried to CIAD laboratory for processing. The pupae were extracted, disinfected with sodium hypochlorite 2% and stored at 7 °C. Also bees and wasps were collected weekly with entomological nets from Apr to Jun 2010 and from May to Sep

2011. Specimens collected were stored in 70% ethanol; some were pinned with entomological needles and identified to genus with taxonomic keys (Lavery & Harder 1988; Michener et al. 1994; Michener 2000; Michener 2007).

A great diversity in species was observed in 3 years, mainly due to an extensive variety of weeds and woody plants surrounding the sited where the pollinators obtained their provisions from many flowering plants including dandelion (*Taraxacum officinale* F. H. Wigg.), clover (*Trifolium repens* L.), Mexican prickly-poppy (*Argemone mexicana* L.), and jagged-edge sunflower (*Helianthus laciniatus* A. Gray.). Various native plant species preferred by these bees and some exotic plants, such as Prickly sow thistle (*Soychus asper* (L.) Hill) and weld (*Reseda luteola* L.), among other forbs bloomed from May to Oct (Cane 2001). According to other studies, where there is great plant diversity, there is great bee diversity (Potts et al. 2003; Hendrix et al. 2010; Wojcik 2011).

We found that the pollinator communities inhabiting the canopies of apple trees orchards of Cuauhtémoc and Guerrero, Chihuahua, were highly diverse, with over 12 bee species found both in nests and trapped in weeds in the orchards. The most taxonomically diverse group in this study was the family Megachilidae. Also during 3 years of sampling we found more than 11 wasp species (Tables 1 and 2). In 2010 with entomological nets, we collected 32 hymenopteran species on weeds, and 22 (68.75%) of these were collected from “La Concepción” orchard, where *Anthidium* sp. (Megachilidae), was the main representative, and 10 (31.25%) species were collected from the “San Martín” orchard (Table 1). In 2011, 45 hymenopteran species were collected in “La Concepción”, which consisted of 21 genera, distributed in 8 families; and the main representative s belonged to the Megachilidae family, with 35.56% of the specimens belonging to the genus *Anthidium*. In the “San Martín” orchard only 15 hymenopteran species were found, which were distributed in 7 genera, 4 genera (9 species) belonging to the Megachilidae and 3 genera belong to Halictidae and Vespidae families (Table 1).

In the 20 traps installed in both orchards in 2011, 245 hymenopterans specimens were collected of which 219 (89.34%) were found in “La Concepción” and 26 (10.61%) in “San Martín” (Table 2). Thus, we was found *Osmia lignaria*, *O. integra* and *Osmia* sp.1 pollinators in “La Concepción”, but in “San Martín” we found only *O. lignaria* specimens (Table 2).

Megachile pugnata Say, was the greatest contributor to the differences between the 2 orchards, and it was the only species collected from a green canopy in “La Concepción” in both years (Matteson et al. 2008; Frankie et al. 2009).

Trachusa sp., *T. ridingsii* Cresson and *Augochlorella aurata* Smith are widely found in northern México (Tonietto et al. 2011). Also *Heriades carinatus* Cresson, *Crabro* sp., *Halictus* sp., *Augochlora* sp. and *Anthidium* sp., *Xylocopa* sp. and *Bombus* spp. were each represented by one specimen (Table 1) (Greenleaf et al. 2007; Tonietto et al. 2011). In this study, an increase in abundance of species that nest in cavities was observed, over those that nest in other nesting types during the 3 years, and similar results were obtained by Matteson et al. (2008) and Dalmazzo (2010).

Other specimens collected from weed flowers with entomological nets were: *Pepsis aciculata* Taschenberg, *Sphex lucae* Saussure, *Dipogon subintermedius* Magretti (recorded in México for first time), *Triscolia ardens* Smith, *Ammophila* spp., *Vespula* sp., *Bicyrtes* sp., and *Lasioglossum* sp. The latter 5 species commonly occur in habitats in apple orchards. Over 29% of the species were represented by a single specimen, which was a similar proportion to those found in other studies (Cane 2001; Tonietto et al. 2011).

Our results are in agreement with earlier surveys in apple orchards of North America, which documented great diversity of bees (Sheffield et al. 2003; Sheffield et al. 2008; Sheffield et al. 2012) and of wasps (Sheffield et al. 2008; Mates et al. 2012). Abundance and diversity of native bees in agricultural systems are positively correlated with proximity and extent of the surrounding habitat (Steffan-Dewenter 2003; Kremen et al. 2004; Taki et al. 2007; Sobek et al. 2009; Hagen & Kraemer 2010). However anthropogenic habitats can be of high value to native bees as evidenced Winfree et al. (2007) and Tonietto et al. (2011) and that the proximity to more natural habitats, as field margins plentiful in floral resources, generally increases bee and wasp diversity (Sheffield et al. 2008; Hagen & Kraemer 2010; Roulston & Goodell 2011; Watson et al. 2011). The pollinator bees associated with crops grown over wide geographic ranges can show remarkable differences from region to region, and each region may have a particular profile of diversity and abundance of species (Michener 2007). In this respect, Neumann & Carreck (2010); Murray et al. (2009); Winfree et al. (2009); Tonietto et al. (2011) mentioned that the abundance of pollinators in the ecosystem is strongly affected by various biotic factors including resource availability (fragmentation and loss) and abiotic factors including loss of natural habitats, human-dominated green spaces and other land-use changes. Restoration of the natural populations of pollinators requires reversal of these changes (Dixon 2009; Winfree 2010; Tonietto et al. 2011), since remnant natural habitats are often insufficient to conserve biodiversity (Rosenzweig 2003). Nevertheless, native bee communities can persist in some anthropogenic

TABLE 1. HYMENOPTERA COLLECTED WITH ENTOMOLOGICAL NETS IN TWO APPLE ORCHARDS IN CUAUHTÉMOC, CHIHUAHUA IN 2010-2011.

Orchard	Year	Family	Genera	Species	Number of Specimens
“La Concepción”	2010	Megachilidae	<i>Trachusa</i>	sp.1	6
				<i>ridingsii</i>	3
				<i>Megachile</i>	3
				<i>Anthidium</i>	8
				<i>Augochlorella</i>	2
	2011	Megachilidae	<i>Trachusa</i>	sp.1	5
				<i>Heriades</i>	3
				<i>Megachile</i>	4
				<i>Anthidium</i>	1
				<i>Osmia</i>	3
		Halictidae	<i>Augochlorella</i>	<i>aurata</i>	1
				<i>Halictus</i>	1
				<i>Augochlora</i>	1
		Crabronidae	<i>Crabro</i>	sp.	2
				sp.	1
		Apidae	<i>Xylocopa</i>	sp.	1
				<i>Bombus</i>	1
		Vespidae	<i>Parancistrocerus</i>	<i>pedestris</i>	3
				<i>Ancistrocerus</i>	2
				<i>Vespula</i>	sp.
		Sphecidae	<i>Chlorion</i>	sp.	2
				<i>Sphex</i>	lucae
				<i>Ammophila</i>	sp.
				<i>Bicyrtes</i>	sp.
		Pompilidae	<i>Pepsis</i>	<i>aciculata</i>	1
				<i>Dipogon</i>	subintermedius
		Scoliidae	<i>Triscolia</i>	<i>ardens</i>	2
“San Martín”	2010	Halictidae	<i>Augochlora</i>	sp. ^p	2
				<i>Lasioglossum</i>	sp.
		Megachilidae	<i>Trachusa</i>	sp.1	2
				<i>Anthidium</i>	sp.
		Crabronidae	<i>Crabro</i>	sp.	2
				sp.	2
	2011	Megachilidae	<i>Trachusa</i>	sp.1	4
				<i>Heriades</i>	carinatus
				<i>Megachile</i>	sp.
				<i>Anthidium</i>	sp.
		Halictidae	<i>Augochlora</i>	sp.	1
		Vespidae	<i>Ancistrocerus</i>	<i>tuberculocephalus</i>	3
				<i>Vespula</i>	sp.

habitats (Cane 2001; Marlin & LaBerge 2001). Rosenzweig (2003) and Tonietto et al. (2011) noted that the return to organic management provides suitable habitats for native species and thus supports native biodiversity

We extend heartfelt thanks to Fundación PRODUCE Chihuahua, for funding the project “Collecta e identificación de *Osmia* spp. y otros géneros polinizadores para el manzano”, with registration number 70035. To Eng. Alejandro Berlanga and

Mr. Martín Chaparro, for allowing us to conduct this research in their apple tree orchards.

SUMMARY

Research on the diversity and abundance of taxa pollinators at the community level are essential to know aspects of their biology, ethology, ecology and the impacts on reproduction and plant diversity. Bees have an important role in providing pollination, which is essential to plant reproduc-

TABLE 2. HYMENOPTERA FOUND IN NESTS OF OSMIA SPP. IN TWO APPLE ORCHARDS IN 2011-2012.

Orchard	Year	Family	Genera	Species	Type of Nest	Number of Specimens
“La concepción”	2011	Megachilidae	<i>Osmia</i>	<i>lignaria</i>	C	6
				<i>integra</i>	C	3
				sp.1	C	180
		Vespidae	<i>Megachile</i>	<i>pugnata</i>	C	8
				<i>pedestris</i>	C-SW	6
				<i>tuberculocephalus</i>	C-SW	12
		Sphecidae	<i>Chlorion</i>	sp.	S	4
	2012	Megachilidae	<i>Osmia</i>	sp.1	C	143
				sp.2	C	20
				sp.	C	4
		Vespidae	<i>Ancistrocerus</i>	<i>tuberculocephalus</i>	C-SW	84
		Sphecidae	<i>Chlorion</i>	sp.	S	2
“San Martín”	2011	Megachilidae	<i>Osmia</i>	<i>lignaria</i>	C	5
				<i>pugnata</i>	C	12
		Vespidae	<i>Ancistrocerus</i>	<i>tuberculocephalus</i>	C-SW	9
	2012	Megachilidae	<i>Osmia</i>	sp.2	C	64
				<i>pugnata</i>	C	69
				sp.	C	20
		Vespidae	<i>Ancistrocerus</i>	<i>tuberculocephalus</i>	C-SW	47

Nesting type (nest location): S, soil; C, cavity; SW, soft wood; according to Tonietto et al. (2011).

tion. The objective of the study was to document the diversity and abundance of bee pollinators, with emphasis in Megachilid species, in 2 unman- aged apple orchards, located in Cuauhtémoc and Guerrero, Chihuahua, México. Both orchards had similar conditions, with intermediate levels of ad- jacent natural/semi-natural habitat. In 2010, 2011 and 2012, special wood nesting traps for capturing *Osmia* spp. were installed. Four species of solitary pollinator bees of *Osmia* genus were captured, of which the most represented was *Osmia* sp.1, with 323 specimens, followed by *Osmia* sp.2 with 84 specimens, *O. lignaria* Say with 11 specimens and 3 specimens of *O. integra* Cresson. Also, other spe- cies of bees and wasps visitors of plants with flow- ers were found such as *Trachusa* sp., *Anthidium* spp., *Megachile pugnata* Say, *Heriades carinatus* Cresson, *Xylocopa* sp., *Bombus* sp., *Ancistrocerus tuberculocephalus* (Saussure) and *Vespula* sp.

Key Words: pollination, insect pollinators, blue orchard mason bee, Megachilid, *Osmia lignaria*, *Osmia integra*

RESUMEN

Las investigaciones sobre la diversidad y abun- dancia de taxas polinizadoras a nivel comunidad son esenciales para conocer aspectos relacionados con su biología, ecología, etología y los impactos sobre la reproducción y diversidad de plantas. Por lo que, mantener la relación entre plantas y polini- zadores es vital para la estabilidad de los ecosiste-

mas y agroecosistemas donde las abejas juegan un rol muy importante para la polinización, la cual es esencial para la reproducción de las plantas. El ob- jetivo del estudio fue estimar la diversidad y abun- dancia de abejas polinizadoras, con énfasis en espe- cies de Megaquílidos, en dos huertos de manzano, sin manejo, localizados en Cuauhtémoc y Guerrero, Chihuahua, México. Ambos huertos muestran con- diciones similares, con niveles intermedios de há- bitats adyacentes naturales y semi-naturales. En 2010, 2011 y 2012, se colocaron trampas especiales de madera para la anidación de polinizadores para la captura de *Osmia* spp. Se encontraron cuatro especies de abejas solitarias polinizadoras pertene- cientes al género *Osmia*, donde *Osmia* sp.1 fue la más representada, con 323 especímenes, 84 especí- menes de *Osmia* sp.2, 11 especímenes de *O. lignaria* Say y tres de *O. integra* Cresson. Se encontraron otras especies de abejas y avispa visitantes de plantas con flores tales como *Trachusa* sp., *Anthi- dium* spp., *Megachile pugnata* Say, *Heriades cari- natus* Cresson, *Xylocopa* sp., *Bombus* sp., *Ancistro- cerus tuberculocephalus* (Saussure) and *Vespula* sp.

Palabras Clave: Polinización, insectos polini- zadores, abeja albañil, Megaquílidos, *Osmia lig- naria*, *Osmia integra*

REFERENCES CITED

BIESMEIJER, J. C., ROBERTS, S. P. M., REEMER, M., OHLEMÜLLER, R., EDWARDS, M., PEETERS, T., SCHAFFERS, A. P., POTTS, S. G., KLEUKERS, R.,

- THOMAS, C. D., SETTELE, J., AND KUNIN, W. E. 2006. Parallel declines in pollinators and insect pollinated plants in Britain and the Netherlands. *Science* 313: 351-354.
- BOSCH, J., AND KEMP, W. P. 2001. How to manage the blue orchard bee as an orchard pollinator. Sustainable Agriculture Network, National Agricultural Library; Beltsville, MD, USA. 88 pp.
- CANE, J. H. 2001. Habitat fragmentation and native bees: A premature verdict? *Conserv. Ecol.* 5(1): 3.
- CANE, J. H. 2008. Bees (Hymenoptera: Apoidea: Apiformes), pp. 419-433 *In* J. L. Capinera [ed.], *Encyclopedia of Entomology* 2nd edition, Springer. 4346 pp.
- DALMAZZO, M. 2010. Diversidad y aspectos biológicos de abejas silvestres de un ambiente urbano y otro natural de la región central de Santa Fe, Argentina. *Rev. Soc. Entomol. Argentina* 69(1-2): 33-4.
- DAS, B., AHMAD, N., SRIVASTAVA, K. K., AND RANJAN, P. 2011. Top working method and bloom density of pollinizers as productive determinant for spur type apple (*Malus x domestica* Borkh.) cultivars. *Sci. Hort.* 129: 642-648.
- DIXON, K. 2009. Pollination and restoration. *Science* 325: 571-573.
- DRUMMOND, F. A., AND STUBBS, C. S. 1997. Potential for management of the blueberry bee, *Osmia atriventris* Cresson. *Acta Hort.* (ISHS) 446:77-86.
- FELICOLI, A., AND PINZAUTI, M. 2008. Pollination by *Osmia* Bees (Hymenoptera: Megachilidae), pp. 2971-2978 *In* J. L. Capinera [ed.], *Encyclopedia of Entomology* 2nd edition, Springer. 4346 p.
- FRANKIE, G., THORP, R. W., HERNANDEZ, J., RIZZARDI, M., ERTTER, B., PAWELEK, J. C., WITT, S. L., SCHINDLER, M., COVILLE, R., AND WOJC, V. A. 2009. Native bees are a rich natural resource in urban California gardens. *California Agric.* 63(3): 113-120.
- GALLAI, N., SALLES, J. M., SETTELE, J., AND VAISSIÈRE, B. E. 2009. Economic valuation of the vulnerability of world agriculture confronted with pollinator decline. *Ecol. Econ.* 68: 810-821.
- GREENLEAF, S. S., WILLIAMS, M., WINFREE R., KREMEN C. 2007. Bee foraging ranges and their relationship to body size. *Oecologia* 153: 589-596.
- HAGEN, M., AND KRAEMER, M. 2010. Agricultural surroundings support flower-visitor networks in an Afro-tropical rain forest. *Biol. Conserv.* 143: 1654-1663.
- HENDRIX, S. D., KWAISER, K. S., AND HEARD, S. B. 2010. Bee communities (Hymenoptera: Apoidea) of small Iowa hill prairies are as diverse and rich as those of large prairie preserves. *Biodivers. Conserv.* 19: 1699-1709.
- KLEIN, A. M., VAISSIÈRE, B. E., CANE, J. H., STEFFAN-DEWENTER, I., CUNNINGHAM, S. A., KREMEN, C., AND TSCHARNTKE, T. 2007. Importance of pollinators in changing landscapes for world crops. *Proc. R. Soc. B: Biol. Sci.* 274: 303-313.
- KRAEMER, M. E., AND FAVI, F. D. 2005. Flower phenology and pollen choice of *Osmia lignaria* (Hymenoptera: Megachilidae) in central Virginia. *Environ. Entomol.* 34(6): 1593-1605.
- KREMEN, C., WILLIAMS, N. M., BUGG, R. L., FAY, J. P., AND THORP, R. W. 2004. The area requirements of an ecosystem service: crop pollination by native bee communities in California. *Ecol. Lett.* 7: 1109-1119.
- LAVERTY, T. M., AND HARDER, L. D. 1988. The bumble bees of eastern Canada. *Canadian Entomol.* 120: 965-987.
- MARLIN, J. C., AND LABERGE, W. E. 2001. The native bee fauna of Carlinville, Illinois, revisited after 75 years: A case for persistence. *Conserv. Ecol.* 5(1): U91-U116.
- MATES, S. G., PERFECTO, I., AND BADGLEY, C. 2012. Parasitoid wasp diversity in apple orchards along a pest-management gradient. *Agric. Ecosys. Environ.* 156: 82-88.
- MATTESON, K. C., ASCHER, J. S., AND LANGELOTO, G. A. 2008. Bee richness and abundance in New York City urban gardens. *Ann. Entomol. Soc. America* 101: 140-150.
- MICHENER, C. D. 2000. The bees of the world. Baltimore and London: The Johns Hopkins University Press. 913 pp.
- MICHENER, C. D. 2007. The bees of the world, 2nd edition, Johns Hopkins University Press, Baltimore, Maryland, USA. 992 pp.
- MICHENER, C. D., MCGINLEY, R. J., AND DANFORTH, B. N. 1994. The bee genera of North and Central America (Hymenoptera: Apoidea). Washington, DC: The Smithsonian Institution. 209 pp.
- MURRAY, T. E., KUHLMANN, M., AND POTTS, S. G. 2009. Conservation ecology of bees: Populations, species, and communities. *Apidologie* 40: 211-236.
- NEUMANN, P., AND CARRECK, N. L. 2010. Honey bee colony loss. *J. Apiculture Res.* 49:1-6.
- POTTS, S. G., VULLIANY, N., DAFNI, A., NÉEMAN, G., AND WILMER, P. 2003. Linking bees and flowers: How do floral communities structure pollinator communities? *Ecology* 84: 2628-2642.
- ROSENZWEIG, M. L. 2003. Reconciliation ecology and the future of species diversity. *Oryx* 37(2): 194-205.
- ROULSTON, T. H., AND GOODELL, K. 2011. The role of resources and risks in regulating wild bee populations. *Annu. Rev. Entomol.* 56: 293-312.
- SHEFFIELD, C. S., KEVAN, P. G., SMITH, R. F., RIGBY, S. M., AND ROGERS, R. E. L. 2003. Bee species of Nova Scotia, Canada, with new records and notes on bionomics and floral relations (Hymenoptera: Apoidea). *J. Kansas Entomol. Soc.* 76: 357-384.
- SHEFFIELD, C. S., KEVAN, P. G., WESTBY, S. M., AND SMITH, R. F. 2008. Diversity of cavity-nesting bees (Hymenoptera: Apoidea) within apple orchards and wild habitats in the Annapolis Valley, Nova Scotia, Canada. *Canadian Entomol.* 140: 235-249.
- SHEFFIELD, C. S., KEVAN, P. P., PINDAR, A., AND PACKER, L. 2012. Bee (Hymenoptera: Apoidea) diversity within apple orchards and old fields in the Annapolis Valley, Nova Scotia, Canada. *Canadian Entomol.* 145: 1-21.
- SOBEK S., TSCHARNTKE T., SCHERBER C., SCHIELE S., STEFFAN-DEWENTER, I. 2009. Canopy vs. understory: Does tree diversity affect bee and wasp communities and their natural enemies across forest strata? *Forest Ecol. Mgt.* 258: 609-615.
- STEFFAN-DEWENTER, I. 2003. Importance of habitat area and landscape context for species richness of bees and wasps in fragmented orchard meadows. *Conserv. Biol.* 17: 1036-1044.
- TAKI, H., KEVAN, P. G., AND ASCHER, J. S. 2007. Landscape effects of forest loss in a pollination system. *Landscape Ecol.* 22: 1575-1587.
- TONIETTO, R., FANT, J., ASCHER, J., ELLIS, K., AND LARKIN, D. 2011. A comparison of bee communities of Chicago green roofs, parks and prairies. *Landscape Urban Planning* 103: 102-108.

- WATSON, J. C., WOLF, A. T., AND ASCHER, J. S. 2011. Forested landscapes promote richness and abundance of native bees (Hymenoptera: Apoidea: Anthophila) in Wisconsin apple orchards. *Environ. Entomol.* 40: 621-632.
- WILLIAMS, N. M., CRONE, E. E., ROULSTON, T. H., MINCKLEY, R. L., PACKER, L., AND POTTS, S. G. 2010. Ecological and life-history traits predict bee species responses to environmental disturbances. *Biol. Conserv.* 143(10): 2280-2291.
- WINFREE, R. 2010. The conservation and restoration of wild bees. *Ann. N. Y. Acad. Sci.* 1195: 169-197.
- WINFREE, R., AGUILAR, R., VASQUEZ, D. P., LEBUHN, G., AND AIZEN, M. A. 2009. A meta-analysis of bees' responses to anthropogenic disturbance. *Ecology*. 90(8): 2068-2076.
- WINFREE, R., GRISWOLD, T., AND KREMEN, C. 2007. Effect of human disturbance on bee communities in a forested ecosystem. *Conserv. Biol.* 21(1): 213-223.
- WOJCIK, V. 2011. Resource abundance and distribution drive bee visitation within developing tropical urban landscapes. *J. Pollination Ecol.* 4(7): 48-56.