

## **First Report of *Neosilba pradoi* and *Dasiops frieseni* (Diptera: Lonchaeidae) in Cultivated and Wild Hosts in Uruguay**

Authors: Calvo, María Victoria, Delgado, Soledad, Scatoni, Iris, and Garcia, Flavio Roberto Mello

Source: Florida Entomologist, 100(4) : 831-832

Published By: Florida Entomological Society

URL: <https://doi.org/10.1653/024.100.0427>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

# First report of *Neosilba pradoi* and *Dasiops friesei* (Diptera: Lonchaeidae) in cultivated and wild hosts in Uruguay

María Victoria Calvo<sup>1\*</sup>, Soledad Delgado<sup>1</sup>, Iris Scatoni<sup>1</sup>, and Flavio Roberto Mello Garcia<sup>2</sup>

The members of the family Lonchaeidae (lance flies) are glossy black color with a blue or green metallic shine, or a mixture of both blue and green. The larvae usually are saprophagous or frugivorous (Bentancourt et al. 2009). This family is very well represented in South America, although knowledge of the frugivorous Lonchaeidae is just being developed; presently, little or nothing is known about the identity and biology of the many South American species (Korytkoioski & Ojeda 1971; Norrbom & McAlpine 1997; Strikis et al. 2011; Gislotti et al. 2017).

The family Lonchaeidae possibly has received little attention because several species are opportunists associated with fruit fly (Diptera: Tephritidae) infestations (McAlpine & Steyskal 1982), rather than being primary pests. However, Araujo & Zucchi (2002), Aguiar-Menezes et al. (2004), Strikis & Lerena (2009), and Nicácio & Uchôa (2011) found that some species were primary invaders, and not dependent on previous oviposition by tephritids.

In Uruguay, there have been few reports of Lonchaeidae associated with fruit. *Dasiops uruguayensis*, described by Enderlein in 1936 (Korytkoioski & Ojeda 1971), and *Lonchaea chalybea* Wiedemann are the only records for this country (Ruffinelli & Carbonell 1954).

Between Nov 2013 and May 2014, mature fruits were collected from potential hosts of Diptera in Salto (31.3865°S, 57.7176°W) and Canelones (34.6207°S, 56.3612°W), Uruguay. Fruits (from plants or recently fallen) were collected from 18 plant species (Table 1). The samples were counted, weighed, and stored individually in screen-covered plastic pots containing

Table 1. Species of plants sampled, fruit weight, number of fruits sampled, number of adult Lonchaeidae emerged, average number of adults per kg, and percentage of fruit infested.

Plant taxa	Fruit weight (g)	No. fruit sampled	No. lonchaeids emerged	No. adults per kg	% fruit infested
Myrtaceae					
<i>Psidium cattleianum</i>	1396.0	145	3	2	2
<i>Myrcianthes pungens</i>	655.4	345			
<i>Psidium guajaba</i>	7986.9	195			
<i>Acca sellowiana</i>	10270.7	373			
<i>Eugenia uniflora</i>	162.6	8	1	6	13
<i>Hexachlamis edulis</i>	449.2	25			
Ebenaceae					
<i>Diospyros kaki</i>	2922.3	46			
Moraceae					
<i>Ficus carica</i>	2147.6	40			
<i>Maclura pomifera</i>	6359.0	21			
Rutaceae					
<i>Fortunella margaritus</i>	192.2	20			
<i>C. reticulata</i> × <i>C. sinensis</i>	13227.2	119			
<i>Citrus sinensis</i> cv. W Navel	14721.6	82			
<i>Citrus sinensis</i> cv. Valencia	3888.4	30	27	7	17
<i>Citrus paradisi</i>	9364.0	39			
Sapotaceae					
<i>Pouteria gardneriana</i>	791.6	56			
Passifloraceae					
<i>Passiflora caerulea</i>	33.1	2	13	393	100
Rosaceae					
<i>Pyrus communis</i>	27001.5	181	1	0.04	0.6
Santalaceae					
<i>Acanthosyris spinescens</i> *	3015.3	308	2	0.001	0.6

\*First record of a lonchaeid infesting this plant.

<sup>1</sup>University of República Oriental del Uruguay, Department of Entomology, Montevideo, 12900, Uruguay; E-mail: vcalvo@fagro.edu.uy; soledaddelgadojorge@hotmail.com; iscatoni@fagro.edu.uy

<sup>2</sup>Federal University of Pelotas, Institute of Biology, Department of Zoology and Genetics, Laboratory of Insect Ecology 96010-900, P.O. Box 354, Pelotas, Rio Grande do Sul, Brazil; E-mail: flavio.garcia@pq.cnpq.br

\*Corresponding author; E-mail: vcalvo@fagro.edu.uy

**Table 2.** Numbers of adults of *Neosilba pradoi* and *Dasiops frieseni* by host, average number of adults per kg, and percent of fruit infested.

Plant Species	<i>Neosilba pradoi</i>			<i>Dasiops frieseni</i>		
	No. adults	No. adults per kg	% fruit infested	No. adults	No. adults per kg	% fruit infested
<i>Citrus sinensis</i> cv. Valencia	1	0.3	3.3	26*	6.7	6.7
<i>Passiflora caerulea</i>	12	362.5	100	—	—	—
<i>Eugenia uniflora</i>	1	6.2	12.5	—	—	—

\*First record of host for the *D. frieseni*.

sterile sand, and kept at 25 °C. The emerged adults were preserved in 70% ethanol for identification.

A total of 47 specimens of Lonchaeidae were obtained from 6 plant species. Forty specimens were identified as *Neosilba pradoi* Strikis & Lerena 2009 (Lonchaeinae) and *Dasiops frieseni* Norrbom & McAlpine, 1997 (Dasiopinae). The rest were identified only to family level. These two species are recorded for first time in Uruguay (Table 2). A total of 5 males and 9 females of *N. pradoi* emerged from fruits collected from plants, and 14 males and 12 females of *D. frieseni* from fruits collected from both the plants and on the soil beneath the trees. In most fruits (85% of those infested) a single species emerged, but in one fruit of *Psidium cattleianum* Sabine (Myrtaceae) and one of *Citrus sinensis* (L.) (Rutaceae), adults of *Anastrepha fraterculus* (Wiedemann) and *Ceratitis capitata* (Wiedemann) (both Diptera: Tephritidae) emerged along with the lonchaeids, respectively. As noted previously, some species of Lonchaeidae should be considered to be of economic importance and regarded as primary invaders. Though sometimes associated with tephritid infestation, lonchaeids seem to be independently capable of attacking fruit (Uchôa 2012).

In Brazil, *N. pradoi* has been reported to occur in *Passiflora caerulea* L. (Passifloraceae), though Marsaro et al. (2012) found lower levels of infestation as compared to us (28.4 larvae per kg of fruit infested by *N. pradoi* or *Lonchaea* sp.). Moreover, *Eugenia uniflora* (L.) (Myrtaceae) and *Citrus sinensis* were described as hosts by Garcia & Norrbom (2011). They found infestation levels (2.7 adults per kg and 4.0 adults per kg, respectively) similar to what we observed, but this is the first time that *N. pradoi* has been found in the orange cv. Valencia. Aguiar-Menezes et al. (2004) found that *D. frieseni* had the highest infestation index (considering lonchaeids and tephritids) infesting *Passifloraceae* spp. in southeastern Brazil.

This study is the first report of these species for Uruguay and it is the most southern detection of them, and thus can aid in the determination of the species' distribution and ecology. In addition, this is the first record of *Acanthosyris spinescens* Griseb. (Santalaceae) as host for a lonchaeid. More extensive and intensive surveys of lonchaeid hosts should be conducted in Uruguay to improve our knowledge of the diversity and ecology of these important dipteran species, which could have important effects on fruit production.

We thank Dr. Pedro Strikis (University of São Paulo, Brazil) for identifying the insects, Wuilliam Techeira (MGAP Salto), Emily Silva Araujo (Federal University of Pelotas) and Facultad de Agronomía (Salto) for helping during fruit collection. We also thank Capes-UdelaR and Agencia Nacional de Investigación e Innovación for financial support.

Summary

We report *Neosilba pradoi* and *Dasiops frieseni* (Diptera: Lonchaeidae) for the first time in Uruguay, which is the most southern distribution yet found for these species. Moreover, new host associations are recorded: *D. frieseni* in *Citrus sinensis* cv. Valencia and an undetermined lonchaeid in *Acanthosyris spinescens*.

Key Words: lance flies; *Eugenia uniflora*; *Passiflora caerulea*; *Citrus sinensis*

Sumario

Se reportan por primera vez para Uruguay *Neosilba pradoi* y *Dasiops frieseni* (Diptera: Lonchaeidae), siendo la distribución más austral. Además, se registran nuevas asociaciones de hospederos: *Citrus sinensis* cv. Valencia para *D. frieseni* y *Acanthosyris spinescens* para una especie indeterminada de Lonchaeidae.

Palabras Clave: lonqueídios; *Eugenia uniflora*; *Passiflora caerulea*; *Citrus sinensis*

References Cited

Aguiar-Menezes EN, Nascimento RB, Menezes EB. 2004. Diversity of fly species (Diptera: Tephritoidea) from *Passiflora* spp. and their hymenopterous parasitoids in two municipalities of the southeastern Brazil. *Neotropical Entomology* 33: 113–116.

Araújo EL, Zucchi RA. 2002. Hospedeiros e níveis de infestação de *Neosilba pendula* (Bezzi) (Diptera: Lonchaeidae) na região de Mossoró/Assu, RN. *Arquivos do Instituto Biológico* 69: 913–94.

Bentancourt CM, Morelli E, Scatoni IB. 2009. Insectos del Uruguay. Hemisferio Sur, Montevideo, Uruguay.

Garcia FRM, Norrbom AL. 2011. Tephritoid flies (Diptera, Tephritoidea) and their plant hosts from the state of Santa Catarina in Southern Brazil. *Florida Entomologist*, 94: 151–157.

Gislotti L, Uchoa MA, Prado A. 2017. New records of fruit trees as host for *Neosilba* species (Diptera, Lonchaeidae) in southeast Brazil. *Biota Neotropica* 17: e20160213. <http://dx.doi.org/10.1590/1676-0611-BN-2016-0213>

Korytkowski C, Ojeda D. 1971. Revisión de las especies de la familia Lonchaeidae en el Perú (Diptera: Acalypratae). *Revista Peruana de Entomología* 14: 87–116.

Marsaro ALJ, Strikis PC, Savaris M, Lampert S, Adaime R. 2012. New host records of Lonchaeidae (Diptera) in Brazil and associated parasitoid. *Revista de Agricultura* 87: 116–118.

McAlpine JF, Steyskal GC. 1982. A revision of *Neosilba* McAlpine with a key to the world genera of Lonchaeidae (Diptera). *The Canadian Entomologist* 114: 105–137.

Nicácio J, Uchoa MA. 2011. Diversity of frugivorous flies (Diptera: Tephritidae and Lonchaeidae) and their relationship with host plants (Angiospermae) in environments of South Pantanal region, Brazil. *Florida Entomologist* 94: 443–466.

Norrbom AL, McAlpine JF. 1997. A revision of the Neotropical species of *Dasiops* Rondani (Diptera: Lonchaeidae) attacking *Passiflora* (Passifloraceae). *Memoirs of the Entomological Society of Washington* 18: 189–211.

Ruffinelli A, Carbonell CS. 1954. Segunda lista de insectos y otros artrópodos de importancia económica en el Uruguay. *Revista de la Asociación de Ingenieros Agrónomos* 94: 33–82.

Strikis PC, Lerena MLM. 2009. A new species of *Neosilba* (Diptera, Lonchaeidae) from Brazil. *Iheringia, Serie Zoologica* 99: 273–275.

Strikis PC, Deus EG, Silva RA, Pereira JDB, Jesus CR, Marsaro Júnior AL. 2011. Conhecimento sobre Lonchaeidae na Amazônia brasileira, pp. 205–216 *In* Silva RA, Lemos WP, Zucchi RA [Eds.], Moscas-das-frutas na Amazônia brasileira: diversidade, hospedeiros e inimigos naturais. Embrapa Amapá, Macapá, Amapá, Brazil.

Uchôa MA. 2012. Fruit flies (Diptera: Tephritoidea): biology, host plants, natural enemies, and the implications to their natural control, pp. 271–300 *In* Soloneski, S, Larramendy, ML [Eds], Integrated Pest Management and Pest Control: Current and Future Tactics. Tech, Rijek, Croatia.