



First Record of *Blattella vaga* (Blattodea: Ectobiidae) from Southern Alabama

Authors: Jeon, Sunghun Alan, Balusu, Rammohan R., Zhang, Lee, Fadamiro, Henry Y., and Appel, Arthur G.

Source: Florida Entomologist, 101(1) : 109-112

Published By: Florida Entomological Society

URL: <https://doi.org/10.1653/024.101.0119>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

First record of *Blattella vaga* (Blattodea: Ectobiidae) from southern Alabama

Sunghun Alan Jeon¹, Rammohan R. Balusu¹, Lee Zhang¹, Henry Y. Fadamiro¹, and Arthur G. Appel^{1,*}

Abstract

The field cockroach, *Blattella vaga* Hebard (Blattodea: Ectobiidae), is native to central Asia including Afghanistan, India, Iran, Pakistan, and Sri Lanka. It was described first in 1935; however, from specimens collected in Arizona and California. Since then, the distribution of *B. vaga* has slowly increased along the southern United States and Mexican border, apparently following major interstate highways. We report the first record of *B. vaga* from Mobile, Alabama, and suggest that this species will spread to Florida and possibly northward into Georgia and South Carolina. The identification was confirmed using morphological, chemical, and molecular methods. We suggest that when possible, multiple independent methods should be used to confirm species identifications.

Key Words: Distribution; cuticular hydrocarbon profile; mitochondrial gene sequence

Resumen

La cucaracha de campo, *Blattella vaga* Hebard (Blattodea: Ectobiidae), proviene de Asia central, incluyendo Afganistán, India, Irán, Pakistán y Sri Lanka. Sin embargo, fue descrita por primera vez en 1935 a partir de especímenes recolectados en Arizona y California. Desde entonces, la distribución de *B. vaga* ha extendido lentamente a lo largo del sur de los Estados Unidos y México, aparentemente siguiendo las principales carreteras interestatales. Reportamos el primer registro de *B. vaga* en la ciudad de Mobile, Alabama y sugerimos que esta especie se extenderá a la Florida y posiblemente hacia el norte a Georgia y Carolina del Sur. La identificación se confirmó utilizando métodos morfológicos, químicos y moleculares. Sugerimos que, cuando sea posible, se utilicen múltiples métodos independientes para confirmar las identificaciones de especies.

Palabras Clave: Distribución; perfil de hidrocarburo cuticular; secuencia del gen mitocondrial

The field cockroach, *Blattella vaga* Hebard (Blattodea: Ectobiidae), was described in 1935 by Morgan Hebard from specimens collected in Arizona and California (Hebard 1935). This species resembles the other established *Blattella* spp. in North America: the Asian cockroach, *B. asahinai* Mizukubo, and the German cockroach, *B. germanica* (L.) (Atkinson et al. 1991; Appel 1995). Adults of these species are similar in length, with a pair of longitudinal stripes on their pronotum. Both *B. asahinai* and *B. vaga* live outdoors, fly to lights, and can become peridomestic pests (Helfer 1987; Atkinson et al. 1991). In contrast, *B. germanica* is almost exclusively (but see Appel & Tucker 1986) a domiciliary pest in apartments, homes, and food preparation areas (Ebeling 1978). Unlike *B. asahinai* and *B. vaga*, *B. germanica* does not fly and is repelled by light (Ebeling et al. 1966).

Since its description in 1935 from specimens collected in 1933, *B. vaga* has been reported periodically from the southern tier of the United States (Hogue 1993; Drees & Jackman 1998). Atkinson et al. (1991) reported the distribution of *B. vaga* to include the southern regions of the contiguous states of California, Nevada, New Mexico, Arizona, Texas, and Louisiana. There have been no reports of *B. vaga* from Alabama or Mississippi (Dakin & Hays 1970; Pratt 1988; GBIF Secretariat 2017). The distribution pattern of this species appears to overlap major interstate highways such as I-10 (east from southern California through Arizona, New Mexico, Texas, and Louisiana) and I-40 (east from south-

ern California through Arizona, New Mexico, and Texas) (GBIF Secretariat 2017). Because there are many distribution records along I-10, it would not be surprising to find *B. vaga* near that interstate considering the proximity of its eastern-most record in Louisiana to Mobile, Alabama. Austin et al. (2007) reported infestations of the closely related *B. asahinai* in Texas also near major highways. Similarly, Snoddy and Appel (2008) concluded that distributions of *B. asahinai* in Alabama and Georgia followed major interstates northward from Florida.

The objective of this study was to determine the identity of cockroaches from Alabama that we hypothesized could be *B. vaga*, and a range extension for this species. We used several independent methods to determine the identification of this species and offer several predictions of new distributions.

Materials and Methods

SPECIMEN COLLECTION AND REARING

Cockroaches were collected on 15 Mar 2016 near Cedar Point Fishing Pier, Mobile County, Mobile, Alabama (30.3103°N, 88.1383°W). A total of 11 specimens (third instar through terminal instar) were collected by hand from under wood and in surrounding grass; the soil was a moist

¹Auburn University, Department of Entomology and Plant Pathology, Auburn, Alabama 36849-5413, USA; E-mail: szj0029@tigermail.auburn.edu (S. A. J.), balusrr@auburn.edu (R. R. B.), zhangl1@auburn.edu (L. Z.), fadamhy@auburn.edu (H. Y. F.), appelag@auburn.edu (A. G. A.)

*Corresponding author; E-mail: appelag@auburn.edu

dark clay. Specimens were reared at room temperature and ambient photoperiod in a 0.47 L jar with moist Eco Earth™ (Zoo Med Laboratories, Inc., San Luis Obispo, California, USA) shredded coconut fiber substrate and leaves from various deciduous trees. Cockroaches were fed dry Purina Dog Chow® (Nestlé Purina, St. Louis, Missouri, USA) and pieces of fresh carrot. The jar was lightly misted with water weekly.

A colony of *B. vaga* that had been obtained from the University of California, Riverside in 1985 and reared continuously at Auburn University was used to compare with wild-caught specimens. The Auburn colony was reared in 3.8 L glass jars at 28 ± 2 °C, 40–55% RH, and a photoperiod of 12:12 h (L:D). Colonies were provided corrugated cardboard for cockroach harborage, dry dog food, and clean water. Colonies were fed, supplied water, and cleaned weekly.

MORPHOLOGICAL EXAMINATION

Adult male specimens were killed by a brief exposure to hydrogen cyanide gas. Tergal glands and genitalia were prepared for examination as described by Roth (1985). The abdomens were removed from the rest of the body briefly and placed in 10% KOH at 37 °C for 24 h to loosen and remove internal tissues. Abdomens then were washed in water, dehydrated in a series of increasingly concentrated ethyl alcohol solutions, and cleared in xylol. Abdomens were dissected to remove the tergites with tergal glands and to separate the genitalia. Tergal glands, subgenital and supraanal plates, and the genitalia were mounted on glass slides with Permount™ (Fisher Scientific, Fair Lawn, New Jersey, USA).

Intact male cockroaches and slide-mounted material were examined with a dissecting microscope at 10–100X. Morphological characters were used to follow the key to *Blattella* spp. presented in Roth (1985).

CUTICULAR HYDROCARBON ANALYSIS

Adult male cockroaches were killed by freezing at –20 °C for 2 h. Cuticular hydrocarbons were extracted by immersing a single dead, thawed male cockroach in 0.5 ml of hexane under laboratory conditions for 15 min. Hexane extracts were evaporated to dryness under a gentle stream of nitrogen and re-dissolved in 250 µL of hexane solvent to achieve equivalent concentration of cuticular hydrocarbons among the samples. Extracts then were analyzed using an Agilent 7890A Gas Chromatograph (Agilent Technology, Santa Clara, California, USA) coupled to a 5975C Mass Selective Detector, with an HP-5ms capillary column (30 m × 0.25 mm i.d., 0.25 µm film thickness). Mass spectra were obtained using electron impact (EI, 70 eV). One µL of each extract was injected into the gas chromatograph-mass selective detector in splitless injection mode. The gas chromatograph oven temperature was programmed from 40 °C for 1 min to 280 °C at 20 °C per min, and then the temperature was ramped at 1 °C per min to 320 °C and held for 1 min. The injector and transfer line temperatures were set at 300 and 320 °C, respectively. The patterns of cuticular hydrocarbon components of wild (unknown) specimens were compared with a laboratory colony of *B. vaga* (known) using their mass spectra, retention indices (Kováts index) (NIST 2005 library search software version 2.0, National Institute of Standards and Technology, Gaithersburg, Maryland, USA), and the mass spectra previously reported for *B. vaga* (Carlson & Brenner 1988).

DNA EXTRACTION AND PCR AMPLIFICATION

Blattella vaga DNA was isolated from fresh tissue using EZNA SP Plant DNA kit (OMEGA Bio-tek, Norcross, Georgia, USA) following the manufacturer's protocol. The extraction process included homogenizing several freshly removed legs in 200 µL of lysis buffer (100 mM NaCl,

100 mM EDTA, 100 mM Tris, and 0.5% SDS, at pH 7.0) followed by incubation steps after addition of proteinase K (4 ml of 20 mg per ml) at 55 °C for 3 h followed by RNAase (2 ml of 10 mg per ml) at 37 °C for 20 min. DNA pellets were precipitated in 100% ethanol and stored overnight at –20 °C. Finally, the DNA was washed in 70% ethanol, centrifuged, and vacuum dried. DNA then was suspended in 100 µL of TE buffer (10 mM Tris, pH 8.0, and 1 mM EDTA) and stored at –20 °C.

PCR amplification was performed with a programmable thermal controller (SureCycler 8800, Agilent Technology, Santa Clara, California, USA). Composition of the reaction mixture (50 µL) used for amplification was 28.5 µL water, 5 µL 10X PCR buffer II, 800 mM dNTP mix, 0.8 mM of each primer, 2.5 U AmpliTaq DNA polymerase, 2 µL DNA, and 3 mM MgCl₂.

The central portion of the mitochondrial CO1 gene was amplified using the primer pairs C1-J-1718 (5'-GGAGGATTGGAAATTGATTAGTTC-3') and C1-N-2191 (5'-CCCGGTAAAATTAATATAAATCTC-3') (Pachamuthu et al. 2000). Amplification was performed using the following temperature profile: 94 °C for 2 min, 35 cycles of 94 °C for 45 s, 48 °C for 1 min, and 72 °C for 1.5 min. After a final extension step at 72 °C for 5 min, PCR products were stored at 4 °C.

PCR products were isolated by loading the PCR product onto a 1.0% agarose TAE gel containing ethidium bromide (0.6 mg per ml), and DNA fragments were purified by E.Z.N.A. Gel Extraction Kit (OMEGA Bio-tek, Norcross, Georgia, USA) using the manufacturer's protocol.

Capillary sequencing was conducted using an Applied Biosystems 3130 Genetic Analyzer (Fisher Scientific, Fair Lawn, New Jersey, USA) at the Genomics and Sequencing Laboratory, Auburn University, Alabama. Sequences were aligned using NCBI's nucleotide blast program.

Results

All adult males were unambiguously identified as *B. vaga* (couplet 16) using Roth's species key to the genus *Blattella* (Roth 1985). Diagnostic characteristics of adult males included similar size non-spine-like styles on the subgenital plate (Fig. 1A), only abdominal tergite 7 uniquely modified (Fig. 1B), and genital phallomere L2d present and flattened (Fig. 1C) (Roth 1985). Additional adult characteristics included wide longitudinal stripes on the pronotum and the area between the eyes darkly colored (Buxton & Freeman 1968). Nymphal characteristics included an overall yellowish coloration with longitudinal brown stripes. All morphological characters of the field-collected and reared specimens matched those of colony specimens.

Cuticular hydrocarbon analysis showed almost identical chromatograms for field-collected specimens and the laboratory colony of *B. vaga* (Fig. 2). The individual retention times of all 14 major cuticular hydrocarbon components were the same for the field-collected specimens and the laboratory colony, indicating a probable match.

Using GenBank data Sequence ID: AF228735.1 as a comparison, there was 98% similarity (Fig. 3) between the partial sequence of the mitochondrial cytochrome c oxidase subunit I-like protein gene from the field-collected cockroach and the published sequence for *B. vaga*, indicating a very high probability of a match.

Discussion

Based on morphological, molecular, and cuticular hydrocarbon analysis, we have identified the cockroaches collected from Mobile, Alabama, as the field cockroach, *Blattella vaga* Hebard. To the best of our knowledge, this is the first report of *B. vaga* collected from Alabama. As with most, if not all, reports of range extensions, it is possible




Fig. 1. Adult male *Blattella vaga* from field-collected colony. (A) Subgenital plate showing 2 non-spinelike styles; (B) abdominal tergite 7 with modification; (C) genital phallomere L2d.

those populations of *B. vaga* have existed in Alabama before 2016 and that populations may be distributed widely in southern Alabama.

There was a very high (98%) similarity in the partial gene sequence between the field-collected and published sequence for *B. vaga*. Even though the sequence similarity was not 100%, it is likely that differences in individual bases are not significant and result in coding for the same amino acids and ultimately proteins (Griffiths et al. 2008). There is no generally agreed upon percentage DNA sequence similarity for separation of insect species (Cognato 2006), although the small difference (2%) is certainly suggestive of a positive match.

Cuticular hydrocarbons are important for resistance to desiccation (Hadley 1981) and have been identified as pheromones (Blomquist 2010). Cuticular hydrocarbon analysis showed identical retention times for the 14 major components (Fig. 2), indicating the presence of the same hydrocarbons from field-collected and laboratory-reared *B. vaga*. Furthermore, our results are consistent with gas chromatographic profiles and mass spectra previously reported for *B. vaga* (Carlson & Brenner 1988). There were, however, quantitative differences between the 2 groups. Several studies have demonstrated quantitative differences in cuticular hydrocarbon profiles due to abiotic and biotic factors. Exposure to different temperatures (Gibbs et al. 1998) and consumption of different diets (Liang & Silverman 2000) can affect cuticular hydrocarbon profiles.

Several factors probably facilitated movement of *B. vaga* from Arizona and southern California to Alabama. Adults of both sexes readily fly during the photophase and are not repelled by light, behaviors




Fig. 2. Gas chromatography traces of cuticular hydrocarbons of (A) a male unknown field-collected cockroach, and (B) a male *Blattella vaga* from a laboratory colony. Numbers near peaks represent retention times.

Score		Expect	Identities	Gaps
809 bits(438)		0.0	460/470(98%)	5/470(1%)
Query 8	CCAGATATAGCAITTTCCCGAATAAATAATATAAGTTTCTCCGACTATTACCACCTTCA	67		
Sbjct 1	CCAGATATAGCAITTTCCCGAATAAATAATATAAGTTTCT--GACTATTACCACCTTCA	57		
Query 68	TTATCACITTTTATTGGCTAGTAGTCTGGTTGAAAGAGGTGCTGGAACAGGATGAACGTGA	127		
Sbjct 58	TTATCACITTTTATTGGCTAGTAGTCTGGTTGAAAGAGGTGCTGGAACAGGATGAACGTGA	117		
Query 128	TATCCACCTCTAGCAAGAGGTATTGCTCATGCTGGTGATCAGTTGATTAGCAATTTT	187		
Sbjct 118	TATCCACCTCTAGCAAGAGGTATTGCTCATGCTGGTGATCAGTTGATTAGCAATTTT	177		
Query 188	TCCTTACATTAGCAGGAGTTTCATCAATTTTAGTGCTGTTAAATTTTATTCAACAATT	247		
Sbjct 178	TCCTTACATTAGCAGGAGTTTCATCAATTTTAGTGCTGTTAAATTTTATTCAACAATT	237		
Query 248	ATTAATATAAAACCTATTAACATAAAACCTGAACGGATTCCCTTTTGTATGATCTGTA	307		
Sbjct 238	ATTAATATAAAACCTATTAACATAAAACCTGAACGGATTCCCTTTTGTATGATCTGTA	297		
Query 308	GGTATTACAGCTTTATTACTTTTATTATCATTACCTGATTAGCAGGGACGCTATTACAA	367		
Sbjct 298	GGTATTACAGCTTTATTACTTTTATTATCATTACCTGATTAGCAGGG--GCTATTACAA	355		
Query 368	TATTATTAACTGATCGTAATTTAAATACATCATCTTTGATCCTGCTGGTGGTGGTGATC	427		
Sbjct 356	TATTATTAACTGATCGTAATTTAAATACATCATCTTTGATCCTGCTGGTGGTGGTGATC	415		
Query 428	CAATTTTATATCAACATTTATTCTGATCTTTTGGTCATCCAGAGTTTAT	477		

Fig. 3. *Blattella vaga* cytochrome c oxidase subunit I-like protein gene, partial sequence; mitochondrial gene for mitochondrial product. Using GenBank data Sequence ID: AF228735.1, Length: 1,235, there was one 98% match confirming that the sequence for the wild-caught male cockroach matched the published sequence for *B. vaga*.

similar to *B. asahinai*, an invasive species that also has extended its range from 1 city in Florida to the entire state and northward into at least Alabama, Georgia, North Carolina, and South Carolina (Snoddy & Appel 2008). Rather than long directional flights, both *B. asahinai* and *B. vaga* tend to take short (0.5–5 m) flights and change direction from 1 flight to the next (Appel & Snoddy personal observations). We believe that these behaviors contribute to *B. vaga* flying onto vehicles and being transported to new locations. In addition, it is likely that *B. vaga*-infested potted plants, sod, waste, or other materials could be transported along interstate highways by commercial trucking.

Physiologically, *B. vaga* is relatively resistant to desiccation (Appel et al. 1983), which would allow it to survive exposure to hot and dry climates, as well as desiccation from exposure to moving air. Like *B. asahinai* (Snoddy 2007), this species avoids extreme conditions (hot and dry, and cold) by burrowing into a substrate.

In conclusion, we have used 3 independent methods (morphological, molecular, and chemical) that confirmed the identification of *B. vaga* from specimens collected in Mobile, Alabama. Because of its behavior, physiology, and distribution pattern along major interstate highways, it is likely this species will become established elsewhere in the southeastern US.

Acknowledgments

We thank Xing Ping Hu, Nannan Liu, and Charles H. Ray (Auburn University) for reviewing the manuscript and Marla J. Eva for technical support. This work was supported in part by the USDA National Institute of Food and Agriculture, AAES Hatch/Multistate Grants ALA-08-058 to AGA and ALA-08-044 to HYF.

References Cited

- Appel AG. 1995. *Blattella* and related species, pp. 1–19 In Rust ML, Owens JM, Reiersen DA [eds.], Understanding and Controlling the German Cockroach, Oxford University Press, New York, USA.
- Appel AG, Tucker JB. 1986. Occurrence of the German cockroach, *Blattella germanica* (Dictyoptera: Blattellidae), outdoors in Alabama and Texas. Florida Entomologist 69: 422–423.
- Appel AG, Reiersen DA, Rust MK. 1983. Comparative water relations and temperature sensitivity of cockroaches. Comparative Biochemistry and Physiology 74A: 357–361.
- Atkinson TH, Koehler PG, Patterson RS. 1991. Catalog and atlas of the cockroaches (Dictyoptera) of North America North of Mexico. Miscellaneous Publications, Entomological Society of America 78: 1–86.
- Austin JW, Glenn GJ, Szalanski AL, McKern JA, Gold RE. 2007. Confirmation of Asian cockroach *Blattella* [sic] *asahinai* (Blattodea: Blattellidae [sic]) introduction to Texas based on genetics, morphology, and behavior. Florida Entomologist 90: 574–576.
- Blomquist GJ. 2010. Biosynthesis of cuticular hydrocarbons, pp. 35–51 In Blomquist GJ, Bagnères AG [eds.], Insect Hydrocarbons. Cambridge University Press, Cambridge, United Kingdom.
- Buxton GM, Freeman TJ. 1968. Positive separation of *Blattella vaga* and *Blattella germanica* (Orthoptera: Blattellidae). Pan-Pacific Entomologist 44: 168–169.
- Carlson DA, Brenner RJ. 1988. Hydrocarbon-based discrimination of three North American *Blattella* cockroach species (Orthoptera: Blattellidae) using gas chromatography. Annals of the Entomological Society of America 81: 711–723.
- Cognato AI. 2006. Standard percent DNA sequence difference for insects does not predict species boundaries. Journal of Economic Entomology 99: 1037–1045.
- Dakin Jr ME, Hays KL. 1970. A synopsis of Orthoptera (*sensu lato*) of Alabama. Alabama Agricultural Experiment Station Bulletin 404: 1–118.
- Drees BM, Jackman JA. 1998. A Field Guide to Common Texas Insects. Gulf Publishing Company, Houston, Texas, USA.
- Ebeling W. 1978. Urban Entomology. Division of Agricultural Sciences, University of California, Berkeley, California, USA.
- Ebeling W, Wagner RE, Reiersen DA. 1966. Influence of repellency on the efficacy of blatticides: I. Learned modification of behavior of the German cockroach. Journal of Economic Entomology 59: 1374–1388.
- Gibbs AG, Louie AK, Ayala JA. 1998. Effects of temperature on cuticular lipids and water balance in a desert *Drosophila*: Is thermal acclimation beneficial? Journal of Experimental Biology 201: 71–80.
- GBIF Secretariat. 2017. *Blattella vaga* Hebard, 1935. In GBIF Backbone Taxonomy. <https://doi.org/10.5072/hufs9m> (last accessed via GBIF.org on 13 Oct 2017)
- Griffiths AJF, Wessler SR, Lewontin RC, Carroll SB. 2008. Introduction to genetic analysis, 9th ed. W. H. Freeman and Company, New York, USA.
- Hadley NF. 1981. Cuticular lipids of terrestrial plants and arthropods: a comparison of their structure, composition, and waterproofing function. Biological Reviews 56: 23–47.
- Hebard M. 1935. Studies in the Orthoptera of Arizona, Part 1. New genera, species and geographical races. Transactions of the American Entomological Society 59: 111–153.
- Helper JR. 1987. How to Know the Grasshoppers, Crickets, Cockroaches and their Allies. Dover Publications, Mineola, New York, USA.
- Hogue CL. 1993. Insects of the Los Angeles Basin, 2nd ed. Natural History Museum of Los Angeles County, California, USA.
- Liang D, Silverman J. 2000. You are what you eat: diet modifies cuticular hydrocarbons and nestmate recognition in the Argentine ant, *Linepithema humile*. Naturwissenschaften 87: 412–416.
- Pachamuthu P, Kamble ST, Clar, TL, Foster JE. 2000. Differentiation of three phenotypically similar *Blattella* spp.: analysis with polymerase chain reaction-restricted fragment length polymorphism of mitochondrial DNA. Annals of the Entomological Society of America 93: 1138–1146.
- Pratt HD. 1988. Annotated checklist of the cockroaches (Dictyoptera) of North America. Annals of the Entomological Society of America 81: 882–885.
- Roth LM. 1985. A taxonomic revision of the genus *Blattella* Caudell (Dictyoptera, Blattaria: Blattellidae). Entomologica Scandinavica, Supplement 22: 1–221.
- Snoddy ET. 2007. Distribution and population dynamics of the Asian cockroach (*Blattella asahinai* Mizukubo) in southern Alabama and Georgia. MS thesis, Auburn University, Auburn, Alabama, USA.
- Snoddy ET, Appel AG. 2008. Distribution of *Blattella asahinai* (Dictyoptera: Blattellidae) in southern Alabama and Georgia. Annals of the Entomological Society of America 101: 397–401.