

First Record of *Liriomyza huidobrensis* (Diptera: Agromyzidae) Disseminating *Alternaria solani* (Pleosporaceae) in Potato Crops in Brazil

Authors: Soares, Walyson Silva, Plata-Rueda, Rosa Angelica, Fernandes, Maria Elisa de Sena, Fernandes, Flávio Lemes, Alves, Flávia Maria, et al.

Source: Florida Entomologist, 102(1) : 234-235

Published By: Florida Entomological Society

URL: <https://doi.org/10.1653/024.102.0139>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

First record of *Liriomyza huidobrensis* (Diptera: Agromyzidae) disseminating *Alternaria solani* (Pleosporaceae) in potato crops in Brazil

Walyson Silva Soares¹, Rosa Angelica Plata-Rueda²,
Maria Elisa de Sena Fernandes², Flávio Lemes Fernandes^{2,*},
Flávia Maria Alves³, and Ítalo Willian da Silva²

Liriomyza leafminers can adversely affect agronomic crops by transmitting a number of disease pathogens, including fungi, bacteria, and viruses. Often this results from the physical damage to leaves following insertion of an ovipositor by females during the act of egg laying. For example, Durairaj et al. (2010) reported that *L. trifolii* (Burgess) (Diptera: Agromyzidae) females perforate leaves of *Solanum lycopersicum* L. (Solanaceae), predisposing the plant to infection by *Alternaria* (Pleosporales: Pleosporaceae) fungi. Similarly, Deadman et al. (2002) reported that leaf perforation during *L. trifolii* oviposition favored leaf necrosis caused by *Alternaria alternata* (Fr.) Keissl in *S. tuberosum* L. (Solanaceae). *Alternaria* spp. can reduce photosynthetic leaf area and subsequent tuber production in infected potato plants (Simmons 2000).

The economic impacts of *Liriomyza* are high. For example, *Liriomyza* spp. in Kenya have caused losses of US\$54 and US\$64.5 million in *Pisum sativum* L. and *Phaseolus vulgaris* L. (Fabaceae), respectively (Pratt et al. 2017). These species also are agronomic pests of other vegetable crops such as *Beta vulgaris* L. and *Spinacia oleracea* L. (Amaranthaceae) (Minkenberg 1988). Mujica and Kroschel (2013) reported that the leafminer *L. huidobrensis* (Blanchard) (Diptera: Agromyzidae) was responsible for leaf damage that subsequently reduced yields of *S. tuberosum* by up to 51% in the Cañete Valley (central coast of Peru). Here, we provide the first report on the association of *L. huidobrensis* as a facilitating agent of *A. solani* (Cooke) Wint infection to *S. tuberosum* in a commercial potato crop in Brazil.

Leafminer sampling was conducted twice between Sep 2013 to Apr 2014 in the municipality of Rio Paranaíba (Minas Gerais State, Brazil) on *S. tuberosum* leaves (cv. Ágata) from a field containing larval *L. huidobrensis*. The field (19.208744°S, 46.164902°W) consisted of 35 ha, with potato plants at the tuber formation stage (50 d after planting) (Meier 2001). The field was divided into 5 areas of 5 ha each. Each area was further divided into 10 parcels of 50 × 50 m (2,500 m²). We evaluated 50 plants per parcel.

Plants were spaced 80 to 87 cm apart with 12 to 15 stems per meter. Mines of *L. huidobrensis* larvae with *A. solani* lesions were visually counted following the method of Alves et al. (2014) (Fig. 1a). Spatial distribution of mines caused by *L. huidobrensis* and *A. solani* lesions was evaluated using the Moran Index (I) ($P \leq 0.05$)

with the Z test (Seffrin et al. 2018). This index measures whether a distribution is irregular, even, or clustered. Indices closest to 1 indicate high spatial dependence (clustered values); those closest to zero are characterized as spatial independence, while indices closest to -1 indicate high level of dispersion.

We found a positive and significant correlation ($r = 0.84$; $n = 500$; $P < 0.001$) between the number of *L. huidobrensis* mines and *A. solani* lesions, with the dissemination of this pathogen in potato plants being influenced by the degree of spatial distribution of *L. huidobrensis* ($I = +0.225$; $P = 0.001$) and *A. solani* ($I = +0.428$; $P = 0.001$). Moreover, the presence of *A. solani* concentric halos in potato leaves associated with *L. huidobrensis* oviposition sites, coincided with growth of this fungus. Our results were similar to those of Durairaj et al. (2010) who reported that *L. trifolii* oviposition sites favored *A. alternata* infection in tomatoes. Indeed, Deadman et al. (2002) observed that as *L. trifolii* oviposition perforations in *S. tuberosum* leaf tissue increased, *A. alternata* lesions increased (Deadman et al. 2002).

Summary

A significant association between *Liriomyza huidobrensis* leafminers and *Alternaria solani* fungus is reported here for the first time in a commercial potato (*Solanum tuberosum*) field in Brazil. We found that there was a high correlation (0.84) between the number of *L. huidobrensis* mines and *A. solani* lesions on infected plants.

Key Words: cluster; miner; oviposition; Solanaceae; spatial distribution; symptoms

Sumário

Uma associação significativa entre a mosca minadora *Liriomyza huidobrensis* e o fungo *Alternaria solani* é reportado aqui pela primeira vez em campo de batata comercial (*Solanum tuberosum*) no Brasil. Nós verificamos alta correlação especial (0,84) entre o número de minas de *L. huidobrensis* e as lesões de *A. solani* em plantas infectadas.

Palavras Chaves: agregação; minador; oviposição; Solanaceae; distribuição espacial; sintomas

¹Universidade Federal de Uberlândia, Instituto de Ciências Agrárias, Uberlândia, Minas Gerais 38400-902, Brazil; E-mail: walysonagronomia@gmail.com (W. S. S.)

²Universidade Federal de Viçosa, Instituto de Ciências Agrárias, Rio Paranaíba, Minas Gerais 38810-000, Brazil; E-mail: angelicaplata@yahoo.com.mx (R. A. P. R.), maria.sena@ufv.br (M. E. S. F.), flaviofernandes@ufv.br (F. L. F.), italo.silva@ufv.br (I. W. S.)

³Universidade Federal de Viçosa, Departamento de Fitotecnia, Viçosa, Minas Gerais 36570-900, Brazil; E-mail: flaviamarialves@gmail.com (F. M. A.)

*Corresponding author; E-mail: flaviofernandes@ufv.br




Fig. 1. (a) Presence of *Liriomyza huidobrensis* mines on a *Solanum tuberosum* leaf infected with *Alternaria solani*; (b) diagram showing representative distribution of mines and fungal lesions.

References Cited

- Alves FM, Diniz JFS, Silva IW, Fernandes FL, Silva PR, Gorri JER. 2014. A sampling plan for *Liriomyza huidobrensis* (Diptera: Agromyzidae) on a potato (*Solanum tuberosum*) plantation. American Journal of Potato Research 91: 663–672.
- Deadman ML, Khan IA, Thacker JRM, Al-Habsi K. 2002. Interactions between leafminer damage and leaf necrosis caused by *Alternaria alternata* on potato in the sultanate of Oman. Plant Pathology Journal 18: 210–215.
- Durairaj C, Karthikeyan G, Ganapathy N, Karuppuchamy P. 2010. Predisposition effect of *Liriomyza trifolii* damage to *Alternaria* leaf spot disease in tomato. Karnataka Journal of Agricultural Science 23: 161–162.
- Meier U. 2001. Growth stages of mono and dicotyledonous plants. BBCH (Biologische Bundesanstalt, Bundessortenamt and Chemical industry) Monograph, Federal Biological Research Centre for Agriculture and Forestry. Berlin, Germany.
- Minkenberg OPJM. 1988. Dispersal of *Liriomyza trifolii*. EPPO Bulletin 18: 173–182.
- Mujica N, Kroschel J. 2013. Pest intensity-crop loss relationships for the leafminer fly *Liriomyza huidobrensis* (Blanchard) in different potato (*Solanum tuberosum* L.) varieties. Crop Protection 47: 6–16.
- Pratt CF, Constantine KL, Murphy ST. 2017. Economic impacts of invasive alien species on African smallholder livelihoods. Global Food Security 14: 31–37.
- Seffrin R, Araújo EC, Bazzi CL. 2018. Regression models for prediction of corn yield in the state of Paraná (Brazil) from 2012 to 2014. Acta Scientiarum Agronomy 40: 1–11.
- Simmons EG. 2000. *Alternaria* themes and variations (244–286): species on Solanaceae. Mycotaxon 75: 1–115.