



Alternative Survey Methods for the Emerald Ash Borer

Authors: Francese, Joseph A., Booth, Everett G., Lopez, Vanessa M., and Sorensen, Benjamin

Source: Florida Entomologist, 102(1) : 243-245

Published By: Florida Entomological Society

URL: <https://doi.org/10.1653/024.102.0142>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Alternative survey methods for the emerald ash borer

Joseph A. Francese^{1,*}, Everett G. Booth¹, Vanessa M. Lopez^{2,3}, and Benjamin Sorensen⁴

The emerald ash borer, *Agrilus planipennis* Fairmaire (Coleoptera: Buprestidae), a highly destructive pest of ash (*Fraxinus* L.) (Oleaceae), was discovered initially in North America in 2002 near Detroit, Michigan, USA. It has subsequently been detected in 31 additional US states and in 2 Canadian provinces (Haack et al. 2002; Emerald Ash Borer Info 2017). Since its arrival in the mid-1990s (Siegert et al. 2014), it is estimated that *A. planipennis* has killed tens of millions of trees throughout this range (Emerald Ash Borer Info 2017).

Since 2008, the US Department of Agriculture, Animal and Plant Health Inspection Service, Plant Protection and Quarantine (USDA-APHIS-PPQ) Emerald Ash Borer Cooperative Program has conducted a multi-state emerald ash borer survey (USDA APHIS PPQ 2017). The traps currently deployed are glue-coated purple prism traps (Francese et al. 2008, 2013a) baited with [3Z]-hexenol, a green leaf volatile found to increase *A. planipennis* trap catch (Grant et al. 2010, 2011; Poland et al. 2011; Crook et al. 2012).

Multi-funnel traps (Lindgren 1983) have been shown to be a promising tool for catching emerald ash borer (Francese et al. 2011). Because these traps lack the messy adhesive-coating found on the prism traps, multi-funnel traps are a more user-friendly option that do not need to be discarded after each use. Green multi-funnel traps, based on a color attractive to emerald ash borer (Crook et al. 2009; Francese et al. 2010) caught more *A. planipennis* than the standard black multi-funnel traps, purple multi-funnel traps, and prism traps in trapping assays conducted in heavily infested areas in southeastern and south-central Michigan. Fluon®-coated intercept panel traps have been shown to increase cerambycid trap catch, and remain effective for more than 1 field season (Graham et al. 2010; Graham & Poland 2012; Allison & Redak 2017). Fluon®-coated green multi-funnel traps also caught 40× more emerald ash borer adults than traps not treated with any coating (Francese et al. 2013b). In low emerald ash borer density areas, green multi-funnel traps have been shown to be as effective in detecting populations as purple prisms (Crook et al. 2014).

Green multi-funnel traps have been available for use as a survey and detection tool for emerald ash borer since 2015, but due to their initial purchase and associated survey costs, they are not in wide programmatic use. The current survey guidelines recommend that during the course of the field season, multi-funnel traps be checked every 2 to 3 wk and prism traps only every 6 wk (USDA APHIS PPQ 2017). This additional maintenance leads to extra travel and personnel hours spent surveying, which increases program costs. Trap upkeep costs would include the additional killing agent that gets added at each trap check. Currently, the recommended killing agent in the US for the green multi-

funnel traps is a propylene glycol solution (about 20–30%) in water, in the form of “recreational vehicle” antifreeze (USDA APHIS PPQ 2017). Several studies found that traps equipped with a wet cup captured more cerambycid and buprestid species than traps equipped with a dry cup; however, because there were no *Agrilus* species captured in these studies, additional investigations are needed to evaluate this trapping method for *A. planipennis* (Allison & Redak 2017).

The overall goal of the research presented here was to assess various trap modifications and survey methods that could help to both reduce the aforementioned costs associated with multi-funnel traps, and thus provide alternatives to sticky traps for survey of emerald ash borer. In particular, the main objectives were to investigate the effect of the trap checking frequency and the effect of wet vs. dry killing agents on emerald ash borer trap catch and detection. To test these objectives, 2 separate trapping studies, the trap check interval study (Study 1) and the trap cup collection method study (Study 2), were conducted using green plastic multi-funnel traps (Chemtica USA, Durant, Oklahoma, USA) previously described by Francese et al. (2011). Traps were unbaited and coated with a 50% dilution of Fluon® that previously had been shown to be as effective at catching *A. planipennis* (Francese et al. 2013b) and other woodborers (Allison et al. 2016) as a 100% dilution. For both studies, traps were hung from ropes in the lower canopy (5–8 m) of host along the edges of white (*Fraxinus americana* L.) and green (*Fraxinus pennsylvanica* Marshall) ash-dominated woodlots in Edmore, Michigan; North Andover, Massachusetts; and Bethel, Ohio, USA. No lures were used on the traps. We replicated treatments in a randomized complete block design with trap lines as blocks. Within each trap line, we put traps in adjacent trees with an average of 5 m between traps.

The trap check frequency study was conducted in both North Andover, Massachusetts (n = 7) and Bethel, Ohio (n = 7) to determine how often multi-funnel traps needed to be checked. Four trap check intervals were compared: 1-wk, 3-wk, 6-wk, and 12-wk. Trap timing intervals were chosen based on the recommended survey guidelines (3 wk), recommended lure change timing (6 wk), and the full location specific predicted *A. planipennis* flight period duration (12 wk). Because 1 of the 12 wk traps placed in Massachusetts fell during the course of the season, the trap line that was a part of that replicate was removed from the study.

A trap cup collection method study was conducted on private land in Edmore, Michigan (n = 5), North Andover, Massachusetts (n = 5), and at East Fork State Park in Bethel, Ohio (n = 8) to compare alternate methods for killing and collecting captured *A. planipennis* adults. Four treatments were compared: (1) standard wet collection cup filled

¹USDA APHIS PPQ CPHST Otis Laboratory, 1398 West Truck Road, Buzzards Bay, Massachusetts 02542, USA; Email: joe.francese@aphis.usda.gov (J. A. F.), everett.g.booth@aphis.usda.gov (E. G. B.)

²USDA Forest Service, State and Private Forestry, Forest Health Protection, 201 14th Street, SW, FHP 3CE, Washington, DC 20250, USA; Email: vanessalopez@fs.fed.us (V. M. L.)

³Department of Biology, Xavier University, 3800 Victory Parkway, Cincinnati, Ohio 45207, USA; Email: vanessalopez@fs.fed.us (V. M. L.)

⁴USDA APHIS PPQ CPHST Brighton Field Station, 5936 Ford Court, Suite 200, Brighton, Michigan 48116, USA; Email: bensoren87@gmail.com (B. S.)

*Corresponding author; E-mail: joe.francese@aphis.usda.gov

with 150 to 200 mL of propylene glycol (Camco Easygoing -50, Camco, Greensboro, North Carolina, USA); (2) dry collection cup with an internal funnel (cone 7.0 cm long, 13.7 cm diam.; stem 6.7 cm long, 3.9 cm diam.) placed in the bottom trap funnel to reduce the chances of escape; (3) dry collection cup containing an insecticidal strip (Vaportape II; 2,2-dichlorovinyl dimethyl phosphate (10%) Hercon Environmental, Emingsville, Pennsylvania, USA); and (4) dry collection cup containing an internal funnel and an insecticidal strip. Internal funnels were coated with Fluon® (50% solution) to prevent beetles from climbing out of the collection cup.

During trap checks, the contents of each trap cup were strained using a paper paint filter. Paint filters were then placed in individual, labeled Whirl-Pak sampling bags (Nasco, Fort Atkinson, Wisconsin, USA), ethanol was added for preservation, and samples were stored in a freezer until sorting and identification could be conducted. In both assays, collected beetles were summed for each trap over the entire field season. Summed catch was log-transformed ($y + 0.5$) prior to statistical analysis to normalize the data, which was confirmed by testing residuals after ANOVA. Separate analyses of variance (ANOVA) were performed on the transformed total number of *A. planipennis* adults captured per trap for each study, to compare treatment effects (cup collection method or trap check interval depending on the study) (JMP 10) (SAS Institute 2012). Confidence intervals (95%) were calculated from the standard error of the transformed trap catch. Means and confidence intervals were then back-transformed for presentation in the text and tables that follow.

In the trap check frequency study, the interval between checks ($F = 0.66$; $df = 3, 52$; $P = 0.58$) did not significantly affect trap catch (Table 1). Because there was greater decomposition and the 6- and 12-wk interval, samples had to be sorted and identified in a fume hood to reduce the smell; however, identification characteristics were not affected. These results suggest that from the program perspective, multi-funnel traps could be checked less frequently than currently recommended, which would greatly reduce costs. However, it also should be noted that leaving the traps in the field for 12 wk without checking them could lead to loss of data as was the case with the 12-wk trap that fell during the course of the study.

In the trap collection method study, collection method did not ($F = 0.03$; $df = 3, 68$; $P = 0.99$) play a significant role in trap catch (Table 2). Based on these results, dry cup methods could be used as an alternative to the propylene glycol wet cups. These results are encouraging, especially for emerald ash borer trapping in more remote areas where transporting or disposing of used propylene glycol is not practical. As expected, the dry cup without a killing agent was still effective, but would not be recommended in most situations because live beetles were found in the traps during the check period. However, this method does show promise for live-trapping other woodborer species attracted to these traps.

The staff members of the USDA APHIS PPQ Otis Laboratory, and Bethel and Brighton Field Stations provided field work assistance: Scott Gula, Mandy Furtado, Erin Schott, Elizabeth Reardon, MacKenzie O’Kane, Alexander Reitz, Brenna Walters, Sam Engle, Alyssa Perry, and

Table 1. Back-transformed mean (\pm 95% confidence intervals) *Agrilus planipennis* trap catch in green multi-funnel traps checked at 1 of 4 intervals ($n = 14$). Trap catch had been summed for each trap for the entire season.

Collection cup method	Trap catch
Weekly	8.03 (4.55 – 14.15)
Every 3 wk	22.31 (13.45 – 37.00)
Twice during the field season (at 6 and 12 wk)	22.72 (12.23 – 42.22)
Once at the end of the field season (at 12 wk)	18.52 (9.14 – 37.54)

Table 2. Back-transformed mean (\pm 95% confidence intervals) *Agrilus planipennis* trap catch in green multi-funnel traps using 1 of 4 different collection methods ($n = 18$). Trap catch had been summed for each trap for the entire season.

Collection cup method	Trap catch
Standard wet cup (propylene glycol)	8.95 (4.89 – 16.37)
Dry cup with internal funnel, no pesticide strip	9.31 (5.31 – 16.31)
Dry cup without funnel, with pesticide strip	9.31 (6.18 – 18.37)
Dry cup with funnel, with pesticide strip	8.29 (4.65 – 14.78)

Patrick Gemperline. Ann Ray of Xavier University provided planning and technical assistance. This work was funded by the USDA APHIS PPQ Emerald Ash Borer Program.

Summary

As part of an ongoing project to improve survey and detection for the emerald ash borer, *Agrilus planipennis*, several field assays were conducted to (1) determine how often traps need to be checked during a given field season, and (2) compare the effectiveness of traps with “dry” (with insecticidal strips or internal funnel) vs. “wet” (with propylene glycol surfactant) collection cups. There were no significant differences among any of the trap check intervals or the trap methods tested. This will provide new tools to surveyors, and allow them more flexibility as they survey for this invasive pest.

Key Words: *Agrilus planipennis*; green multi-funnel traps; wet cup; dry cup; trap check interval

Sumario

Se realizaron varios ensayos de campo como parte de un proyecto en marcha para mejorar el sondeo y la detección del barrenador esmeralda del fresno, *Agrilus planipennis* para (1) determinar con qué frecuencia se deben revisar las trampas durante una temporada de campo determinada, y (2) comparar la efectividad de trampas “secas” (con tiras de insecticida o embudo interno) vs. “húmedas” (con surfactante de propilenglicol). No hubo diferencias significativas entre ninguno de los intervalos de control de trampa o los métodos de trampa probados. Estos resultados proporcionan nuevas herramientas a los topógrafos y les permitirá una mayor flexibilidad a medida que realizan el sondeo para la detección de esta plaga invasiva.

Palabras Clave: *Agrilus planipennis*; trampas verdes multi-embudo; taza mojada; taza seca intervalo de revisar trampas

References Cited

- Allison JD, Redak RA. 2017. The impact of trap type and design features on the survey and detection of bark and wood boring beetles and their associates: a review and meta-analysis. *Annual Review of Entomology* 62: 127–146.
- Allison JD, Graham EE, Poland TM, Strom BL. 2016. Dilution of fluon before trap surface treatment has no effect on longhorned beetle (Coleoptera: Cerambycidae) captures. *Journal of Economic Entomology* 109: 1215–1219.
- Crook DJ, Francese JA, Rietz ML, Lance DR, Hull-Sanders HM, Mastro VC, Silk PJ, Ryall KL. 2014. Improving detection tools for emerald ash borer (Coleoptera: Buprestidae): comparison of multifunnel traps, prism traps and lure types at varying population densities. *Journal of Economic Entomology* 107: 1496–1501.
- Crook DJ, Francese JA, Zylstra KE, Fraser I, Sawyer AJ, Bartels DW, Lance DR, Mastro VC. 2009. Laboratory and field response of the emerald ash borer

- (Coleoptera: Buprestidae) to selected regions of the electromagnetic spectrum. *Journal of Economic Entomology* 102: 2160–2169.
- Crook DJ, Khrimian A, Cosse A, Fraser I, Mastro VC. 2012. Influence of trap color and host volatiles on capture of the emerald ash borer (Coleoptera: Buprestidae). *Journal of Economic Entomology* 105: 429–437.
- Emerald Ash Borer Info. 2017. Emerald ash borer. (online) <http://www.emeraldashborer.info> (last accessed 24 Nov 2018).
- Francese JA, Crook DJ, Fraser I, Lance DR, Sawyer AJ, Mastro VC. 2010. Optimization of trap color for emerald ash borer (Coleoptera: Buprestidae). *Journal of Economic Entomology* 103: 1235–1241.
- Francese JA, Fraser I, Lance DR, Mastro VC. 2011. Efficacy of multi-funnel traps for capturing emerald ash borer (Coleoptera: Buprestidae): effect of color, glue, and other trap coatings. *Journal of Economic Entomology* 104: 901–908.
- Francese JA, Oliver JB, Fraser I, Lance DR, Youssef N, Sawyer AJ, Mastro VC. 2008. Influence of trap placement and design on capture of the emerald ash borer (Coleoptera: Buprestidae). *Journal of Economic Entomology* 101: 1831–1837.
- Francese JA, Rietz ML, Crook DJ, Fraser I, Lance DR, Mastro VC. 2013a. Improving detection tools for the emerald ash borer (Coleoptera: Buprestidae): comparison of prism and multifunnel traps at varying population densities. *Journal of Economic Entomology* 106: 2407–2414.
- Francese JA, Rietz ML, Mastro VC. 2013b. Optimization of multifunnel traps for emerald ash borer (Coleoptera: Buprestidae): influence of size, trap coating and color. *Journal of Economic Entomology* 106: 2415–2423.
- Graham EE, Mitchell RF, Reigel PF, Barbour JD, Millar JG, Hanks LM. 2010. Treating panel traps with a fluoropolymer enhances their efficiency in capturing cerambycid beetles. *Journal of Economic Entomology* 103: 641–647.
- Graham EE, Poland TM. 2012. Efficacy of fluron conditioning for capturing cerambycid beetles in different trap designs and persistence on panel traps over time. *Journal of Economic Entomology* 105: 395–401.
- Grant GG, Poland TM, Ciaramitaro T, Lyons DB, Jones GC. 2011. Comparison of male and female emerald ash borer (Coleoptera: Buprestidae) responses to phoebe oil and (Z)-3-hexenol lures in light green prism traps. *Journal of Economic Entomology* 104: 173–179.
- Grant GG, Ryall KL, Lyons DB, Abou-Zaid MM. 2010. Differential response of male and female emerald ash borers (Col., Buprestidae) to (Z)-3-hexenol and manuka oil. *Journal of Applied Entomology* 134: 26–33.
- Haack RA, Jendek E, Liu H, Marchant KR, Petrice TR, Poland TM, Ye H. 2002. The emerald ash borer: a new exotic pest in North America. *Newsletter of the Michigan Entomological Society* 47: 1–5.
- Lindgren BS. 1983. A multiple funnel trap for scolytid beetles (Coleoptera). *The Canadian Entomologist* 115: 299–302.
- Poland TM, McCullough DG, Anulewicz AC. 2011. Evaluation of double-decker traps for emerald ash borer (Coleoptera: Buprestidae). *Journal of Economic Entomology* 104: 517–531.
- USDA-APHIS-PPQ (US Department of Agriculture, Animal and Plant Health Inspection Service, Plant Protection and Quarantine). 2017. 2017 emerald ash borer survey guidelines. (online) http://www.aphis.usda.gov/plant_health/plant_pest_info/emerald_ash_b/downloads/survey_guidelines.pdf (last accessed 24 Nov 2018).
- SAS Institute. 2012. JMP version 10. SAS Institute, Cary, North Carolina, USA.
- Siegert NW, McCullough DG, Liebhold AM, Telewski FW. 2014. Dendrochronological reconstruction of the epicenter and early spread of emerald ash borer in North America. *Diversity and Distributions* 20: 847–858.