



Lance Flies (Diptera: Lonchaeidae) Associated with *Acca Sellowiana* (Myrtaceae) in Lages, Santa Catarina, Brazil

Authors: Souza, Gabriely Koerich, Pikart, Tiago Georg, Strikis, Pedro Carlos, Oliveira, Viviane Lunelli de, Bender, Mariana, et al.

Source: Florida Entomologist, 102(1) : 249-250

Published By: Florida Entomological Society

URL: <https://doi.org/10.1653/024.102.0144>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Lance flies (Diptera: Lonchaeidae) associated with *Acca sellowiana* (Myrtaceae) in Lages, Santa Catarina, Brazil

Gabriely Koerich Souza^{1,*}, Tiago Georg Pikart², Pedro Carlos Strikis³, Viviane Lunelli de Oliveira¹, Mariana Bender¹, Pedro Boff⁴, and Mari Inês Carissimi Boff¹

Feijoa (*Acca sellowiana* [Berg] Burret) (Myrtaceae) is a small fruit tree native to the highlands of southern Brazil and northeast Uruguay (Barni et al. 2004). In its natural habitat in Brazil, this species occurs more often in woodland environs (capons) and Araucaria forests at altitudes between 900 and 1,100 masl (Lorenzini et al. 2007). The flesh of feijoa fruit is white, sweet, and aromatic with a long-lasting pineapple-like scent, and a taste reminiscent of barren strawberries (*Potentilla*) (Rosaceae) and pineapples (*Ananas comosus* [L.] Merr.) (Bromeliaceae) (Pasquariello et al. 2015). Feijoa fruit is also a source of vitamins, minerals, and secondary metabolites, with antibacterial, antioxidant, antiallergenic, and immunological properties (Weston 2010).

Because local feijoa production has been limited to regions where it naturally occurs in Brazil, fruit pests such as the South American fruit fly, *Anastrepha fraterculus* (Wiedemann) (Diptera: Tephritidae), have been reported to infest *A. sellowiana* up to 100% during the fruit ripening period (Ducroquet et al. 2000). Recently, the spotted wing drosophila, *Drosophila suzukii* (Matsumura) (Diptera: Drosophilidae), also was reported attacking feijoa fruit in Brazil (Souza et al. 2017), reinforcing the importance of frugivorous flies to the cultivation of feijoa.

Another group of frugivorous flies, the lance flies (Diptera: Lonchaeidae) are very small, glossy dipterans that have been reported to be primary invaders of commercial plant species, with some fly species considered major agricultural pests of South American fruits and vegetables (Strikis & Prado 2005; Souza-Filho et al. 2009). Although a few species of Lonchaeidae already have been reported in feijoa (Gattelli et al. 2008; Nunes et al. 2012; Gislotti et al. 2017), little information is available about this association in the state of Santa Catarina. We report here on a study to document the species of Lonchaeidae associated with fruits of *A. sellowiana* in an orchard located in Lages, Santa Catarina, Brazil.

In Feb 2018, fruits (11.6 kg, 671 fruit) were harvested from 13 feijoa trees in an experimental orchard in the Estação Experimental de Lages of the Empresa de Pesquisa Agropecuária e Extensão Rural de Santa Catarina (EPAGRI) in Lages, Santa Catarina, Brazil (27.8086°S, 50.3306°W). Harvested fruits were transported to the Laboratório de Pesquisa em Entomologia of the Centro de Ciências Agroveterinárias da Universidade do Estado de Santa Catarina. Infested fruits were individually placed into plastic pots (750 mL) containing sterile moist vermiculite, and incubated at 25 ± 2 °C, 70 ± 10% RH, and a 12:12 h (L:D) photoperiod to allow fly larvae to pupate. After 30 d, Lonchaei-

dae pupae were removed from the vermiculite and placed in a 750 mL plastic pot cage for adult emergence. At emergence, adults were collected and killed in a freezer, then stored in 70% alcohol for subsequent identification.

A total of 224 pupae was obtained, resulting in an infestation level of 0.3 puparia per fruit and 19.3 puparia per kg of fruit. From the pupae, 71 adults of Lonchaeidae emerged. Thirty-six female specimens were not identified beyond the family level, because Lonchaeidae identification is based primarily on the morphology of male terminalia. The 35 male specimens recovered from the same fruits were identified as: *Neosilba pradoi* Strikis & Lerena (N = 27), *Neosilba delvechioi* Strikis 2011 (N = 4), *Neosilba zadolicha* McAlpine & Steyskal (N = 2), *Neosilba bella* Strikis & Prado (N = 1), and *Neosilba bifida* Strikis & Prado (N = 1). *Neosilba bella* and *N. zadolicha* had been recorded already from *A. sellowiana* (Gattelli et al. 2008; Nunes et al. 2012), but this is the first report of *N. bifida*, *N. delvechioi*, and *N. pradoi* infesting fruits of feijoa. Moreover, this is the first report that *N. bella*, *N. bifida*, and *N. delvechioi* have been reported from the state of Santa Catarina.

In our study, *N. pradoi* was the predominant species obtained from fruits of *A. sellowiana*, representing 77.1% of the male adults recovered. Garcia & Norrbom (2011) reported similar results, where they sampled the fruits of 46 plant species belonging to 25 families from 6 municipalities of Santa Catarina; the next most abundant species they collected was *N. zadolicha*. In a recent survey conducted by Calvo et al. (2017), *N. pradoi* was collected from 18 cultivated and wild plant species in Uruguay.

Neosilba bella, *N. bifida*, *N. delvechioi*, and *N. zadolicha* were present in very low numbers in our study, which could suggest their occurrence as being accidental infestations of *A. sellowiana* fruit. According to Gislotti et al. (2017), this situation may suggest that immigration from other nearby host species and surrounding forest areas was occurring. The capability of fruit flies to move from adjacent native vegetation, particularly forest fragments, into orchards was previously demonstrated by Kovalski et al. (1999) and Vargas et al. (2001). Indeed, the feijoa orchard sampled in our study is surrounded by some woodland areas where other Myrtaceae species are present that may serve as reservoirs for these lonchaeid species.

It is important to state that *N. delvechioi* was described by Strikis (2011) from drawings made by Maria Cecilia Delvechio, a researcher that started the studies of *Neosilba* species in the state of São Paulo.

¹Centro de Ciências Agroveterinárias, Universidade do Estado de Santa Catarina, Lages, Santa Catarina, 88520-000, Brazil; E-mail: gaby.florestal@gmail.com (G. K. S.), vivilunelli.oliveira@gmail.com (V. L. O.), bendermari97@hotmail.com (M.B.), mari.boff@udesc.br (M. I. C. B.)

²Centro de Ciências Biológicas e da Natureza, Universidade Federal do Acre, Rio Branco, Acre, 69920-900, Brazil; E-mail: tiago.florestal@gmail.com (T. G. P.)

³Laboratório de Paleontologia, Instituto de Geociências, Universidade de São Paulo, São Paulo, São Paulo, 05508-080, Brazil; E-mail: pcstrikis@gmail.com (P. C. S.)

⁴Estação Experimental de Lages, Empresa de Pesquisa Agropecuária e Extensão Rural de Santa Catarina, Lages, Santa Catarina, 88502-970, Brazil; E-mail: boff.pedro@yahoo.com.br (P. B.)

*Corresponding author; E-mail: gaby.florestal@gmail.com (G. K. S.)

She found only 1 specimen in 1978 in a peach orchard in the municipality of Monte Alegre do Sul, which is situated on the border of the state near Minas Gerais. In that region, the climate is characterized by an oceanic climate with warm summers (designated as “Cfb” according to the Köppen & Geiger classification), with a mean temperature of 18.9 °C. For more than 30 yr, surveys were carried out to try to find more males of *N. delvechioi* in several regions of the state of São Paulo, where this species was thought to be extinct. After finding it in Santa Catarina, we believe that it is a species adapted to colder climates where it may be important as a pest. In the state of Paraná, this species was once found in considerably larger numbers attacking apples (varieties ‘Fuyu’ and ‘Eva’) along with *N. zadolicha*, *N. certa*, and *N. pradoi*, the 3 last species in minor numbers.

Although studies recently have increased our knowledge on Lonchaeidae with new distribution and host records for Santa Catarina State (Garcia & Norrbom 2011; Strikis 2011), this is the first time that *Neosilba* species are reported damaging fruits of feijoa in Santa Catarina State. Considering that species of *Neosilba* can be primary invaders of commercial plant species, and that *N. pradoi* has been recorded in the region of natural occurrence of feijoa (Gattelli et al. 2008; Nunes et al. 2012; Calvo et al. 2017), our results suggest that this species has the potential to become an economically important pest in feijoa orchards.

The authors thank the Conselho Nacional de Desenvolvimento Científico e Tecnológico (CNPq), Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES), Rede Guarani/Serra Geral - FAPESC TO 2015TR10672, and the Núcleo Agroecologia e Saúde Ambiental FAPESC/CNPq TO TR2012000363 for financial support.

Summary

This study documents the species of lance flies (Diptera: Lonchaeidae) associated with fruits of feijoa (*Acca sellowiana* [Berg] Burret) (Myrtaceae), as well as their infestation level in an orchard located in Lages, Santa Catarina, Brazil. *Neosilba bifida*, *N. delvechioi*, and *N. pradoi* are reported for the first time damaging fruits of feijoa. Moreover, this is the first report of the presence of *N. bella*, *N. bifida*, and *N. delvechioi* from the state of Santa Catarina.

Key Words: feijoa; *Neosilba bifida*; *Neosilba delvechioi*; *Neosilba pradoi*; *Neosilba bella*; frugivorous flies

Sumario

Este estudo apresenta as espécies de moscas-das-frutas (Diptera: Lonchaeidae) associadas a frutos de feijoa (*Acca sellowiana* [Berg] Burret) (Myrtaceae) bem como seu nível de infestação em um pomar em Lages, Santa Catarina, Brasil. *Neosilba bifida*, *N. delvechioi*, e *N. pradoi*

são observadas pela primeira vez danificando frutos de feijoa. Além disso, esta é a primeira vez que *N. bella*, *N. bifida*, e *N. delvechioi* são registradas para o estado de Santa Catarina.

Palavras Chave: feijoa; *Neosilba bifida*; *Neosilba delvechioi*; *Neosilba pradoi*; *Neosilba bella*; moscas frugívoras

References Cited

- Barni EJ, Ducroquet JP, Silva MC, Neto RB, Presser RF. 2004. Potencial de mercado para goiabeira-serrana catarinense. Empresa de Pesquisa Agropecuária e Extensão Rural de Santa Catarina, Florianópolis, Santa Catarina, Brazil.
- Calvo MV, Delgado S, Scatoni I, Garcia, FRM. 2017. First report of *Neosilba pradoi* and *Dasiops friesei* (Diptera: Lonchaeidae) in cultivated and wild hosts in Uruguay. Florida Entomologist 100: 831–832.
- Ducroquet JPHJ, Hickel ER, Nodari RO. 2000. Goiabeira-serrana (*Feijoa sellowiana*). Fundação de Apoio a Pesquisa, Ensino e Extensão, Jaboticabal, São Paulo, Brazil.
- Garcia FRM, Norrbom AL. 2011. Tephritoid flies (Diptera, Tephritoidea) and their plant hosts from the state of Santa Catarina in southern Brazil. Florida Entomologist 94: 151–157.
- Gattelli T, Silva FF, Meirelles RN, Redaelli LR, Dal Soglio FK. 2008. Moscas frugívoras associadas a mirtáceas e laranjeira “Céu” na região do Vale do Rio Caí, Rio Grande do Sul, Brasil. Ciência Rural 38: 236–239.
- Gislotti LJ, Uchoa MA, Prado A. 2017. New records of fruit trees as host for *Neosilba* species (Diptera, Lonchaeidae) in southeast Brazil. Biota Neotropica 17: e20160213. doi.org/10.1590/1676-0611-bn-2016-0213
- Kovaleski A, Sugayama RL, Malavasi A. 1999. Movement of *Anastrepha fraterculus* from native breeding sites into apple orchards in Southern Brazil. Entomologia Experimentalis et Applicata 91: 457–463.
- Lorenzini AR, Boff MIC, Rech TD, Boff P. 2007. Fitogeografia da goiabeira serrana no Planalto Serrano Catarinense. Agropecuária Catarinense 20: 86–89.
- Nunes AM, Müller FA, Gonçalves RS, Garcia MS, Costa VA, Nava DE. 2012. Moscas frugívoras e seus parasitoides nos municípios e Pelotas e Capão do Leão, Rio Grande do Sul, Brasil. Ciência Rural 42: 6–12.
- Pasquariello MS, Mastrobuoni F, Di Patre D, Zampella L, Capuano LR, Scortichini M, Petriccione M. 2015. Agronomic, nutraceutical and molecular variability of feijoa (*Acca sellowiana* (O. Berg) Burret) germplasm. Scientia Horticulturae 191: 1–9.
- Souza GK, Pikart TG, Oliveira VL, Boff P, Boff MIC. 2017. *Acca sellowiana* (Myrtaceae): a new alternative host for *Drosophila suzukii* (Diptera: Lonchaeidae) in Brazil. Florida Entomologist 100: 190–191.
- Souza-Filho MF, Raga A, Azevedo-Filho JA, Strikis PC, Guimarães JÁ, Zucchi RA. 2009. Diversity and seasonality of fruit flies (Diptera: Tephritidae and Lonchaeidae) and their parasitoids (Hymenoptera: Braconidae and Figitidae) in orchards of guava, loquat and peach. Brazilian Journal of Biology 69: 31–40.
- Strikis PC. 2011. Description of 11 new species of genus *Neosilba* (Diptera: Lonchaeidae) from Brazil, its hosts and geographical distribution. Trends in Entomology 7: 67–79.
- Strikis PC, Prado AP. 2005. A new species of genus *Neosilba* (Diptera: Lonchaeidae). Zootaxa 828: 1–5.
- Vargas RI, Peck SL, McQuate GT, Jackson CG, Stark JD, Armstrong JW. 2001. Potential for area wide integrated management of Mediterranean fruit fly (Diptera: Tephritidae) with a braconid parasitoid and a novel bait spray. Journal of Economic Entomology 94: 817–825.
- Weston RJ. 2010. Bioactive products from fruit of the feijoa (*Feijoa sellowiana*, Myrtaceae): a review. Food Chemistry 121: 923–926.