

Rediscovery of *Aedes scapularis* (Diptera: Culicidae) in the Florida Keys

Authors: Hribar, Lawrence J., and Cerminara, Carrie

Source: Florida Entomologist, 104(2) : 145

Published By: Florida Entomological Society

URL: <https://doi.org/10.1653/024.104.0213>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Rediscovery of *Aedes scapularis* (Diptera: Culicidae) in the Florida Keys

Lawrence J. Hribar^{1,*}, and Carrie Cerminara²

Aedes scapularis (Rondani) (Diptera: Culicidae) is a widely distributed mosquito that can be found from southern Texas, to Argentina, and most of the Caribbean islands (Arnell 1976). Pritchard et al. (1947) reported collecting this mosquito from Vaca Key, Florida, USA, an island now within the current city of Marathon (Monroe County), Florida, USA. No further collections of *Ae. scapularis* have been reported in the Keys (Hribar et al. 2016). Recently Reeves et al. (2020) recorded this mosquito on the Florida mainland from multiple localities in Broward and Miami-Dade counties.

On 21 May 2020, 3 female mosquitoes that conformed to the morphological description of *Ae. scapularis* were collected in a dry ice-baited American Biophysics Company light trap (Clarke, St. Charles, Illinois, USA) on Boca Chica Key, Florida. Later, 5 more specimens were collected from the same trap location but collection data were not recorded. All specimens were tentatively identified as *Ae. scapularis* and sent to N. Burkett-Cadena, University of Florida, Vero Beach, Florida, USA, for verification; identification was confirmed independently as *Ae. scapularis* by N. Burkett-Cadena and L. Reeves. The 3 specimens originally collected in May 2020 were deposited into the Florida State Collection of Arthropods (accession number E2021-336-1). On 18 Feb 2021 another female specimen was collected in an ABC light trap on the island of Key West, Florida, USA.

Aedes scapularis is one of the most widely distributed mosquitoes in the Americas, and like other members of the Scapularis group, larvae can be found in a variety of temporary and semi-permanent freshwater habitats including swamp margins and crab holes (Arnell 1976). In Jamaica, *Ae. scapularis* larvae often are found with those of *Aedes tortilis* (Theobald) (Diptera: Culicidae); the latter species is found throughout the Florida Keys (Hribar et al. 2016). Reeves et al. (2020) have reviewed the pertinent literature related to *Ae. scapularis*. As the authors point out, the 2020 collections may indicate that a reintroduction of this species occurred in the recent past. Whether the occurrence of this species at multiple sites in multiple counties represents 1 reintroduction event or several is unknown and likely unknowable. However, its vector status and importance to human and animal health, although not entirely clear, are of concern. Arnell (1976) briefly reviewed the literature pertaining to the medical importance of *Ae. scapularis*. The causative agent of Bancroftian filariasis, *Wuchereria bancrofti* (Cobbold) Seurat (Spirurida: Onchocercidae), has been detected in this mosquito. At least 15 viruses have been isolated from *Ae. scapularis*, including yellow fever, Venezuelan equine encephalitis, and Mayaro viruses, the latter an emerging pathogen in Latin America and the Caribbean with potential to reach North America (Blohm et al. 2019; Mota et al. 2019). Ecological niche modeling suggests that *Ae. scapularis* could become established in more counties in Florida (Campbell et al. 2021). The re-introduction of this mosquito species underscores the importance of continuing surveil-

lance as well as adequate vigilance in identification of those collections to proactively detect the presence of possible invasive vectors.

We thank N. Burkett-Cadena and L. Reeves, Florida Medical Entomology Laboratory, University of Florida, Vero Beach, Florida, USA, for confirming the identification of *Ae. scapularis*.

Summary

The first collections of *Aedes scapularis* (Rondani) (Diptera: Culicidae) from the Florida Keys in 70 yr are reported from Boca Chica Key and Key West, Florida, USA. The importance of this species as a disease vector is briefly discussed.

Key Words: geographic distribution; reintroduction; island; surveillance; Boca Chica Key; species inventory

Sumario

Se registran las primeras colecciones de *Aedes scapularis* (Rondani) (Diptera: Culicidae) en 70 años de Boca Chica Key y Key West, Florida, EE. UU. Se discute brevemente la importancia de esta especie como vector de enfermedades.

Palabras Claves: distribución geográfica; reintroducción; isla; vigilancia; Cayo Boca Chica; inventario de especies

References Cited

- Arnell JH. 1976. Mosquito studies (Diptera, Culicidae). XXXIII. A revision of the Scapularis Group of *Aedes* (*Ochlerotatus*). Contributions of the American Entomological Institute 13 (3): 1–144.
- Blohm G, Elbadry MA, Mavian C, Stephenson C, Loeb J, White S, Telisma T, Chavannes S, De Rocher VM, Salemi M, Lednický JA. 2019. Mayaro as a Caribbean traveler: evidence for multiple introductions and transmission of the virus into Haiti. International Journal of Infectious Diseases 87: 151–153.
- Campbell LP, Burkett-Cadena ND, Miqueli E, Unlu I, Sloyer KE, Medina J, Vasquez C, Petrie W, Reeves LE. 2021. Potential distribution of *Aedes* (*Ochlerotatus*) *scapularis* (Diptera: Culicidae): a vector mosquito new to the Florida peninsula. Insects 2021 12: 213. <https://doi.org/10.3390/insects12030213>.
- Hribar LJ, Leal AL, DeMay DJ, Tambasco AN. 2016. Mosquitoes (Insecta: Diptera: Culicidae) of the Florida Keys, Florida, United States of America. Check List 7: 458–464.
- Mota MT de O, Avilla CM, Nogueira ML. 2019. Mayaro virus: a neglected threat could cause the next worldwide viral epidemic. Future Virology 14: 375–377.
- Pritchard AE, Seabrook EL, Mulrennan JA. 1947. The mosquitoes of the Florida Keys. Florida Entomologist 30: 8–15.
- Reeves LE, Medina J, Miqueli E, Sloyer KE, Petrie W, Vasquez C, Burkett-Cadena ND. 2020. Establishment of *Aedes* (*Ochlerotatus*) *scapularis* (Diptera: Culicidae) in mainland Florida, with notes on the *Ochlerotatus* Group in the United States. Journal of Medical Entomology 58: 717–729.

¹Florida Keys Mosquito Control District, 503 107th Street, Marathon, Florida 33050, USA; Email: lhribar@keysmosquito.org (L. J. H.)

²Florida Keys Mosquito Control District, 18 Aquamarine Drive, Key West, Florida 33050, USA; Email: ccerminara@keysmosquito.org (C. C.)

*Corresponding author; E-mail: lhribar@keysmosquito.org