

Cellular Automata: Simulating Alpine Tundra Vegetation Dynamics in Response to Global Warming

Authors: Zhang, Yanqing A., Peterman, Michael R., Aun, Dorin L., and Zhang, Yanming

Source: Arctic, Antarctic, and Alpine Research, 40(1) : 256-263

Published By: Institute of Arctic and Alpine Research (INSTAAR),
University of Colorado

URL: [https://doi.org/10.1657/1523-0430\(06-048\)\[ZHANG\]2.0.CO;2](https://doi.org/10.1657/1523-0430(06-048)[ZHANG]2.0.CO;2)

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Cellular Automata: Simulating Alpine Tundra Vegetation Dynamics in Response to Global Warming

Yanqing A. Zhang*†§

Michael R. Peterman*

Dorin L. Aun* and

Yanming Zhang‡

*Department of Geography, Simon Fraser University, Burnaby, British Columbia, V5A 1S6, Canada

†School of Computing Science, Simon Fraser University, Burnaby, British Columbia, V5A 1S6, Canada

‡Northwest Plateau Institute of Biology, the Chinese Academy of Science, Xining, Qinghai, 810001, P.R. China

§Corresponding author. Present address: 3-2025 Clarke St., Port Moody, British Columbia, V3H 1Y3, Canada. instca@yahoo.com

Abstract

This study attempts to model alpine tundra vegetation dynamics in a tundra region in the Qinghai Province of China in response to global warming. We used Raster-based cellular automata and a Geographic Information System to study the spatial and temporal vegetation dynamics. The cellular automata model is implemented with IDRISI's Multi-Criteria Evaluation functionality to simulate the spatial patterns of vegetation change assuming certain scenarios of global mean temperature increase over time. The Vegetation Dynamic Simulation Model calculates a probability surface for each vegetation type, and then combines all vegetation types into a composite map, determined by the maximum likelihood that each vegetation type should distribute to each raster unit. With scenarios of global temperature increase of 1 to 3°C, the vegetation types such as Dry *Kobresia* Meadow and Dry *Potentilla* Shrub that are adapted to warm and dry conditions tend to become more dominant in the study area.

DOI: 10.1657/1523-0430(06-048)[ZHANG]2.0.CO;2

Introduction

Global temperature has increased due to the effects of greenhouse gases emissions, and significant impacts have occurred in the biomes and ecosystem types in the region of the Qinghai-Tibetan plateau (Johnes and Briffa, 1992; Ni, 2000; Leemans, 2004). Based on 38 years (1959–1996) of climate observations and statistical analysis, the annual mean temperature increases during this period ranged from 0.4 to 0.6°C in the area of the Haibei Alpine Tundra Ecosystem Research Station (Li et al., 2004). The equilibrium pattern of plateau biomes in response to climate change was modeled using the BIOME3China model (Ni, 2000). However, depicting the vegetation change in response to global warming is a challenge in terms of using spatial modeling to detect and simulate the vegetation response mechanism to warmer weather. In this paper, we model the alpine tundra vegetation changes by integrating the environmental factors in a raster-based Geographic Information System (GIS) environment. Assuming each vegetation type in the raster cell unit reacts as a homogeneous entity, we conduct a spatial and temporal simulation by integrating cellular automata and Multi-Criteria Evaluation (MCE) provided in the IDRISI software (Eastman, 2003).

Spatial modeling processes are available in current GIS software such as IDRISI, which is capable of dealing with a large set of raster data and manipulating the data via operations in a series of discrete time steps, where single raster cells can be influenced by their neighborhood or other data in an overlay. All map layers are imposed on the same grid system. This type of GIS environment provides a sophisticated tool to help us target the real problem in a complex system (Wolfram, 1984; Coulelis, 1985; Itami, 1994; White and Engelen, 2000; Giles, 2002). In our study, we use GIS analysis, linear regression, MCE, cellular automata (CA), and a raster image calculator to build a unique Vegetation Dynamic Simulation Model (VDSM). Global warming scenarios

are interpreted as inputs of the spatial parameters. Large processing tasks are completed by the computer system.

The predicted outcome of this study is that individual vegetation types will respond to a global mean temperature increase (GMTI) in 2100 of 1 or 3°C by either expanding or shrinking their range because of plant species' suitability to the warmer and drier climate conditions. This corresponds to 0.1 or 0.3°C per decade, respectively (Leemans, 2004).

Methods

DATA SET

Our study site is located near the Haibei Alpine Tundra Ecosystem Research Station, Qinghai Province, China (37°29'–37°45'N, 101°12'–101°33'E) (Zhang and Zhou, 1992). The elevation in the study area varies from 3000 to 4500 m a.s.l. The model uses the following data:

- A 90 m × 90 m resolution DEM;
- Temperature derived from the DEM using an empirical linear regression model (Zhang, 2005);
- Land surface parameters (aspect, slope, stream channel density) derived from the DEM using GIS analysis tool provided in the IDRISI software (Eastman, 2003);
- A raster vegetation map (30 m × 30 m pixels) produced in 1988 (Zhang, 2005). The vegetation map with a total of 10 vegetation classes is resampled to match the DEM resolution (90 m × 90 m) (Fig. 1).

MULTI-CRITERIA EVALUATION

Constraints are the limited areas that are not considered to be natural vegetation, such as water bodies, glaciers, gravel slopes,




FIGURE 1. Haibei alpine tundra vegetation distribution, 1988.

and artificial grasslands. They represent areas where the natural vegetation cannot grow or are otherwise constrained. These areas are excluded from the vegetation map. A Boolean image is created to display inclusion and exclusion of the constraint conditions.

$$y = \text{MCE} (F1, F2, F3, \dots Fn) \quad (1)$$

The factors ($F1, F2, \dots Fn$) used in the MCE are selected based on the most important variables that determine the output y in Equation (1). In our study, we use MCE in step 1 (Fig. 2) to determine a normalized surface temperature, which calibrates the temperature by these spatial parameters: aspect, suitable surface temperature, and distance to the nearest stream channel.

The temperature has a certain spatial pattern in the study area: (1) It increases from the north ($a = 0^\circ$) to the southeast ($b = 145^\circ$), with the highest values from the southeast to the southwest ($c = 275^\circ$), then decreasing to the north ($d = 360^\circ$). The change pattern can be described as a sigmoidal fuzzy function type, in a symmetric shape with specific values (a, b, c, d) in Table 1. (2)

The distance to a stream affects the temperature. If the area is within 10 m of a stream or water body, its temperature is closer to the stream or water temperature; beyond 600 m from a stream or water body, the temperature is minimally affected by the nearest water bodies. The values are defined as $a = 10$ m, $b = 600$ m, with a sigmoidal fuzzy function type and monotonically decreasing shape. (3) In the lower valley of our study area, the monthly mean temperature in July is 10.1°C (Li et al., 2004). The temperature decreases with increasing elevation. We define the temperature less than $a = 0^\circ\text{C}$ as unsuitable for alpine plant growth. The temperature from 0°C to $b = 5^\circ\text{C}$ is defined as less suitable for alpine plant growth; the temperature from 5°C to $c = 13^\circ\text{C}$ is defined as most suitable. The temperature from 15.5°C to $d = 18^\circ\text{C}$ reduces the suitability for alpine plant growth, representing the dry, south-facing areas (Table 1). In the Analytical Hierarchy Process (AHP), the most important temperature is in the range of $b = 5^\circ\text{C}$ to $c = 13^\circ\text{C}$, which present the sigmoidal fuzzy function type and symmetrical shape.




FIGURE 2. Vegetation dynamic simulation model.

In step 2 (Fig. 2), we use the MCE method and create a suitability map for each vegetation type using the aspect, slope, and the distance to stream channels. The suitability values (a, b, c, d) in Table 2 are defined based on the vegetation distribution in the study area (Zhang and Zhou, 1992). The fuzzy membership function type and shape and AHP weights for each vegetation are calculated and reported in Table 2 (Eastman, 2003).

MACRO MODELER

The macro model is created to simulate changes in each vegetation type through time and space, integrating operations such as overlay, scalar, fuzzy module, and cellatom (Fig. 3). These operations are available in the IDRISI software package (Eastman, 2003) and have to be built in Macro Modeler with an initial scalar value (0.0–1.0) and fuzzy set values (0–255).

The GMTI scenarios are implemented in the Macro Modeler by adjusting the scalar operation to increase the temperature by 0.1°C in a discrete time period. The same logic is applicable to 0.3°C in a discrete time period (Leemans, 2004). Running the simulation for 10 iterations, the effects of increasing temperature on each vegetation type are accumulated in the output image.

The cellular automata module is implemented in the Macro Modeler (Fig. 3) and used to form a uniform raster image to represent global warming effects in a spatial context, which operates over discrete time steps (Coulelis, 1985; Giles, 2002; White and Engelen, 2000). The change in each cell depends on the parameters or requirements set by the user and the surrounding neighbors (Wolfram, 1984; Itami, 1994; Ruxton and Saravia, 1998). This project used the CELLATOM module with a 3*3 filter and reclassifies an output cell if at least 3 neighbors contain non-null values. We define the 10 iterations using the DynaLink module (Eastman, 2003). Within each iteration, each vegetation map is dynamically updated by running Cellatom, then it is overlaid with the vegetation suitability map altered by a GMTI of 0.1 or 0.3°C. Thus, after 10 iterations, a final suitable vegetation map is produced. The same dynamic processing is repeated for each vegetation type resulting in a total of 10 vegetation suitability maps in response to warmer weather.

COMPOSITE FINAL VEGETATION MAP

Our objective is to create a composite vegetation map for each global warming scenario, GMTI of 1 or 3°C over time (Fig. 2). All of the 10 vegetation suitability maps with a GMTI of 1 or 3°C are combined in order to produce a composite map using the image calculator module in IDRISI (Eastman, 2003). The highest suitability among the vegetation types is selected to represent the successful vegetation type in that cell.

$$\text{Veg_dominant} = \text{MAX}\{\text{Veg1}, \text{Veg2}, \dots, \text{Veg10}\} \quad (2)$$

Equation (2) above creates a map where the value at each pixel corresponds to the vegetation type with the highest probability of thriving, where Veg1, Veg2 ... Veg10 correspond to each of the 10 vegetation types. The 10 resulting maps of dominant vegetation types are overlaid to produce a composite distribution of the most probable vegetation types.

Results

NORMALIZED TEMPERATURE SPATIAL DISTRIBUTION

Temperature changes across the study area are not only due to elevation, but also due to aspect and distance from the nearest stream channel. The linear regression model provided a temperature spatial distribution based on elevation alone, which is our primary step. Furthermore, the normalized temperature surface created by the MCE is highly representative of the potential temperature distribution in a normalized fuzzy format (Fig. 4).

TABLE 1

Normalized surface temperature's fuzzy membership function types and shapes, factor's suitability values (a, b, c, and d), and AHP weights.

Factors to standardize with fuzzy		Membership function type	Membership function shape	a	b	c	d	AHP weights
Temperature	aspect	Sigmoidal	Symmetric	0	145	275	360	0.2631
	temperature suitability	Sigmoidal	Symmetric	−2	5	13	16	0.5472
	distance to stream	Sigmoidal	Monotonically decreasing	10	600	—	—	0.1897

Notes: aspect: 0–360°; temperature suitability: °C; distance to stream: meters; a, b, c, d: suitabilities for the factors; AHP weights derived from pairwise comparison up to an acceptable level.

TABLE 2
Ten vegetation types' fuzzy membership function types and shapes, factor's suitability values (a, b, c, and d), and AHP weights.

Vegetation type	Factors to standardize with fuzzy	Membership function type	Membership function shape	a	b	c	d	AHP weights
Wet <i>Potentilla</i> Shrub	aspect	Sigmoidal	Monotonically Increasing	0	3	—	—	0.5842
	Slope	Sigmoidal	Symmetric	0	10	20	48	0.2318
	distance to stream	Sigmoidal	Symmetric	0	1500	1800	3700	0.184
Dry <i>Potentilla</i> Shrub	aspect	Sigmoidal	Symmetric	0	160	240	360	0.2627
	Slope	Sigmoidal	Monotonically Increasing	0	25	—	—	0.1591
	distance to stream	Sigmoidal	Monotonically Increasing	0	1800	—	—	0.5782
Wet Willow Shrub	aspect	Sigmoidal	Symmetric	0	250	300	360	0.4434
	slope	Sigmoidal	Symmetric	0	15	22	48	0.3874
	distance to stream	Sigmoidal	Symmetric	0	500	1900	3700	0.1692
<i>Canagana</i> Shrub	aspect	Sigmoidal	Symmetric	0	240	255	351	0.4836
	slope	Sigmoidal	Symmetric	0	6	12	20	0.1677
	distance to stream	Sigmoidal	Symmetric	0	500	600	1000	0.3487
Wet <i>Kobresia</i> Meadow	aspect	Sigmoidal	Symmetric	0	240	315	360	0.625
	slope	Sigmoidal	Symmetric	0	10	16	50	0.2385
	distance to stream	Sigmoidal	Monotonically decreasing	—	—	200	600	0.1365
Typical <i>Kobresia</i> Meadow	DEM	Sigmoidal	Monotonically decreasing	—	—	3100	3300	0.2599
	slope	J-shape	Monotonically decreasing	—	—	5	10	0.3275
	distance to stream	Sigmoidal	Symmetric	0	50	1900	3000	0.4126
Dry <i>Kobresia</i> Meadow	aspect	Sigmoidal	Symmetric	0	200	270	300	0.5936
	slope	Sigmoidal	Symmetric	0	1	8	18	0.2493
	distance to stream	Sigmoidal	Symmetric	0	200	1200	1800	0.1571
<i>Carex</i> Meadow	aspect	Sigmoidal	Symmetric	0	70	120	200	0.4934
	slope	Sigmoidal	Symmetric	0	10	15	20	0.3108
	distance to stream	Sigmoidal	Symmetric	0	500	900	1500	0.1958
Riverside <i>Blysmus</i> Meadow	DEM	Sigmoidal	Monotonically decreasing	—	—	3200	3300	0.6483
	slope	Sigmoidal	Monotonically decreasing	—	—	0	3	0.2297
	distance to stream	Sigmoidal	Monotonically decreasing	—	—	0	600	0.122
Riverside <i>Kobresia</i> Meadow	DEM	Sigmoidal	Monotonically decreasing	—	—	3200	3300	0.5396
	slope	Sigmoidal	Monotonically decreasing	—	—	0	10	0.297
	distance to stream	Sigmoidal	Monotonically decreasing	—	—	0	600	0.1634

Note: slope: 0–90°; DEM: indicates elevation range for relatively flat area; for others, refers to the note on Table 1.




FIGURE 3. Example of Macro Modeler incorporating Cellular Automata for *Caragana* Shrub.

VEGETATION CHANGE COMPARISON

We calculate the percent area change for each vegetation type (Fig. 5) before we compose the final vegetation map. By increasing the global mean temperature (0.1 or 0.3°C per decade), the percent area change of each vegetation type indicates the potential expansion from their original ranges. For example, without




FIGURE 4. Normalized temperature map with values from 0 to 255.

competing with other types of vegetation, both Dry *Potentilla* Shrub and Dry *Kobresia* Meadow could expand their vegetation area by 23%. The difference between them is that Dry *Potentilla* Shrub responds more positively to a GMTI of 1°C, and Dry *Kobresia* Meadow responds more positively to a GMTI of 3°C. Similar phenomena are also observed in other vegetation types.

After we compose the final vegetation map, the highest suitability among the vegetation types is selected to represent the successful vegetation type in every cell. For instance, Dry *Potentilla* and Dry *Kobresia* expand into areas previously occupied by wet types of vegetation (Fig. 6). The Riverside *Blysmus* meadow, which requires moist conditions, disappears completely with a 3°C temperature increase. In general, the dry vegetation types demonstrate significant expansion from their original ranges and tend to become more dominant in the study area.

VEGETATION DYNAMICS OVER TIME

The time dimension in the CA module of IDRISI is interpreted as a discrete time step, which corresponds to a temperature-time dimension. The CA module is integrated into the macro modeler, and uses the DynaLink module (Eastman, 2003) to simulate vegetation change within each iteration. The GMTI is defined as a spatial parameter in the deterministic model with a temperature increase of 0.1 or 0.3°C per decade.




FIGURE 5. Percent change in vegetation area with GMTI at 1 and 3 °C.




FIGURE 6. Final composite vegetation maps for GMTI at 1 and 3 °C.

VEGETATION DYNAMIC SIMULATION MODEL (VDSM)

The VDSM is an example showing that spatial modeling can solve complex ecosystem problems in terms of having the capability to simulate spatial and temporal vegetation dynamics. In this case study, we model the spatial distribution of temperature and create GMTI scenarios as a spatial grid image. The vegetation dynamics are simulated in discrete time by applying CA in a Macro Modeler. In future studies, this model will be capable of modeling the water-time dimension that makes the simulation more adaptable to global warming research. The VDSM could be potentially incorporated with a normal climate change model to assist in a long-term ecosystem simulation. Alternatively, the VDSM is capable of linking with a stochastic model of temperature change, in which we might be able to forecast an ecosystem disaster.

Discussions

TEMPERATURE DISTRIBUTION

Temperature distribution is correlated with and controlled primarily by elevation. Numerous spatial interpretation methods have been applied to estimate the spatial distribution of temperature (Li, 2005). The interpolation results do not always agree with the actual sample points, including using geo-statistical methods, and spatio-temporal spline. These methods are highly dependent on the distance to the sample points, and the surface equation. In our study, our first step is to create the primary temperature surface based a linear relation with elevation. The objective is to obtain a more accurate temperature map in terms of aspect, suitable temperature, and distance to the stream. We use

the Multi-Criteria Evaluation with Weighted Linear Combination (MCE_WLC) to calibrate the spatial temperature distribution. The fuzzy memberships between the temperature and each factor (aspect, suitable temperature, distance to stream) are based on previous research works (Zhang and Zhou, 1992; Zhang and Welker, 1996; Zhang, 2005). The output, the normalized temperature surface is set into fuzzy format (0–255). Since the temperature is major factor on determining vegetation composition, structure, and distribution, the normalized temperature surface plays an important role when we simulate vegetation dynamics in spatial and temporal dimensions.

ERROR AND FUZZY SETS

Data from multiple sources are unlikely to be the same resolution, scale, or nominal context. The errors occurred during data conversion, and propagated through the simulation. This model did not take competition among plant species and grazing effects into account (Zhang, 1990; Wu et al., 1996). Clearly, some species are more aggressive and dominant than others, and this would affect their probability of survival when compared to other less aggressive plant species. The lack of a grazing factor resulted in some uncertainty in the modeled vegetation changes.

Fuzzy membership function types and shapes for the factors are based mainly on previous research. We had difficulty validating the accuracy of values of a, b, c, d on Tables 1 and 2. But, the weights of AHP illustrate the importance of each factor in terms of determining the vegetation distribution. For example, the *Wet Potentilla* Shrub is distributed on north-facing slopes, where the north aspect is more important than the slope and the distance

to a stream (e.g., aspect weight 0.5842 > slope weight 0.2318 > distance to stream weight 0.184). Similarly, the Wet Willow Shrub, *Caragana* Shrub, Wet *Kobresia* Meadow, Dry *Kobresia* Meadow, and *Carex* Meadow are distributed and correlated with the different aspects of the terrain. However, the Dry *Potentilla* Shrub and Typical *Kobresia* Meadow are distributed on open valley, and the aspect and slope become less important to their distributions. Also, the Riverside *Blasmus* Meadow and Riverside *Kobresia* Meadow are mainly distributed between 3200 and 3300 m within a buffer area 600 m from the stream or water bodies. As a result, the spatial factors' fuzzy sets agree strongly with the vegetation distributions.

VEGETATION DYNAMIC SIMULATION MODEL

The VDSM starts from thinking about how to solve a complex vegetation dynamics problem using CA. The VDSM is built by combining the MCE, Macro Modeler, CA, image calculator, Scalar, and Fuzzy functions in IDRISI. Making a clear objective helps us to look into many available modelers and functions in order to solve the problem within the IDRISI software environment. For instance, using Decision Wizard, we create an objective and define a set of constraints, which eliminates the areas that are not natural vegetation. The spatial patterns of the factors (temperature, aspect, slope, and distance to stream) are created as continuous surfaces using Fuzzy functions. The transition rules in Macro Modeler are defined as the maximum potential suitable vegetation in each cell as well as over the study area. The frame work of VDSM (Fig. 2) is a summary of the model structure and functionalities. VDSM is flexible enough to be integrated with other sub-models that are available from GIScience technology. Figure 3 provides an example of Macro Modeler incorporating Cellular Automata for the modeling of *Caragana* Shrub. In the case of a GMTI of 0.1°C, the suitability map for *Caragana* is weighted at 0.4 and the normalized temperature map is weighted at 0.6. These two factors are combined using the overlay module to produce a map of suitability for *Caragana* Shrub with an incremental temperature increase of 0.1°C. Ten iterations on one vegetation layer are simulated and updated after each iteration using the Cellatom and Dynalink modules. At the end of the iterations, we obtained an accumulated effect of GMTI of 1°C on the vegetation layer.

It would also be possible to incorporate the water layer with the vegetation layer and temperature layer, which can be linked by its weighted value in the VDSM. Thus, the VDSM not only provides a discrete time representation, but also demonstrates how we could develop our model for use with more complex scenarios (Chapin et al., 2006; McGuire et al., 2006).

Compositing a final vegetation map demonstrates the power of GIS analysis in IDRISI. The image calculator module in IDRISI successfully carries out interpolation of the image calculation presented in Equation (2) at the grid cell level. The VDSM illustrates how we could study vegetation dynamics and model many other spatio-temporal phenomena.

Conclusions

The VDSM integrates the suitability maps created from MCE, Macro-Modeler, CA, and spatial environmental factors. And the temperature-time dimension model is incorporated into the VDSM, which makes the temperature a spatial parameter that affects the vegetation dynamics over a discrete time step. The simulating processes conducted by Macao Modeler generate the temperature

increase of 0.1 to 0.3 °C per decade, which represents the influences of the different global warming scenarios. The results from Figures 5 and 6 demonstrate that global temperature increase reduces moisture availability (Zhang and Welker, 1996) such that dry vegetation can invade areas previously occupied by vegetation adapted to moist conditions. The structure of the model is generally applicable to other situations, but the particular factors and constraints used in this model are unique to the Haibei alpine tundra ecosystem.

Global warming has strong effects on the alpine ecosystems in terms of altering the biomes and ecosystem biodiversity (Zhang, 1993; Ni, 2000; Song et al., 2005). The alpine ecosystem in the region of the Qinghai-Tibetan plateau is sensitive and vulnerable to the changing climate (Zhang and Welker, 1996). The VDSM illustrates that altering global mean temperature changes the alpine vegetation dynamics. With the future integration of water condition (Hodkinson et al., 1999) and disturbance regimes (Zhang, 1990; Zhang and Liu, 2003; Chapin et al., 2006) into the VDSM, the simulation could model more detailed mechanisms and complex feedbacks (McGuire, 2006) of the alpine tundra ecosystem to the changing climate.

Acknowledgments

We thank Dr. Dragicevic, Verda Kocabas, and the Simon Fraser University Spatial Information Systems (SIS) lab for their support of this research project. We thank the two reviewers who gave us the critical suggestions to improve our revised manuscripts. We thank the Northwest Plateau Institute of Biology, the Chinese Academy of Sciences, for academic support and encouragement of this project.

References Cited

- Chapin, F. S., III, McGuire, A. D., Randerson, J., Pielke, R. Sr., Baldocchi, D., Hobbie, S. E., Roulet, N., Eugster, W., Kasischke, E., Rastetter, E. B., Zimov, S. A., and Running, S. W., 2000: Arctic and boreal ecosystems of western North America as components of the climate system. *Global Change Biology*, 6: 211–223.
- Chapin, F. S., III, Robards, M. D., Huntington, H. P., Johnstone, J. F., Trainor, S. F., Kofinas, G. P., Ruess, R. W., Fresco, N., Natcher, D. C., and Naylor, R. L., 2006: Directional changes in ecological communities and social-ecological systems: a framework for prediction based on Alaskan examples. *The American Naturalist*, 168: S36–S49.
- Coulelis, H., 1985: Cellular world: a framework for modeling micro-macro dynamics. *Environment and Planning A*, 17: 585–596.
- Eastman, J. R., 2003: *IDRISI Kilimanjaro Tutorial*. Manual Version 14.0. Worcester, Massachusetts: Clark Labs of Clark University, 61–123.
- Giles, J., 2002: What kind of science is this? *Nature*, 417: 216–218.
- Hodkinson, I. D., Webb, N. R., Bale, J. S., and Block, W., 1999: Hydrology, water availability and tundra ecosystem function in a changing climate: the need for a closer integration of ideas? *Global Change Biology*, 5(3): 359–369.
- Itami, R. M., 1994: Simulating spatial dynamics: cellular automata theory. *Landscape and Urban Planning*, 30: 27–47.
- Johnes, P. D., and Briffa, K. R., 1992: Global surface air temperature variations during the twentieth century. Part I: spatial, temporal and seasonal details. *The Holocene*, 2: 165–179.
- Leemans, R. E., 2004: Another reason for concern: regional and global impacts on ecosystems for different levels of climate change. *Global Environmental Change*, 14: 219–228.
- Li, X., Cheng, G. D., and Lu, L., 2005: Spatial analysis of air temperature in the Qinghai-Tibet Plateau. *Arctic, Antarctic, and Alpine Research*, 37(2): 246–252.

- Li, Y. N., Zhao, X. Q., Cao, G. M., Zhao, L., and Wang, Q. X., 2004: Analysis on climates and vegetation productivity background at Haibei Alpine Meadow Ecosystem Research Station. *Plateau Meteorology*, 23(4): 558–567.
- Ni, J., 2000: A simulation of biomes on the Tibetan Plateau and their responses to global climate change. *Mountain Research and Development*, 20(1): 80–89.
- McGuire, A. D., Chapin, F. S., III, Walsh, J. E., and Wirth, C., 2006: Integrated regional changes in arctic climate feedbacks: implications for the global climate system. *Annual Review of Environmental Resources*, 31: 61–91.
- Ruxton, G. D., and Saravia, L. A., 1998: The need for biological realism in the updating of cellular automata models. *Ecological Modeling*, 107(2–3): 105–112.
- Song, M. H., Zhou, C. P., Ouyang, H., 2005: Simulated distribution of vegetation types in response to climate change on the Tibetan Plateau. *Journal of Vegetation Science*, 16(3): 341–350.
- White, R., and Engelen, G., 2000: High-resolution integrated modeling of the spatial dynamics of urban and regional system. *Environment and Urban Systems*, 24: 383–400.
- Wolfram, S., 1984: Cellular automata as models of complexity. *Nature*, 311(4): 419–424.
- Wu, H., Li, B. L., Stoker, R., and Li, Y., 1996: A semi-arid grazing ecosystem simulation model with probabilistic and fuzzy parameters. *Ecological Modeling*, 90: 147–160.
- Zhang, X. S., 1993: The Tibetan Plateau in relation to the vegetation of China. *Annals of the Missouri Botanical Garden*, 70: 564–570.
- Zhang, X. S., Yang, D. A., Zhou, G. S., Liu, C. Y., and Zhang, J., 1996: Model expectation of impacts of global climate change on biomes of the Tibetan Plateau. In Omasa, K., Kai, K., Taoda, H., Uchijima, Z., and Yoshino, M. (eds.), *Climate change and plants in East Asia*. Tokyo: Springer-Verlag, 25–38.
- Zhang, Y. M., and Liu, J. K., 2003: Effects of plateau zokors (*Myospalax fontanierii*) effects on plant community and soil in an alpine meadow. *Journal of Mammalogy*, 84(2): 644–651.
- Zhang, Y. Q., 1990: A quantitative study on the characteristics and succession pattern of alpine shrublands under the different grazing intensities. *Acta Phytocologia and Geobotanica Sinica*, 14(4): 358–365.
- Zhang, Y. Q., 2005: Raster multi-criteria evaluation for experimental design with aggregating weighed factors. *GeoTec Event Proceeding*. Poster Presentation in Vancouver Canada. (<http://www.sfu.ca/geog/geog355fall04/yqz/index.htm>).
- Zhang, Y. Q., and Welker, J. M., 1996: Tibetan alpine tundra responses to simulated changes in climate: aboveground biomass and community responses. *Arctic and Alpine Research*, 28(2): 203–209.
- Zhang, Y. Q., and Zhou, X. M., 1992: The quantitative classification and ordination of Haibei alpine meadow. *Acta Phytocological ET Geobotanica Sinica*, 16(1): 36–42.

Ms accepted March 2007