

Contrasting Responses of Nitrogen-Fixation in Arctic Lichens to Experimental and Ambient Nitrogen and Phosphorus Availability

Authors: Weiss, Marissa, Hobbie, Sarah E., and Gettel, Gretchen M.

Source: Arctic, Antarctic, and Alpine Research, 37(3) : 396-401

Published By: Institute of Arctic and Alpine Research (INSTAAR),
University of Colorado

URL: [https://doi.org/10.1657/1523-0430\(2005\)037\[0396:CRONIA\]2.0.CO;2](https://doi.org/10.1657/1523-0430(2005)037[0396:CRONIA]2.0.CO;2)

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

Contrasting Responses of Nitrogen-Fixation in Arctic Lichens to Experimental and Ambient Nitrogen and Phosphorus Availability

Marissa Weiss*†‡

Sarah E. Hobbie* and

Gretchen M. Gettel†

*Department of Ecology, Evolution, and Behavior, University of Minnesota, 1987 Upper Buford Circle, St. Paul, MN 55108, U.S.A.

†Department of Ecology and Evolutionary Biology, Cornell University, Corson Hall, Ithaca, NY 14853, U.S.A.

‡msw27@cornell.edu

Abstract

We investigated the influence of nitrogen (N) and phosphorus (P) on N₂-fixation and abundance of two of the most common N₂-fixing arctic lichens, *Peltigera aphthosa* and *P. polydactyla*, in two common moist upland tundra types, acidic and non-acidic tundra, at Toolik Lake, Alaska. Acidic tundra has higher N and lower P availability than non-acidic tundra. We measured the abundance of the lichens in control (no fertilization), N- and P-fertilized plots, and N₂-fixation using the acetylene reduction assay method on lichens from control and P-fertilized plots from both tundra types. Lichens on N-treated plots were too scarce to include in our N₂-fixation estimates. Lichen abundance was lower in plots fertilized with N than in control and P-fertilized plots, while per-biomass N₂-fixation rates were higher in P-fertilized plots than in control plots. Per-biomass rates of N₂-fixation did not differ between acidic and non-acidic tundra, but both lichen species are more abundant on acidic tundra. Thus, despite per-biomass stimulation of N₂-fixation by experimental P addition and reduction in lichen abundance with N fertilization, *Peltigera* contributes more N to the acidic tundra site, indicating that soil N and P availability are not the primary controls of N₂-fixation and abundance of these lichens.

Introduction

Nitrogen-fixation is an important source of new nitrogen (N) to arctic ecosystems because N inputs from atmospheric deposition are very low (Alexander and Schell, 1973; Alexander, 1974; Barsdate and Alexander, 1975; Chapin et al., 1991; Chapin and Bledsoe, 1992). Yet, surprisingly little research on N₂-fixation has been done in upland tundra, despite its widespread distribution in the circumpolar Arctic (Bliss and Matveyeva, 1992; Walker, 1998). Because N₂-fixation is a physiological process subject to abiotic and biotic controls, determining influences on N₂-fixation at the organismal level can offer insights into the important controls of N₂-fixation at the ecosystem level (Chapin and Bledsoe, 1992).

Lichens are major components of arctic ecosystems; thus, the contribution of N₂-fixing lichens to arctic nutrient cycles is likely substantial (Nash and Olafsen, 1995). Additionally, N₂-fixing lichens may be particularly important in supplying new N in extreme environments where N₂-fixing plants are rare (Gunther, 1989). N₂-fixation in arctic lichens occurs in species that host cyanobacterial symbionts *Nostoc* or *Anabaena*, and has been measured in several lichen genera, including *Peltigera*, *Nephroma*, and *Stereocaulon* in the arctic and alpine tundra (Alexander, 1974).

N₂-fixation is inhibited by increasing available N (Alexander et al., 1978; Chapin et al., 1991; Vitousek and Howarth, 1991; Chapin and Bledsoe, 1992; Vitousek and Field, 1999). Phosphorus (P) stimulates N₂-fixation, and low P availability constrains N₂-fixation (Chapin et al., 1991; Chapin and Bledsoe, 1992; Crews, 1993; Hendzel et al., 1994; Bowman et al., 1996; Thompson and Vitousek, 1997; Kurina and Vitousek, 1999; Vitousek, 1999; Vitousek and Field, 1999; but see Alexander et al., 1978). These studies support the suggestion that N₂-fixation may be limited by P, and that the ratio of N to P availability is important in regulating N₂-fixation (Vitousek and Howarth, 1991). In spite of substantial work at other sites, the influence of N or P availability on N₂-fixation in upland tundra has not previously been

investigated. Furthermore, it is unclear whether N and P have similar influences over growth, abundance, and N₂-fixation activity of lichens.

Here we investigate the influence of N and P on abundance and N₂-fixation of two N₂-fixing arctic lichens, *Peltigera aphthosa* and *P. polydactyla*, in moist acidic and moist non-acidic tundra near Toolik Lake, Alaska. Moist acidic and non-acidic tundra are the two most common tundra types in the region (Walker, 1998), and they differ greatly in pH, plant species composition, biomass (Walker et al., 1994; Walker et al., 1995), and N and P availability (Hobbie and Gough, 2002). We focus on these particular lichens because previous studies have demonstrated significant N₂-fixation in these species at other sites (Alexander, 1974; Chapin and Bledsoe, 1992), and lichen surveys indicated that these two species are more abundant and have higher rates of N₂-fixation than other N₂-fixing lichens in tundra near Toolik Lake.

We measured the influence of N and P availability on lichen abundance by comparing lichen cover among plots receiving either no fertilization (control), or N or P fertilization. We also compared lichen cover between acidic and non-acidic tundra. We compared lichen N₂-fixation rates in the two species between control and P fertilized plots at both sites. N treatment plots were not included in the N₂-fixation comparison because lichens were too scarce. Specifically, we address the following hypotheses:

- (1) Fertilization with N causes a decline in lichen abundance and therefore a decline in the contribution of lichens to N₂-fixation per unit area of tundra.
- (2) Rates of N₂-fixation by lichens are stimulated by fertilization with P.
- (3) Lichens from acidic tundra, where N availability is greater, have lower rates of N₂-fixation and are less abundant than lichens from non-acidic tundra where P availability is higher; thus the contribution of lichens to N₂-fixation per unit area of tundra is lower in acidic tundra compared to non-acidic tundra.




FIGURE 1. Percent cover for *Peltigera aphthosa* and *P. polydactyla* in control (C), phosphorus (P), and nitrogen (N) treatments on acidic and non-acidic tundra. Quadrats were averaged within plots, and data presented are plot means for each treatment, with standard error bars ($n = 4$ for acidic tundra or 3 for non-acidic tundra).

Site Description

Research was conducted at the Arctic Long Term Ecological Research (LTER) site at Toolik Lake, in the northern foothills of the Brooks Range, Alaska ($68^{\circ}38'N$, $149^{\circ}38'W$). We studied lichens on moist acidic and moist non-acidic tundra adjacent to Toolik Lake. Because the two sites are <2 km apart, they experience the same climate conditions. However, they differ considerably in substrate age, pH, vegetation, and N and P availability (Hobbie and Gough, 2002). Moist acidic tundra is found on older glacial surfaces, while moist non-acidic tundra is found on younger surfaces or surfaces that receive relatively high deposition of loess (M. D. Walker et al., 1994; D. A. Walker et al., 1998). The acidic tundra site investigated here was deglaciated approximately 50,000–120,000 years ago, while the non-acidic site was deglaciated approximately 11,500–25,000 years ago (Hamilton, 2003). Moist acidic tundra has a soil pH of 3–4. The vegetation is mostly comprised of sedges, dwarf shrubs, mosses, and fruticose lichens. Dominant species include the vascular plants *Eriophorum vaginatum*, *Betula nana*, *Carex bigelowii*, *Ledum palustre*, and *Vaccinium vitis-idaea*; the mosses *Sphagnum* spp. and *Hylocomium splendens*; and the lichens *P. aphthosa*, *P. polydactyla*, and (in relatively low abundance) *Nephroma arcticum*. Non-acidic tundra has a circumneutral pH (6–7) and is dominated by dwarf shrubs, sedges, and lichens. Forbs are more abundant and diverse in non-acidic than acidic tundra. Major species include the vascular plants *Eriophorum angustifolium* ssp. *triste*, *Carex bigelowii*, *Dryas integrifolia*, and *Salix reticulata*; the moss *Tomen-thypnum nitens*; and the lichens *P. aphthosa* and *P. polydactyla* (Walker et al., 1994). Net N mineralization rates are higher in acidic tundra than in non-acidic tundra, while cation exchange capacity, extractable calcium, and extractable P in the mineral soil are higher in non-acidic tundra (Hobbie and Gough, 2002). Plant biomass is significantly greater in acidic tundra than in non-acidic tundra in this region (Walker et al., 1995; S. E. Hobbie, L. Gough, and G. R. Shaver, unpublished data). N limits production in acidic tundra, while N and P together strongly

TABLE 1

Three-way ANOVA comparing abundance of *Peltigera aphthosa* and *P. polydactyla* between sites, and among control (C), nitrogen (N), and phosphorus (P) treatments, with Tukey's HSD posthoc test. Three-way ANOVA comparing N_2 -fixation rates per unit area and per unit mass between *P. aphthosa* and *P. polydactyla*, acidic and non-acidic tundra, and control and P treatments.

Source	Abundance		N_2 -fixation per m^2 lichen		N_2 -fixation per g lichen	
	F	P	F	P	F	P
Site	5.35 _{1,30}	0.03	0.26 _{1,16}	0.61	0.09 _{1,16}	0.77
Treatments	18.79 _{2,30} *	0.00	8.62 _{1,16} †	0.01	8.98 _{1,16} †	0.01
Species	31.48 _{1,30}	0.00	0.19 _{1,16}	0.67	2.68 _{1,16}	0.12
Site \times Treatment	3.26 _{2,30}	0.05	0.44 _{1,16}	0.52	0.07 _{1,16}	0.79
Site \times Species	8.79 _{1,30}	0.01	0.70 _{1,16}	0.42	0.94 _{1,16}	0.35
Species \times Treatment	0.20 _{2,30}	0.82	0.37 _{1,16}	0.55	0.89 _{1,16}	0.36
Site \times Species \times Treatment	0.26 _{2,30}	0.77	0.62 _{1,16}	0.44	0.44 _{1,16}	0.52
Treatment Posthoc Comparisons:						
C and P		0.73				
P and N		0.00				
N and C		0.00				

* Treatments = C, N, P.

† Treatments = C, P.

limit production in non-acidic tundra (Shaver et al., 1986; Gough and Hobbie, 2003).

Methods

FERTILIZATION EXPERIMENTS

We studied lichens in plots treated with N or P fertilizer in both acidic and non-acidic tundra. Treatment plots are 5×20 m and have received either no fertilization (control) or fertilization as single annual dry applications of granular fertilizer immediately following snowmelt, with N as NH_4NO_3 or P as P_2O_5 at a rate of $10 \text{ g N m}^{-2} \text{ y}^{-1}$ and $5 \text{ g P m}^{-2} \text{ y}^{-1}$, respectively (Bret-Harte et al., 2001). There are four blocks at the acidic tundra site and three blocks at the non-acidic site. Fertilization began in 1989 on acidic tundra and in 1997 on non-acidic tundra as part of the core LTER experiments.

LICHEN COVER SURVEYS

Lichen abundance was quantified by estimating percent lichen cover at ambient lichen field moisture within the experimental fertilization plots and outside of the fertilization plots at acidic and non-acidic study sites. Percent lichen cover was estimated using a 1-m^2 quadrat that was subdivided into a 25-box grid and used to estimate the percent of the quadrat area that was covered by the two lichens. Within each of the 21 fertilization plots, we surveyed six 1-m^2 quadrats. *Peltigera* area values from six quadrats were averaged for each plot, and the averages for the plots were compared among treatments and between sites and species (21 plots \times 2 species = 42 observations). Outside of the fertilization plots, percent cover in 1-m^2 quadrats was determined every 5 m along eight 50-m transects at each site (10 quadrats per transect \times 8 transects \times 2 sites \times 2 species = 320 observations). These transects were parallel to the slope and were approximately 5 m apart. Because this is a nondestructive measure, we assume that any differences in size due to variability in ambient lichen moisture on measurement days are negligible, certainly less than the precision of the visual percent cover measurement.




FIGURE 2. Acetylene reduction rates for *Peltigera aphthosa* and *P. polydactyla* in control (C) and phosphorus (P) treatments on acidic and non-acidic tundra. Values are means with standard error bars ($n = 3$).

We compared cover among fertilization plots using a three-way ANOVA, with site, treatment, and species as factors, and we used Tukey's HSD posthoc test to make pairwise comparisons among treatments. We compared lichen cover between acidic and non-acidic tundra using two-way analysis of variance (ANOVA) with site and species as factors.

ACETYLENE REDUCTION ASSAYS

N_2 -fixation in the lichens was estimated using the acetylene reduction assay (ARA) (Hardy et al., 1968). Lichens were collected within 24 h of the start of the ARA, placed in plastic bags, and transferred into uncapped 250-mL mason jars within 5 h of collection. The mason jars were modified by punching a hole in each lid and fitting the hole with a silicone Hungate septum. In preparation for the incubation, lichens were hydrated in water from Toolik Lake for 2 min and drained for 1 min to ensure moisture was not limiting and that all lichens were experiencing the same moisture conditions. We used sufficient biomass of lichen to cover the surface of the 24-cm² mason jar lid. Using a syringe, 20 mL of air were removed from the chamber, and 20 mL of acetylene were injected into the chamber. We incubated the lichens under constant light (285 $\mu\text{mol m}^{-2} \text{s}^{-1}$, compact fluorescent bulbs) in a water bath that maintained the temperature in the range of 11.5–12.5°C, which falls within one standard deviation of the mean air temperature ($9.6 \pm 5.1^\circ\text{C}$) during the growing season (10 June to 15 August) in 2000 and 2001 (Arctic LTER database, 2002). The jars were inverted for the incubation, orienting the lichen surface to face the light source. We first did several preliminary experiments. We analyzed acetylene blanks directly from our acetylene supply, and noted minimal contamination of ethylene in our acetylene. That small amount was subtracted from the final measure of ethylene when analyzing data. We incubated lichens with no acetylene and measured no ethylene production in 10 h, and we incubated empty jars with acetylene and measured no ethylene production. After establishing that the rate of N_2 -fixation is linear between 2 and 10 h of incubation, we incubated lichens for 2 to 6 h, at which time a 5-mL gas sample was removed from the jars and analyzed for ethylene using a gas chromatograph (Shimadzu GC-8A, Shimadzu Scientific Instruments, Inc., Columbia, Maryland) with flame-ionization detection (FID) and a Porapak N column. After the incubation, lichens were dried at 65°C for at least 36 h, weighed, and the rate of acetylene reduction was calculated on a per-unit moist lichen area and per-unit dry mass basis.

On 21 July 2000, 24 lichens were collected to compare acetylene reduction rates between control and P fertilized plots. We were unable to harvest sufficient lichens from the N fertilized plots. The experiment is a $2 \times 2 \times 2$ factorial design with three replicates of each species by site by treatment combination. We compared rates of N_2 -fixation in lichens from fertilization plots using a three-way ANOVA with site, treatment, and species as factors. The N concentration of each incubated lichen was determined after the conclusion of the experiment in finely ground oven-dried lichen tissue using a Perkin-Elmer CHN Analyzer (Perkin-Elmer, Boston, Massachusetts).

A total of 72 lichens were collected from outside of the treatment plots on 2 July, 19 July, and 3 August 2001 to compare acetylene reduction rates between the acidic and non-acidic tundra sites. Six replicates of each species from each site were collected on each of the three dates (3 dates \times 2 species \times 2 sites factorial design, with 6 replicates). We compared N_2 -fixation in lichens from the two sites using a three-way ANOVA with site, species, and date as factors.

Results

FERTILIZATION PLOTS: RESPONSE OF PELTIGERA TO EXPERIMENTAL N AND P

The abundance of both *P. aphthosa* and *P. polydactyla* was reduced in the N fertilization plots at both the acidic and non-acidic tundra sites (Fig. 1, Table 1). In fact, lichens were virtually eliminated from plots fertilized with N at the acidic site. In contrast, P addition had no effect on lichen abundance at either site (Fig. 1, Table 1). In the analyses of abundance from the treatment plots, *P. aphthosa* was significantly more abundant at the non-acidic site than at the acidic site, while the abundance of *P. polydactyla* did not differ between the two sites (Fig. 1, Table 1). However, the area within the treatment plots was likely too small to adequately characterize these sites for lichen abundance, and the site comparison of abundance outside of the treatment plots better represents true site differences (see below).

Rates of N_2 -fixation in lichens from plots receiving P were significantly greater than in lichens from control plots (Fig. 2, Table 1) on both a per-lichen area and per-lichen biomass basis, although trends were stronger for N_2 -fixation expressed per unit lichen mass for *P. aphthosa* (Table 1). Because of this similarity, graphs show only the per unit mass data, though the analyses for both are presented (Tables 1 and 2). Thallus N concentration was higher in lichens fertilized by P

TABLE 2

Two-way ANOVA comparing abundance of *Peltigera aphthosa* and *P. polydactyla* between moist acidic and moist non-acidic tundra. Three-way ANOVA comparing N₂-fixation rates per unit area and per unit mass between *P. aphthosa* and *P. polydactyla*, between moist acidic and moist non-acidic tundra, and among the three sampling dates.

Source	Abundance		N ₂ -fixation per m ² lichen		N ₂ -fixation per g lichen	
	F	P	F	P	F	P
Site	58.17 _{1,316}	0.00	0.01 _{1,59}	0.92	0.25 _{1,59}	0.62
Species	6.19 _{1,316}	0.01	5.72 _{1,59}	0.02	0.20 _{1,59}	0.66
Date			8.42 _{2,59}	0.00	7.31 _{2,59}	0.00
Site × Date			2.91 _{2,59}	0.06	2.20 _{2,59}	0.12
Date × Species			1.59 _{2,59}	0.21	0.67 _{2,59}	0.51
Site × Species	19.01 _{1,316}	0.00	0.16 _{1,59}	0.69	0.18 _{1,59}	0.67
Site × Species × Date			0.99 _{2,59}	0.38	0.28 _{2,59}	0.76

(Fig. 3, Table 3), which corresponds with higher lichen N₂-fixation in plots receiving P. Though we were unable to characterize the response of N₂-fixation rates to N fertilization, clearly the contribution of N by *Peltigera* to the N fertilized plots is not substantial.

SITE COMPARISON: RESPONSE OF PELTIGERA TO AMBIENT N AND P

While 72 lichens were incubated with acetylene, one gas sample, a replicate of *P. aphthosa* on non-acidic tundra, was lost due to a leak in an incubation chamber. As a result, our analysis is based on the remaining 71 gas samples.

In contrast to our hypothesis, abundance of both species, but especially *P. polydactyla*, was greater at the acidic site than at the non-




FIGURE 3. Nitrogen concentrations in *Peltigera aphthosa* and *P. polydactyla* from control (C) and phosphorus (P) treatments on acidic and non-acidic tundra. Values are means with standard error bars ($n = 3$).

TABLE 3

Three-way ANOVA comparing lichen N concentration between *Peltigera aphthosa* and *P. polydactyla* and between the control (C) and phosphorus (P) treatments at the acidic and non-acidic sites.

Source	F	P
Site	4.47 _{1,16}	0.05
Treatment	51.45 _{1,16}	0.00
Species	93.65 _{1,16}	0.00
Site × Treatment	0.48 _{1,16}	0.50
Site × Species	1.30 _{1,16}	0.27
Species × Treatment	3.91 _{1,16}	0.07
Site × Species × Treatment	0.12 _{1,16}	0.73

acidic site (Fig. 4, Table 2). However, both species of lichens had similar rates of N₂-fixation at the two sites (Fig. 5, Table 2). Thus, despite similar rates of per-biomass N₂-fixation at the two sites, the contribution of N by N₂-fixation in *P. aphthosa* and *P. polydactyla* is greater in acidic tundra than in non-acidic tundra. Though N₂-fixation rates were significantly higher on 2 July than 19 July and 3 August 2001, there were no significant interactions of date with species or site (Table 2), so we present only the seasonal averages in Figure 4.

Discussion

Our results present an interesting paradox. With P fertilization, the average N₂-fixation rate for both species increased from 0.77 ± 0.19 $\mu\text{moles acetylene g lichen}^{-1} \text{ h}^{-1}$ to 1.65 ± 0.39 $\mu\text{moles acetylene g lichen}^{-1} \text{ h}^{-1}$. However, N₂-fixation rates did not vary between the more P-rich non-acidic site and the acidic site with higher N availability. Furthermore, *Peltigera* percent cover is threefold greater at the acidic site, with percent cover values of 6% and 2% at the acidic site and non-acidic site, respectively. Thus, while P fertilization approximately doubles the rate of N₂-fixation per gram lichen, the areal contribution of N fixed by *Peltigera* under ambient nutrient conditions is three times less on the more P-rich non-acidic site.

Potential explanations for our contradictory results include soil moisture, soil pH, micronutrient limitation, and the physiological constraints of *Peltigera*. Chapin et al. (1991) measured a positive response of cyanobacterial N₂-fixation to P fertilization in a high-arctic lowland, but were unable to measure differing rates of N₂-fixation along a natural P gradient. They attributed this result to differences in moisture among the sites that overrode the influence of P on N₂-fixation rates. However, the two sites studied here do not differ significantly in soil moisture (L. Gough, unpublished data). There is an obvious difference in pH between these two sites, though the pH




FIGURE 4. Percent cover of *Peltigera aphthosa* and *P. polydactyla* on acidic and non-acidic tundra. Values are means with standard error bars ($n = 80$).

optimum for N_2 -fixation in *P. aphthosa* is 5–7 (Englund, 1978), suggesting that rates of N_2 -fixation would be higher on non-acidic tundra if pH were an important factor influencing in situ rates of N_2 -fixation there. N_2 -fixation may be limited by a micronutrient such as iron or molybdenum at the non-acidic site, but this seems unlikely because N_2 -fixation rates in lichens at that site responded positively to P fertilization. The most likely explanation is that since lichens do not have a root system or other absorptive organs, their nutrient sources are limited to atmospheric deposition (Nash and Gries, 1995; Hyvärinen and Crittenden, 1998) and nutrients concentrated at the surface of the substrate at their immediate location. The more abundant P at the non-acidic site is concentrated in the mineral soil (Hobbie and Gough, 2002). The two lichens studied here are separated from the mineral soil by a layer of organic peat, and may be unable to access mineral soil P. Lichens on acidic tundra may not be inhibited by N because while N mineralization rates are higher at that site, they still represent quite low rates of N availability (Hobbie and Gough, 2002).

Our finding that P fertilization stimulated N_2 -fixation rates and increased lichen N concentrations at both sites and for both species is consistent with the findings of others (Chapin et al., 1991; Chapin and Bledsoe, 1992; Crittenden et al., 1994; Thompson and Vitousek, 1997; Vitousek and Field, 1999; but see Alexander et al., 1978). These results support the interpretation that N_2 fixers are limited by P in upland tundra as they are in some other ecosystems (Vitousek and Howarth, 1991).

Declines in abundance of N_2 -fixing lichens with N fertilization have been documented previously in the arctic tundra (Cornelissen et al., 2001), as well as in the boreal forest (Nohrstedt, 1998), where many of the same lichen species are found. The negative effect of N fertilization on lichens likely arises from shading of lichens or burial of lichens under more abundant leaf litter associated with greater plant biomass (Cornelissen et al., 2001; Vitousek and Field, 1999).

Contrary to our expectation, both *P. aphthosa* and *P. polydactyla* are more abundant in acidic than in non-acidic tundra. One possible explanation for these results is that favorable microsites for lichen establishment and growth are more abundant in acidic tundra. For example, we observed that these lichens commonly occur in *Sphagnum* spp. mats at the acidic site, perhaps because the moss provides a moist, cool microhabitat. *Sphagnum* spp. is not present on non-acidic tundra; microhabitats well-suited to the two lichens may be generally less abundant at the non-acidic site.

In summary, while P stimulates N_2 -fixation in *P. aphthosa* and *P. polydactyla* in experimental fertilization studies at Toolik Lake, rates of N_2 -fixation do not differ between lichens from acidic and non-acidic tundra despite differences in N and P availability at the two sites. Since the two lichens are more abundant on acidic tundra, the overall contribution of N_2 fixed by lichens is greater on acidic tundra, perhaps contributing to the overall higher N availability and productivity in acidic versus non-acidic tundra. However, further studies are necessary to determine the factors that control the abundance of *Peltigera* in tundra communities. Such studies will be critical for understanding the contribution of N_2 -fixing lichens to N budgets of arctic tundra.

Acknowledgments

We are grateful to the Toolik Lake Field Research Station and the Toolik Lake Long Term Ecological Research Program for logistic support, and to Kei Koba, Tiffany Miley, Jacques Finlay, Jim Laundre, Gus Shaver, and Lauren McSherry for field assistance and/or use of resources. We also wish to thank two anonymous reviewers who provided helpful comments on the manuscript. This research was supported by a Research Experience for Undergraduates supplement to a grant from the National Science Foundation, OPP-9902695, to S. E. Hobbie. Establishment and maintenance of the fertilization treatments was funded by the Toolik Lake LTER, DEB-9810222.




FIGURE 5. Acetylene reduction rates for *Peltigera aphthosa* and *P. polydactyla* on acidic and non-acidic tundra. Values are means of three dates with standard error bars ($n = 18$ or 17 for *P. aphthosa* on non-acidic tundra).

References Cited

- Alexander, V., 1974: A synthesis of the IBP tundra biome circumpolar study of nitrogen fixation. In: Holding, A. J., Heal, O. W., MacLean Jr., S. F., and Flanagan, P. W. (eds.), *Soil Organisms and Decomposition in Tundra*. Stockholm: Tundra Biome Steering Committee, 109–121.
- Alexander, V., Billington, M., and Schell, D. M., 1978: Nitrogen fixation in Arctic and Alpine tundra. In: Tieszen, L. L. (ed.), *Vegetation and Production Ecology of an Alaskan Arctic Tundra*. New York: Springer-Verlag, 539–558.
- Alexander, V., and Schell, D. M., 1973: Seasonal and spatial variation of nitrogen fixation in the Barrow, Alaska tundra. *Arctic and Alpine Research*, 5: 77–88.
- Arctic LTER database, 2002: <http://ecosystems.mbl.edu/arc>. Cited 18 May 2002.
- Bardsdale, R. J., and Alexander, V., 1975: The nitrogen balance of Arctic tundra: Pathways, rates, and environmental implications. *Journal of Environmental Quality*, 4: 111–117.
- Bliss, L. C., and Matveyeva, N. V., 1992: Circumpolar Arctic vegetation. In: Chapin, F. S. I., Jeffries, R. L., Reynolds, J. F., Shaver, G. R., and Svoboda, J. (eds.), *Arctic Ecosystems in a Changing Climate: An Ecophysiological Perspective*. San Diego: Academic Press, 59–89.
- Bowman, W. D., Schardt, J. C., and Schmidt, S. K., 1996: Symbiotic N_2 fixation in alpine tundra: Ecosystem input and variation in fixation rates among communities. *Oecologia*, 108: 345–350.
- Bret-Harte, M. S., Shaver, G. R., Zoerner, J. P., Johnstone, J. F., Wagner, J. L., Chavez, A. S., Gunkelman, R. F., Lippert, S. C., and Laundre, J. A., 2001: Developmental plasticity allows *Betula nana* to dominate tundra subjected to an altered environment. *Ecology*, 82: 18–32.
- Chapin, D. M., Bliss, L. C., and Bledsoe, L. J., 1991: Environmental regulation of nitrogen fixation in a high Arctic lowland ecosystem. *Canadian Journal of Botany*, 69: 2744–2755.
- Chapin, D. M., and Bledsoe, C. S., 1992: Nitrogen fixation in arctic plant communities. In: Chapin, F. S. I., Jeffries, R. L., Reynolds, J. F., Shaver, G. R., and Svoboda, J. (eds.), *Arctic Ecosystems in a Changing Climate: An Ecophysiological Perspective*. San Diego: Academic Press, 301–319.
- Cornelissen, J. H. C., Callaghan, T. V., Alatalo, J. M., Michelsen, A., Graglia, E., Hartley, A. E., Hik, D. S., Hobbie, S. E., Press, M. C., Robinson, C. H., Henry, G. H. R., Shaver, G. R., Phoenix, G. K., Jones, D. G., Jonasson, S., Chapin, F. S., Molau, U., Neill, C., Lee, J. A., Melillo, J. M., Sveinbjornsson, B., and Aerts, R., 2001: Global change and arctic ecosystems: is lichen decline a function of increases in vascular plant biomass? *Journal of Ecology*, 89: 984–994.
- Crews, T. E., 1993: Phosphorus regulation of nitrogen fixation in a traditional Mexican agroecosystem. *Biogeochemistry*, 21: 141–166.

- Crittenden, P. D., Kaluka, I., and Oliver, E., 1994: Does nitrogen supply limit the growth of lichens? *Cryptogamic Botany*, 4: 143–155.
- Englund, B., 1978: Effects of environmental factors on acetylene reduction by intact thallus and excised cephalodia of *Peltigera aphthosa* Willd. In: Granhall, U. (ed.), *Ecological Bulletins: Environmental Role of Nitrogen-fixing Blue-green Algae and Asymbiotic Bacteria*. Stockholm, 234–246.
- Gough, L., and Hobbie, S. E., 2003: Responses of moist non-acidic arctic tundra to altered environment: Productivity, biomass, and species richness. *Oikos*, 103: 204–216.
- Gunther, A. J., 1989: Nitrogen fixation by lichens in a subarctic Alaskan watershed. *Bryologist*, 92: 202–208.
- Hamilton, T. D., 2003: Glacial geology of the Toolik Lake and Upper Kuparuk River region. In: Walker, D. A. (ed.), *Biological Papers of the University of Alaska. No. 26*. Fairbanks: University of Alaska Printing Services.
- Hardy, R. W. F., Holsten, R. D., Jackson, E. K., and Burns, R. C., 1968: Acetylene-ethylene assay for N₂ fixation: Laboratory and field evaluation. *Plant Physiology*, 43: 1185–1207.
- Hendzel, L. L., Hecky, R. E., and Findlay, D. L., 1994: Recent changes of N₂ fixation in Lake 227 in response to reduction of the N/P loading ratio. *Canadian Journal Of Fisheries And Aquatic Sciences*, 51: 2247–2253.
- Hobbie, S. E., and Gough, L., 2002: Foliar and soil nutrients in tundra on glacial landscapes of contrasting ages in northern Alaska. *Oecologia*, 131: 453–462.
- Hyvärinen, M., and Crittenden, P. D., 1998: Relationships between atmospheric nitrogen inputs and the vertical nitrogen and phosphorus concentration gradients in the lichen *Cladonia portentosa*. *New Phytologist*, 140: 519–530.
- Kurina, L. M., and Vitousek, P. M., 1999: Controls over the accumulation and decline of a nitrogen-fixing lichen, *Stereocaulon vulcani*, on young Hawaiian lava flows. *Journal of Ecology*, 87: 784–799.
- Nash, T. H., and Gries, C., 1995: The response of lichens to atmospheric deposition with an emphasis on the Arctic. *Science Of The Total Environment*, 161: 737–747.
- Nash, T. H., and Olafsen, A. G., 1995: Climate change and the ecophysiological response of Arctic lichens. *The Lichenologist*, 27: 559–565.
- Nohrstedt, H. O., 1998: Residual effects of N fertilization on soil-water chemistry and ground vegetation in a Swedish Scots Pine forest. *Environmental Pollution*, 102: 77–83.
- Shaver, G. R., Chapin, F. S., and Gartner, B. L., 1986: Factors limiting seasonal growth and peak biomass accumulation in *Eriophorum vaginatum* in Alaskan tussock tundra. *Journal of Ecology*, 74: 257–278.
- Thompson, M. V., and Vitousek, P. M., 1997: Asymbiotic nitrogen fixation and litter decomposition on a long soil-age gradient in Hawaiian montane rain forest. *Biotropica*, 29: 134–144.
- Vitousek, P. M., 1999: Nutrient limitation to nitrogen fixation in young volcanic sites. *Ecosystems*, 2: 505–510.
- Vitousek, P. M., and Field, C. B., 1999: Ecosystem constraints to symbiotic nitrogen fixers: a simple model and its implications. *Biogeochemistry*, 46: 179–202.
- Vitousek, P. M., and Howarth, R. W., 1991: Nitrogen limitation on land and in the sea: How can it occur? *Biogeochemistry*, 13: 87–115.
- Walker, D. A., 1998: *The Geobotanical Atlas of the Toolik Lake and Kuparuk River Region, Alaska*. URL <http://www.geobotany.uaf.edu/arcticgeobot/index.html>. Cited 9 June 2002.
- Walker, D. A., Auerbach, N. A., Bockheim, J. G., Chapin, F. S., Eugster, W., King, J. Y., McFadden, J. P., Michaelson, G. J., Nelson, F. E., Oechel, W. C., Ping, C. L., Reeburg, W. S., Regli, S., Shiklomanov, N. I., and Vourlitis, G. L., 1998: Energy and trace-gas fluxes across a soil pH boundary in the arctic. *Nature*, 394: 469–472.
- Walker, D. A., Auerbach, N. A., and Shippert, M. M., 1995: NDVI, biomass, and landscape evolution of glaciated terrain in northern Alaska. *Polar Record*, 31: 169–178.
- Walker, M. D., Walker, D. A., and Auerbach, N. A., 1994: Plant communities of a tussock tundra landscape in the Brooks Range foothills, Alaska. *Journal Of Vegetation Science*, 5: 843–866.

Revised ms submitted April 2005