

## **Floristics and Distribution of Plant Communities across Moisture and Topographic Gradients in Tso Kar Basin, Changthang Plateau, Eastern Ladakh**

Authors: Rawat, Gopal S., and Adhikari, Bhupendra S.

Source: Arctic, Antarctic, and Alpine Research, 37(4) : 539-544

Published By: Institute of Arctic and Alpine Research (INSTAAR),  
University of Colorado

URL: [https://doi.org/10.1657/1523-0430\(2005\)037\[0539:FADOPC\]2.0.CO;2](https://doi.org/10.1657/1523-0430(2005)037[0539:FADOPC]2.0.CO;2)

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

# Floristics and Distribution of Plant Communities across Moisture and Topographic Gradients in Tso Kar Basin, Changthang Plateau, Eastern Ladakh

Gopal S. Rawat\*† and  
Bhupendra S. Adhikari\*‡

\*Department of Habitat Ecology,  
Wildlife Institute of India, P.O. Box 18,  
Chandrabani, Dehra Dun 248 001,  
Uttaranchal, India

†Corresponding author.  
rawatg@wii.gov.in  
‡adhikarib@wii.gov.in

## Abstract

This paper deals with the floristic structure and plant-community composition in relation to environmental gradients in the Tso Kar basin, an ~300 km<sup>2</sup> area of the Changthang plateau. A total of 131 sites covering various landscape units—namely, sandy plains, marsh meadows, moist meadows, scrub steppe, fell-fields, and stream courses—were intensively sampled by using random quadrats of 1 m<sup>2</sup>. In addition, these sites were searched for the presence of other vascular plants. In all, 232 species of vascular plants belonging to 38 families and 101 genera were collected and recorded from the study area. Poaceae (39 species), Asteraceae (27 species), Cyperaceae (25 species), Brassicaceae (14 species), Fabaceae (12 species), and Ranunculaceae (12 species) are the dominant families. It is interesting to note that the study area has a very high monocot:dicot ratio (1:2). Analysis of Raunkiaer's life-forms reveals that the area is dominated by two growth forms—namely, 57% hemicryptophytes (perennial grasses and sedges) and 24% chamaephytes (dwarf herbs and matted shrubs). The density of stems ranges from 28 to 744 plants per square meter. Species richness varies from 4 to 15, and diversity values are from 0.143 to 1.679. Among communities, the highest similarity (57.5%) is between *Scirpus-Carex* and *Scirpus-Kobresia-Ranunculus*. Conservation implications of the findings and the need for a participatory approach of the rangeland management are discussed.

## Introduction

The Changthang plateau in eastern Ladakh represents an important biogeographic province within Indian Trans-Himalaya (Rodgers and Panwar, 1988). This tableland forms the western extension of the Tibetan plateau and is mostly higher than 4500 m above sea level (a.s.l.). The metasedimentary rocks originated by deposition on an ancient Tethyan sea-bed, and they have undergone frequent phases of glaciation and aridity during the Pleistocene and Holocene. Now this region exhibits harsh climatic conditions, low primary productivity, and unique assemblages of flora and fauna. Much of the plateau is characterized by lake basins of varying size as well as sandy plains and rolling mountains fringed by snowy peaks. The internal drainage in some of the basins has resulted in concentration of salts and minerals over the millennia, making the water bodies brackish. Less than 1% of the geographical area on the plateau is cultivated, and most of the vegetated zone is used by migratory pastoral communities—i.e., *Changpa* herders—for livestock grazing. *Changpas* rear goats, sheep, yaks, and horses. Despite a harsh climate, poor vegetation cover, and relatively low standing biomass, this area sustains a high livestock population. The area also supports diverse but low populations of typical Trans-Himalayan mammalian fauna, such as blue sheep or bharal (*Pseudois nayaur*), Tibetan argali (*Ovis ammon hodgsoni*), Tibetan antelope (*Pantholops hodgsoni*), Tibetan gazelle (*Procarpa picticaudata*), snow leopard (*Uncia uncia*), brown bear (*Ursus arctos*), Tibetan wolf (*Canis lupus chanco*), lynx (*Lynx isabellina*), red fox (*Vulpus vulpus*), and wild dog (*Cuon alpinus*). Besides, there are several species of migratory waterfowl, including the threatened black-necked crane (*Grus nigricollis*) and bar-headed goose (*Anser indicus*).

Despite immense ecological and conservation significance, no detailed ecological studies have been conducted in these rangelands until now. A few authors previously conducted floristic surveys in the

region (e.g., Kachroo et al., 1977; Chaurasia and Singh, 1997; and Murti, 2001). Hartmann (1987), Rawat et al. (2001), Kala and Mathur (2002), Rawat and Adhikari (2002), and Klimes (2003) conducted vegetation surveys in other parts of the Indian Trans-Himalaya. Richard (1999) dealt with the rangelands and livestock as a niche opportunity for this region. Systematic surveys on various components of biodiversity in Changthang have started only recently (Anonymous, 2001).

This paper deals with the floristic structure and plant-community composition across various landscape units and environmental gradients in the Tso Kar basin of the Changthang plateau. The data were collected during a rapid ecological survey conducted in the summer of 2001. The role of biotic and abiotic factors in life-form structure and the distribution of plant communities are discussed herein.

## Materials and Methods

### STUDY AREA

The survey was conducted in the Tso Kar basin and the adjacent Rupshu plains of the Indian Changthang, eastern Ladakh (Fig. 1). This area (~300 km<sup>2</sup>) lies approximately between 33°10' and 33°30'N and between 77°55' and 78°20'E; the altitudinal range is ~4400 m to 5800 m a.s.l. It is bordered by the Zaskar range on the south, the Sumdo Nala (a tributary of the Indus) on the east, and Tanglang La (5360 m a.s.l.; a high mountain pass and the ridge extending from it) on the north. Administratively the study area lies in the Nyoma block of the Ladakh district. According to the local herders (*Changpas*), this area is divisible into three units: Samad, Kharnak, and Puga Valley. Meteorological data for the study area are not available, but it is regarded as colder and drier than other parts of Ladakh. A considerable proportion of the study area is flat with extensive sandy plains. Tso Kar, a broad pan-shaped basin, has two large water bodies adjacent to




FIGURE 1. A diagrammatic sketch of Tso Kar basin.

each other—namely, Tso Kar (~19.5 km<sup>2</sup>, 4534 m in elevation) and Tsartsapuk Tso (~2.5 km<sup>2</sup>, 4545 m in elevation). However, Tso Kar, a dumbbell-shaped brackish-water lake receives drainage from most of the basin, whereas Tsartsapuk Tso is a freshwater lake drained mainly by two perennial streams. Tsartsapuk Tso also drains into the former lake. The northwestern and eastern banks of Tso Kar are rich in deposits of borax and common salt and hence are unfavorable for plant growth except for a few salt-tolerant species such as *Axyris amaranthoides* and *Sueda microsperma*.

The vegetation of the study area can be broadly categorized into scrub formations, desert steppe, and marsh meadows. The major plant communities include *Caragana-Eurotia*, *Artemisia-Tanacetum*, *Stipa-Oxytropis-Alyssum*, and *Carex melanantha-Leymus secalinus*. The parts of the study area at very high altitudes (>5000 m) have sparse fell-field communities with moss or cushion-like growth forms, e.g., *Thylacospermum caespitosum*, *Arenaria bryophylla*, *Androsace sarmentosa*, and a variety of lichens. Stream banks and marsh meadows around both the lakes (except areas of borax and salt deposits) exhibit a characteristic sedge-dominated vegetation represented by species of *Carex*, *Kobresia*, *Scirpus*, *Triglochin*, *Pucciniella*, *Ranunculus*, and *Polygonum*. The shallow parts of Tsartsapuk support dense growths of aquatic plants such as *Hippuris vulgaris*, *Potamogeton pectinatus*, *P. perfoliatus*, *Zannichellia palustris*, and *Ranunculus nutans*.

TABLE 1

Summary of the vascular plants recorded in the Tso Kar Basin.

Group	Family	Genera	Species
Dicots	30	74	157
Monocots	7	26	74
Gymnosperms	1	1	1
Tso Kar basin (present study)	38	101	232
Ladakh (Kachroo et al. 1977)	51	190	611

The soil texture varies from sandy to sandy clay. The soil under most of the landscape units is basic in nature (pH 7.9–9.3). The soil's water-holding capacity, organic carbon, and nitrogen were observed to be highest (109%, 5.3%, and 0.623%, respectively) in boggy areas, which were mainly dominated by the *Scirpus* community (B. S. Adhikari and G. S. Rawat, unpublished data).

#### METHODS

The study area was traversed on foot as well as by vehicle (wherever feasible) to cover all the landscape features and landforms. The plant species were collected and recorded systematically for all landforms. These include marsh meadows, moist meadows, sandy plains, alluvial fans, lower (grassy) slopes, higher slopes, scrub steppe, fell-fields, and livestock camps. We used the standard literature (Polunin and Stainton, 1987; Aswal and Mehrotra, 1994; Murti, 2001) for plant identification in the field. Voucher specimens were collected for confirmation of species from the National Herbaria based at Dehra Dun. The species were categorized according to their growth habits (life-form) by using Raunkiaur's classification (Mueller-Dombois and Ellenberg, 1974).

A total of 131 sites were sampled for the structure and composition of vegetation. The study area was stratified into various landscape units. At each site, we used 10 random quadrats of 1 m<sup>2</sup> each following standard methods (Mueller-Dombois and Ellenberg, 1974; Kent and Coker, 1992). Other parameters recorded at each site were altitude, geographical coordinates (determined through the use of the Global Positioning System [GPS]), aspect, terrain, soil type, and vegetation cover. The data on the abundance (number) of individual species were recorded within each quadrat along with their approximate cover (%).

We used simple measures such as density (plants per square meter), diversity (ShannonWiener index,  $H'$ ), and richness (Menhnick's Index) as given in Magurran (1988) to describe the vegetation characteristics within various landscape units. The plant communities

TABLE 2

## Composition of vascular plants in Tso Kar basin.

Family	Genera	Species
Dicotyledons		
Amaranthaceae	1	2
Apiaceae	3	4
Asteraceae	9	27
Boraginaceae	2	2
Brassicaceae	9	14
Caprifoliaceae	1	1
Caryophyllaceae	4	13
Chenopodiaceae	5	7
Crassulaceae	1	3
Euphorbiaceae	1	1
Fabaceae	4	12
Fumariaceae	1	3
Gentianaceae	3	5
Geraniaceae	1	1
Hippuridaceae	1	1
Lamiaceae	5	10
Malvaceae	1	1
Parnassiaceae	1	1
Plantaginaceae	1	1
Polygonaceae	2	6
Potamogetonaceae	1	2
Primulaceae	2	3
Ranunculaceae	4	12
Rosaceae	2	9
Salicaceae	1	1
Saxifragaceae	1	2
Scrophulariaceae	3	8
Solanaceae	2	2
Tamaricaceae	1	1
Urticaceae	1	1
Violaceae	1	1
Monocotyledons		
Amaryllidaceae	1	3
Cyperaceae	4	25
Poaceae	17	39
Juncaceae	1	2
Juncaginaceae	1	2
Zannichelliaceae	1	1
Gymnospermae		
Ephedraceae	1	1
Total	101	232

were classified by using two-way indicator-species analysis (TWINSPAN; Hill, 1979), a computer-based polythetic divisive clustering package. The similarity (qualitative) was calculated following Magurran (1988).

## Results

### FLORISTIC STRUCTURE AND COMPOSITION

In total, 232 species of vascular plants, belonging to 38 families and 101 genera, were collected and recorded from the study area (Table 1). The dominant families include Poaceae (39 species), Asteraceae (27 species), Cyperaceae (25 species), Brassicaceae (16 species), and Fabaceae (12 species, Table 2). The ratio of monocots to dicots in the area is 1:2.1. This ratio is very high compared to the reported ratio of the Ladakh flora as a whole (1:8.30; Kachroo et al., 1977) as well as the alpine flora of the Kumaun Himalaya (1:5.2; Rawat, 1984), Deosai Plains (1:3.6; Woods et al., 1997), and the Valley of Flowers (1:4.2;

TABLE 3

Life-forms spectrum of the Tso Kar basin. Definitions of the various life-forms: Phanerophytes are trees >2 m high. Nannophanerophytes are small trees or shrubs <2 m high. Chamaephytes are woody or herbaceous perennials with buds close to the ground. Hemicryptophytes are rosette- or tussock-forming plants in which aboveground parts die back during unfavorable periods, e.g., graminoids. Geophytes are plants with subterranean bulbs, tubers, or thick rhizomes. Hydrophytes are aquatic plants. Therophytes are annual herbs that survive the unfavorable periods as seeds (for more details, see Kent and Coker, 1992).

Life-forms	Percentage of total species by region			
	Normal (Ramkaier, 1934)	Ladakh (Kachroo et al., 1977)	Changthang (Klimes, 2003)	Changthang (Present study)
Phanerophytes (Ph)	28	2.6	3.5	0
Nannophanerophytes (N)	15	4.8	0	0.9
Chamaephytes (Ch)	9	26.8	5.4	23.5
Hemicryptophytes (H)	26	9.6	62.1	56.5
Geophytes (G)	4	21.3	4.2	4.8
Hydrophytes (Hh)	2	0.6	1.7	2.2
Therophytes (T)	13	33.2	22.3	11.7

Kala et al., 1998). Such a high proportion of monocots in the area may be attributed to a long history of grazing by ungulates on the plateau and the ability of grasses and sedges to withstand heavy grazing.

As the area has a cold desert environment, certain groups of plants such as orchids, ferns, and liverworts are completely absent. A low diversity of *Astragalus*, *Pedicularis*, and *Juncus* and a high ecotypic variation among *Elymus*, *Poa*, *Artemisia*, *Oxytropis*, *Carex*, and *Kobresia* are interesting features of the flora. *Hippuris vulgaris*, *Alyssum canescens*, *Zannichellia palustris*, and *Microula tibetica* are species little known in India.

The flora is dominated by hemicryptophytic (graminoids) and chamaephytic (dwarf herbs with perennial rhizome or root stock) life-forms having ~57 and 23% species, respectively. The herbaceous dicots are generally aromatic (23%), dwarf, cushion-like (17.3%) or woolly and hairy (12%). A comparative account of life-form spectrum for the study area, Ladakh, and world is given in Table 3.

### VEGETATION CHARACTERISTICS

Four distinct physiognomic units of vegetation are discernible in the study area: (1) scrub steppe (*Caragana-Eurotia*, *Artemisia*, and *Tanacetum* formations), (2) desert steppe (*Stipa-Alyssum-Oxytropis*, *Leymus secalinus*, and *Carex melanantha* types), (3) wet and marsh meadows dominated by sedges, and (4) cushion-like vegetation or fell-fields at higher altitudes (>4800 m). The scrub steppe, desert steppe, sedge meadows, and fell-fields formed ~20, 40, 5, and 10% of the land surface, respectively. The remaining 25% of the area consists of steep rocky slopes and snow fields.

Sixteen plant communities were segregated by using TWINSPAN. The distribution of these communities with respect to major environmental gradients—namely, soil moisture, altitude, and landscape units—is shown in the Figure 2. In terms of areal coverage, the desert steppe (*Stipa-Alyssum-Oxytropis*) and scrub steppe (*Caragana-Poa*) communities are the most extensive. Communities 1–6 (*Scirpus-Carex*, *Scirpus-Carex-Ranunculus*, *Scirpus-Kobresia-Ranunculus*, *Pucciniella-Polygonum-Carex*, *Polygonum-Glaux-Carex*, and *Carex-Polygonum-Elymus*) are confined to wet meadows along the gradients of moisture in the valley bottom and stream courses. In the higher altitudes (>4800 m), especially on the wind-blown rocky slopes, *Poa-Elymus-Artemisia* and *Tanacetum tibeticum* communities are prominent. *Eurotia ceratoides* (a




FIGURE 2. Distribution pattern of plant communities along environmental gradients.

perennial shrub) and *Leymus secalinus* (a coarse grass) form pure stands of their own near hill bases. Moist areas above 5000 m are usually dominated by the *Thylacospermum-Arenaria* community.

Percent cover, plant density, species richness, diversity, and evenness for various communities are compared in Table 4. Communities 1–6 have higher density, diversity, and species richness than all other communities. Overall cover, density, species richness, and diversity follow the gradients of soil moisture and landscape units rather than altitudinal gradient. Most of the livestock camps are characterized by the prominence of an unpalatable nitrophilic herb, *Chenopodium glaucum*, and a coarse grass, *Leymus secalinus*.

The similarity between communities is observed to be highest (57.5%) for *Carex-Oxytropis-Artemisia* and *Stipa-Alyssum-Oxytropis*, followed by *Chenopodium-Leymus* and *Elymus* (57.4%, Table 5).

## Discussion

Earlier authors have interpreted Changthang plateau as floristically impoverished compared to other areas of Ladakh (e.g., Kachroo

et al., 1977; Chaurasia and Singh, 1997). An update on the flora based on the present survey reveals that the plateau harbors at least 232 species of vascular plants. This study also underlines the importance of graminoids in the flora of the region. Although forming less than 1% of the geographical area of Ladakh, the flora of the Tso Kar basin includes nearly 25% of the species reported from the region. Because the area forms a continuum with the vast Tibetan plateau, we expected very few endemic species to be found in the study area. Species sensitive to heavy trampling, grazing, and aridity (e.g., orchids and ferns) are completely absent in the present study area. The following species (which are otherwise common in other parts of the Trans-Himalaya) are either absent or extremely rare: *Ephedra Gerardiana*, *Rosa webbiana*, *Lonicera spinosa*, *Salix* spp., *Juniperus* spp., *Berberis* spp., *Hippophae tibetana*, *Cicer microphylla*, and *Cotoneaster* spp. Palynological studies in the area could reveal some interesting details of phytogeography.

The plant communities in the study area follow the gradients of soil moisture and microtopography, which determine the soil type. Of

TABLE 4  
Cover, density, species richness, diversity, and evenness of different plant communities in Tso Kar basin.

Community type	Cover (% $\pm$ 1 SD)	Density ( $m^{-2} \pm$ 1 SD)	Species richness ( $\pm$ 1 SD)	Diversity ( $H' \pm$ 1 SD)	Evenness ( $\pm$ 1 SD)
<i>Scirpus-Carex</i>	88.5 $\pm$ 8.5	468.2 $\pm$ 115.4	12.0 $\pm$ 3.2	1.301 $\pm$ 0.511	0.520 $\pm$ 0.167
<i>Scirpus-Carex-Ranunculus</i>	94.0 $\pm$ 6.2	743.7 $\pm$ 406.7	12.3 $\pm$ 3.3	1.472 $\pm$ 0.266	0.603 $\pm$ 0.157
<i>Scirpus-Kobresia-Ranunculus</i>	91.3 $\pm$ 11.1	708.2 $\pm$ 519.5	15.3 $\pm$ 3.9	1.679 $\pm$ 0.387	0.619 $\pm$ 0.113
<i>Puccinella-Polygonum-Carex</i>	83.3 $\pm$ 20.2	348.1 $\pm$ 300.0	4.0 $\pm$ 1.0	0.740 $\pm$ 0.743	0.496 $\pm$ 0.483
<i>Polygonum-Glaux-Carex</i>	85.5 $\pm$ 11.4	227.1 $\pm$ 62.3	8.3 $\pm$ 1.5	1.556 $\pm$ 0.390	0.735 $\pm$ 0.136
<i>Carex-Polygonum-Elymus</i>	90.0 $\pm$ 8.7	235.1 $\pm$ 32.9	9.7 $\pm$ 3.1	1.622 $\pm$ 0.605	0.712 $\pm$ 0.168
<i>Stipa-Alyssum-Oxytropis</i>	36.6 $\pm$ 12.5	34.7 $\pm$ 36.1	7.3 $\pm$ 2.2	1.321 $\pm$ 0.357	0.675 $\pm$ 0.124
<i>Carex-Oxytropis-Artemisia</i>	55.9 $\pm$ 22.6	99.4 $\pm$ 116.4	8.5 $\pm$ 3.7	1.232 $\pm$ 0.788	0.541 $\pm$ 0.308
<i>Poa-Elymus-Artemisia</i>	61.7 $\pm$ 23.6	27.6 $\pm$ 29.8	5.3 $\pm$ 1.5	1.006 $\pm$ 0.177	0.612 $\pm$ 0.067
<i>Leymus</i>	60.0 $\pm$ 0.0	55.7 $\pm$ 57.8	3.0 $\pm$ 0.0	0.143 $\pm$ 0.193	0.130 $\pm$ 0.176
<i>Chenopodium</i>	40.0 $\pm$ 14.1	77.8 $\pm$ 36.1	4.5 $\pm$ 0.7	0.341 $\pm$ 0.479	0.213 $\pm$ 0.298
<i>Chenopodium-Leymus</i>	46.0 $\pm$ 12.9	61.2 $\pm$ 45.5	3.8 $\pm$ 1.3	0.537 $\pm$ 0.217	0.496 $\pm$ 0.331

**TABLE 5**  
**Similarity (%) between different plant communities in Tso Kar basin.**

Community	Community											
	1	2	3	4	5	6	7	8	9	10	11	12
<i>Chenopodium</i>	100	40.0	16.7	0.0	20.3	17.2	0.0	0.0	0.0	3.8	0.0	0.0
<i>Chenopodium-Leymus</i>		100	54.5	10.5	17.2	15.2	12.5	8.0	8.7	3.8	0.0	0.0
<i>Leymus</i>			100	12.5	14.5	8.9	30.8	18.2	20.0	4.1	0.0	0.0
<i>Poa-Elymus-Artemisia</i>				100	28.6	16.5	0.0	13.3	0.0	14.0	5.3	4.3
<i>Carex-Oxytropis-Artemisia</i>					100	57.4	16.7	26.1	17.9	25.0	15.6	25.6
<i>Stipa-Alyssum-Oxytropis</i>						100	14.9	25.2	17.8	32.3	23.4	26.7
<i>Puccinella-Polygonum-Carex</i>							100	44.4	47.0	22.2	11.4	18.2
<i>Carex-Polygonum-Elymus</i>								100	47.1	38.1	31.8	37.8
<i>Polygonum-Glaux-Carex</i>									100	39.3	28.6	39.2
<i>Scripus-Carex</i>										100	50.7	57.5
<i>Scripus-Carex-Ranunculus</i>											100	42.6
<i>Scripus-Kobresia-Ranunculus</i>												100

the total species recorded from the study area, 132 species (57%) were found within the sample plots. General cover, density, species richness, and diversity are more clearly defined along the gradients of soil moisture and landscape units, instead of being controlled by altitude, as indicated by very low correlation coefficients. The vegetation cover, density, and species richness are higher in mesic sites than at drier sites. Diversity and evenness values are higher in valleys within *Polygonum*-mixed and *Carex*-mixed communities. Compared to the herbaceous alpine meadows of the Greater Himalaya (e.g., Kala et al., 1998; Singh and Rawat, 1999), the steppe formations in the study area have higher proportions of graminoids (Table 6). Compared to moist and smaller pockets in the alpine areas of the Greater Himalaya, the higher diversity and biomass of grazing ungulates—both wild and domestic—in the Trans-Himalaya suggest that the latter region has had a much longer history of grazing ungulates, which could have resulted in the prominence of graminoids (Bock et al., 1995). The vegetation cover during the peak of the growing season is highest within marsh meadows (92%) and lowest within the desert steppe (37%). The

highest stem densities were observed in marsh meadows (544 plants per square meter), followed by the densities in moist meadows (507 plants per square meter). Such a high density is largely due to dominance of tussock- and tiller-forming sedges. The densities are low in alluvial-fan (14 plants per square meter) and scrub-steppe (23 plants per square meter) areas. Species richness is lowest around livestock camps (4.8), followed by lower slopes (6.2), sandy plains (6.6), and alluvial fans (6.8). Although moist and marsh meadows have the highest values of species richness, the higher slopes, which are dominated by a variety of grasses and forbs, have equally high species diversity (Rawat and Adhikari, 2005).

An updated inventory of Changthang (Ladakh) flora highlights the preponderance of graminoids, which is attributed to the area's history of grazing by wild and domestic ungulates and to the better adaptability of graminoids to grazing. Continued biotic pressure coupled with harsh climate conditions in the area has resulted in the dominance of hemicryptophytic and chamaephytic life-forms in most of the communities (Table 6). As many as 16 plant communities have been identified in the area during this study; they are distributed along gradients of soil moisture, altitude, and landscape units.

**TABLE 6**

**Life-form spectrum in different communities in Tso Kar basin**  
**(for abbreviations, see Table 4).**

Community	Percentage of total species by life-form						
	Ph	N	Ch	H	G	Hh	T
<i>Chenopodium-Leymus</i>	—	12.5	25.0	50.0	12.5	—	—
<i>Chenopodium</i>	—	14.3	—	85.7	—	—	—
<i>Leymus</i>	—	25.0	—	75.0	—	—	—
<i>Poa-Elymus-Artemisia</i>	—	—	33.3	50.0	16.7	—	—
<i>Carex-Oxytropis-Artemisia</i>	—	2.0	33.3	49.0	5.9	—	9.8
<i>Stipa-Alyssum-Oxytropis</i>	—	1.2	31.8	51.8	4.7	—	10.6
<i>Carex-Polygonum-Elymus</i>	—	—	—	55.6	11.1	—	33.3
<i>Polygonum-Glaux-Carex</i>	—	—	11.1	72.2	—	—	16.7
<i>Puccinella-Polygonum-Carex</i>	—	—	—	62.5	12.5	—	25.0
<i>Scripus-Kobresia-Ranunculus</i>	—	—	8.9	62.2	8.9	4.4	15.6
<i>Scripus-Carex-Ranunculus</i>	—	—	7.7	69.2	11.5	3.8	7.7
<i>Scripus-Carex</i>	—	—	14.3	68.6	2.9	—	14.3

## Acknowledgments

This study was funded by the International Centre for Integrated Mountain Development (ICIMOD, Kathmandu) and Wildlife Institute of India (WII). We thank the director of WII for providing necessary facilities, Ms. Camille Richard (ICIMOD) and Dr. Y.V. Bhatnagar (WII) for their suggestions, and Mr. P.L. Saklani for his assistance in floral surveys and vegetation sampling.

## References Cited

- Anonymous, 2001: Conserving biodiversity in the Indian Trans-Himalaya: New initiatives of field conservation in Ladakh. *In First Annual Technical Report (1999–2000)*. Dehra Dun, India: Wildlife Institute of India, International Snow Leopard Trust, and U.S. Fish and Wildlife Service, 169 pp.
- Aswal, B. S., and Mehrotra, B. N., 1994: *Flora of Lahaul-Spiti (a cold desert in north west Himalaya)*. Dehra Dun, India: Bishen Singh Mahendra Pal Singh, 761 pp.
- Bock, J. H., Jolls, C. L., and Lewis, A. C., 1995: The effects of grazing on alpine vegetation: A comparison of the Central Caucasus, Republic of Georgia, with Colorado Rocky Mountains, U.S.A. *Arctic, Antarctic, and Alpine Research*, 27: 130–136.

- Chaurasia, O. P., and Singh, B., 1997: *Cold desert plants. Volume III: Changthang Valley*. Leh, India: DRDO [Defence Research and Development Organization] Field Research Laboratory, 84 pp.
- Hartmann, H., 1987: Pflanzengesellschaften trockener Standorte aus der subalpine und alpine Stufe im südlichen und östlichen Ladakh. *Candollea*, 42: 277–326.
- Hill, M. O., 1979: *TWINSPAN*—A Fortran programme for arranging multivariate data in an ordered two way table by classification of the individuals and attributes. Ithaca, New York: Cornell University.
- Kachroo, P., Sapru, B. L., and Dhar, U., 1977: *Flora of Ladakh: An ecological and taxonomic appraisal*. Dehra Dun, India: Bishen Singh Mahendra Pal Singh, 172 pp.
- Kala, C. P., and Mathur, V. B., 2002: Patterns of plant species distribution in the Trans-Himalayan region of Ladakh, India. *Journal of Vegetation Science*, 13: 751–754.
- Kala, C. P., Rawat, G. S., and Uniyal, V. K., 1998: *Ecology and conservation of the Valley of Flowers National Park, Garhwal Himalaya*. Dehra Dun, India: Wildlife Institute of India, 99 pp.
- Kent, M., and Coker, P., 1992: *Vegetation description and analysis. A practical approach*. London: Belhaven Press, 363 pp.
- Klimes, L., 2003: Life-forms and clonality of vascular plants along an altitudinal gradient in E Ladakh (NW Himalayas). *Basic and Applied Ecology*, 4: 317–328.
- Magurran, A. E., 1988: *Ecological diversity and its measurement*. Princeton: Princeton University Press, 179 pp.
- Mueller-Dombois, D., and Ellenberg, H., 1974: *Aims and methods of vegetation ecology*. New York: John Wiley and Sons, 547 pp.
- Murti, S. K., 2001: *Flora of cold deserts of western Himalaya: Volume I. Monocotyledons*. Calcutta: Botanical Survey of India, 452 pp.
- Polunin, O., and Stainton, A., 1984: *Flowers of the Himalaya*. Delhi: Oxford University Press.
- Raunkaier, C., 1934: *The life-forms of plants and statistical plant geography*. Oxford: Clarendon Press, 632 pp.
- Rawat, G. S., 1984: *Studies on the high altitude flowering plants of Kumaon Himalaya*. Ph.D. thesis, Kumaun University, Nainital, India, 453 pp.
- Rawat, G. S., and Adhikari, B. S., 2002: *Vegetation characteristics and patterns of livestock grazing in Changthang plateau, eastern Ladakh*. Report submitted to International Centre for Integrated Mountain Development, Kathmandu, Nepal, and Wildlife Institute of India, Dehra Dun, Uttaranchal, India: 26 pp.
- Rawat, G. S., and Adhikari, B. S., 2005: Millennia of grazing history in eastern Ladakh, India, reflected in rangeland vegetation. Proceedings of II<sup>nd</sup> Global Mountain Biodiversity Assessment, La Paz, Bolivia. Boca Raton, Florida: CRC Press, 14: 201–212.
- Rawat, G. S., Adhikari, B. S., and Rana, B. S., 2001: Vegetation surveys in the Indian Trans-Himalaya. In *Conserving Biodiversity in the Indian Trans-Himalaya: New Initiatives of Field Conservation in Ladakh*. Dehra Dun, India: Wildlife Institute of India, International Snow Leopard Trust, and U.S. Fish and Wildlife Service, 7–14.
- Richard, C., 1999: *Rangelands and livestock as a niche opportunity for Ladakh*. Kathmandu, Nepal: International Centre for Integrated Mountain Development.
- Rodgers, W. A., and Panwar, H. S., 1988: *Planning a protected area network for India: Volumes I and II*. Dehra Dun, India: Wildlife Institute of India, 341 and 267 pp.
- Singh, S. K., and Rawat, G. S., 1999: *Flora of Great Himalayan National Park, Himachal Pradesh*. Dehra Dun, India: Bishen Singh Mahendra Pal Singh, 304 pp.
- Woods, C. A., Kilpatrick, C. W., Rafiq, M., Shah, M., and Khan, W., 1997: Biodiversity and conservation of the Deosai Plateau, northern areas, Pakistan. In Mufti, S. A., Woods, C. A. and Hasan, S. A. (eds.), *Biodiversity of Pakistan*. Islamabad, Pakistan: Pakistan Museum of Natural History, 33–61.

*Ms submitted November 2003*