

## **Epigeal Spider (Araneae) Communities in Moist Acidic and Dry Heath Tundra at Toolik Lake, Alaska**

Authors: Wyant, Karl A., Draney, Michael L., and Moore, John C.

Source: Arctic, Antarctic, and Alpine Research, 43(2) : 301-312

Published By: Institute of Arctic and Alpine Research (INSTAAR),  
University of Colorado

URL: <https://doi.org/10.1657/1938-4246-43.2.301>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

# Epigeal Spider (Araneae) Communities in Moist Acidic and Dry Heath Tundra at Toolik Lake, Alaska

Karl A. Wyant\*†

Michael L. Draney‡ and

John C. Moore†§

\*Corresponding author. Present address: Arizona State University, School of Life Sciences, Tempe, Arizona 85287, U.S.A.  
kawyant@asu.edu

†Natural Resource Ecology Laboratory, Colorado State University, Fort Collins, Colorado 80523, U.S.A.

‡Cofrin Center for Biodiversity and Department of Natural & Applied Sciences, University of Wisconsin–Green Bay, Green Bay, Wisconsin 54311-7001, U.S.A.  
draneym@uwgb.edu  
§jcmoore@nrel.colostate.edu

## Abstract

For the 2004–2006 growing seasons, we trapped a total of 6980 spiders (5066 adults, 1914 immatures) using pitfall traps at the Arctic Long Term Experimental Research (LTER) site in Toolik Lake, Alaska. We found 10 families and 51 putative species, with 45 completely identified, in two distinct habitats: Moist Acidic Tundra (MAT) and Dry Heath (DH) Tundra. We captured spiders belonging to the following families (number of species captured): Araneidae (1), Clubionidae (1), Dictynidae (1), Gnaphosidae (4), Linyphiidae (26), Lycosidae (11), Philodromidae (2), Salticidae (1), Theridiidae (1), and Thomisidae (3). Statistical comparisons of families captured at MAT and DH Tundra indicate that the habitats have significantly different spider communities (Chi Square Test:  $p < 0.0001$ , and Fisher's Exact Test:  $p = 0.0018$ ). This finding is further supported by differences in similarity, diversity, evenness, and species richness between the two habitats. In this report, we present eight new state records and five extensions of previously described ranges for spider species. The following species are new state records for Alaska: *Emblyna borealis* (O.P.-Cambridge 1877), *Horcotes strandi* (Sytschevskaja 1935), *Mecynargus monticola* (Holm 1943), *Mecynargus tungusicus* (Eskov 1981), *Metopobactrus prominulus* (O.P.-Cambridge 1872), *Poecilonea theridiformis* Emerton 1911, and *Poecilonea vakkhanka* (Tanasevitch 1989). The following five species have been reported previously in Alaska, but not near Toolik Lake: *Hypsosinga groenlandica* Simon 1889, *Gnaphosa borea* Kulczyn'ski 1908, *Gnaphosa microps* Holm 1939, *Haplodrassus hiemalis* (Emerton 1909), and *Islandiana cristata* Eskov 1987. Pairwise similarity indices were calculated across 13 other arctic and subarctic spider communities and statistical tests show that all sites are dissimilar ( $p = 0.25$ ). These results fit the general pattern of both the patchiness and habitat specificity of arctic spider fauna.

DOI: 10.1657/1938-4246-43.2.301

## Introduction

Taxonomic knowledge and descriptive reports of northern arctic spider faunas have lagged behind those made in temperate latitudes (Dondale et al., 1997). Within the Arctic, reports of spider assemblages from the northern Nearctic regions (e.g., northern Alaska and Canada) are outnumbered by reports from the Palearctic and southern Nearctic regions (e.g., Cotton, 1979; Cutler and Saltmarch, 1986; Koponen, 1992; Dondale et al., 1997; Pickavance, 2006). Of the assemblages from the Nearctic, the arctic spider fauna is best described from Greenland, where historical connections to European scientists, naturalists, and universities have resulted in a well-documented spider fauna (Danks, 1981; Pickavance, 2006). Despite the existence of Alaska spider species lists and small collections of specimens (see Jackson, 1933; Davis, 1936; Chamberlin and Ivie, 1947; Holm, 1960, 1970; Cutler and Saltmarch, 1986; Slowik, 2006; Paquin et al., 2010), the spider fauna of northern Alaska is still largely unknown except near Barrow (Weber, 1949, 1950). Furthermore, reports focused on large collections of arctic Nearctic spiders are scarce (Pickavance, 2006).

The material reported here represents the culmination of a three year, intensive sampling effort at Toolik Lake in the North Slope Borough, which encompasses 230,510 km<sup>2</sup> of territory across northern Alaska. Toolik Lake is located north of the

Brooks Range at the Arctic Long Term Ecological Research (LTER) site. To the authors' knowledge, this study is the first of its kind for this area. Large collections, made over multiple habitat areas in the Arctic, are valuable because they can record changes in the range of species, report new taxa, compile biogeographical data of previously known taxa, and contribute to our overall understanding of arctic biodiversity (Pickavance, 2006).

In this paper, we report on spider species, captured via pitfall trap, from two distinct tundra habitat types, Moist Acidic Tundra (MAT) and Dry Heath (DH) Tundra in northern Alaska. We compare the community structure of ground spiders of the two tundra types in terms of species richness, evenness, and diversity in relation to soil and vegetation features. We then extend our analysis of epigeal spider community comparisons to include published data sets for communities from other arctic and subarctic sites from within the Nearctic zone.

## Methods

### SITE DESCRIPTION

We conducted surveys in the summers of 2004–2006 at the Arctic Long Term Ecological Research (LTER) site at Toolik Lake, Alaska (Fig. 1). Toolik Lake is located in the northern




FIGURE 1. The Moist Acidic Tundra (MAT) and Dry Heath (DH) Tundra study sites are located at Toolik Lake LTER, Alaska, which is located on the north slope of the Brooks Range along the Dalton Highway.

foothills of the Brooks Range ( $68^{\circ}38'N$  and  $149^{\circ}43'W$ , elevation 760 m) approximately 240 km north of the Arctic Circle. The site is maintained by the Toolik Field Station operated by the Institute of Arctic Biology of the University of Alaska–Fairbanks and was formally established as a site within the LTER network in 1987. The mean annual air temperature at Toolik Lake is  $-7^{\circ}C$ . The growing season is approximately 50–70 days. The mean temperature during the growing season is  $10^{\circ}C$  (Hobbie and Chapin, 1998). Annual precipitation ranges from 200 to 400 mm with 50% falling as snow. Distribution of snow cover is highly dependent on wind and topographical relief.

The Toolik Lake LTER site is typical of the northern foothills region near the Brooks Range. This area has continuous permafrost, no large trees, complete snow cover and ice for most of the year, and cessation of river flow during winter. Waterlogged soils are commonplace as the permafrost prevents the downward drainage of water. At the landscape scale, the tundra vegetation is composed of sedges, grasses, dwarf birches, low willows, mosses, and lichens. The presence and combinations of these plant types is dependent on age since glaciation, soil type, soil moisture, exposure, drainage, and topography (Doles, 2000). Abiotic properties, coupled with the vegetative community, give rise to four unique ecosystem types surrounding Toolik Lake. These include moist acidic, dry heath, wet sedge, and shrub tundra. Ecosystem types discussed here are Moist Acidic Tundra (MAT) and Dry Heath (DH) Tundra.

MAT is the most common vegetation type surrounding Toolik. This vegetation type constitutes 38.7% of the total land cover near Toolik Lake (Walker, 1998). The MAT habitats generally have moist soils and are covered by an organic layer 0–30 cm thick (Shaver and Chapin, 1991). The MAT vegetation type occurs on rolling topography with silt to gravelly soils with varying glaciation histories. Maximum thaw depth is 30–50 cm and, in many locations, the soil does not thaw to the mineral layer (Doles, 2000). The plant community is dominated by an equal biomass of graminoids (*Eriophorum vaginatum* and *Carex bigelowii*), shrubs

(*Betula nana*), evergreens, and mosses (Chapin et al., 1995). Of particular note, the grass species *E. vaginatum* forms dense, compact tufts called tussocks, which are typical of moist acidic tundra. Average plant species density is 11 species/ $m^2$  (Gough et al., 2007). This habitat type is named for the slightly acidic soil and high soil water content. Net primary productivity (NPP) is estimated at  $145\text{ g m}^{-2}\text{ y}^{-1}$  (Gough et al., 2007).

DH Tundra is an arctic vegetation type comprising 7.8% of the land cover near Toolik Lake (Walker, 1998). DH Tundra is characterized by drier conditions, a predominance of dwarf shrubs (*B. nana*), fruticose-lichen plant biomass, and graminoids (*Hierochloa alpine* and *Carex microchaeta*, (Shaver and Chapin, 1991; Gough et al., 2007). Average plant species density is 6 species/ $m^2$  (Gough et al., 2007). This type of tundra occurs on exposed, rocky interfluve areas with thin organic layers (0–10 cm) that retain little snow cover during winter because of wind exposure (Nadelhoffer et al., 1991; Cheng et al., 1998). These areas are well drained and have a deeply thawed mineral layer ( $>2\text{ m}$ ). NPP is estimated at  $60\text{ g m}^{-2}\text{ y}^{-1}$  (Gough et al., 2007). The two sites used in this study are roughly 3.2 km apart.

#### PITFALL TRAPPING

We estimated the diversity and total abundance of surface active invertebrate populations over the growing season (June–August) with pitfall traps in both the MAT and DH Tundra. The traps were 10 cm deep with an 8 cm opening diameter. Traps were one-third filled with 70% ethanol and collected and emptied after 3 days following each date listed in Table 1. We chose pitfall traps for epigeal spider sampling because of the need for passive, minimally destructive sampling techniques on long term ecological research plots.

At the MAT and DH Tundra sites, plot size measured  $5 \times 5\text{ m}$ , with 4 plots per block. The MAT site had four blocks while the design in the DH site had three due to space constraints. This resulted in 16 total plots at the MAT and 12 plots at the DH

TABLE 1

**Sample dates for pitfall traps for this study. All dates include Moist Acidic Tundra (MAT) and Dry Heath (DH) Tundra sites, unless noted. Traps were set and left open three days following the collection date listed below.**

| Sample Dates   |
|--|
| <b>Year: 2004</b>  |
| 14 June, 25 June, 03 July, 10 July, 31 July, 19 August                                     |
| <b>Year: 2005</b>  |
| 06 June, 13 June, 20 June, 27 June, 11 July, 18 July, 04 August, 15 August, 18 August      |
| <b>Year: 2006</b>  |
| 11 June (DH only), 14 June, 17 June, 20 June (DH only), 15 July, 25 July, 28 July, 31 July |

Tundra. We placed the traps roughly 1 m from each other on the east (MAT) or south (DH) sides of the plots. In 2004, we placed three traps within each plot. We increased the number of traps to four per plot in 2005 and 2006 and repeated for each block. This resulted in 84 (2004) and 112 (2005 and 2006) total samples per date. We sampled 6 dates in 2004 and 2006, and 9 dates in 2005 during the arctic growing season (May–August). Furthermore, for 2006, we sampled an additional two times in DH before snowmelt allowed sampling at the MAT site. This resulted in 2280 total possible samples (1248 total MAT samples; 1032 total DH samples). However, we lost a few traps and samples due to wind, precipitation events, animal disturbance, and shipping error. We lost 66 (2.89% loss of the total) samples between 2004 and 2006 (34 missing samples in the MAT and 32 samples in the DH). This represents a loss of 2.72% of possible total MAT samples and 3.1% loss of total DH Tundra samples. Thus, the data analyzed here are from 2214 total samples (1214 samples in the MAT and 1000 samples in the DH Tundra) representing 6642 total trap days of effort. At each site, there were 3642 total trap days of effort in the MAT site and 3000 trap days of effort in the DH Tundra.

Contents were preserved in 70% ETOH and identified to species when possible using appropriate keys (Dondale and Redner, 1978, 1990; Platnick and Dondale, 1992; Buckle et al., 2001, Dondale et al., 2003, Ubick et al., 2005). We followed Platnick (2010) for taxonomy, except for Linyphiidae, where Buckle et al. (2001) was preferred. Maturity of specimens was determined by the presence of sclerotized genitalia. The first author identified most of the material; second author identified most of the individuals belonging to the family Linyphiidae. Mr. Don Buckle also helped in the identification and confirmation of linyphiid species. We identified the early instars to family level or genus when possible and these data are not presented here, as this information is not appropriate for our species list and richness calculations. Range information was derived from Dondale and Redner (1978, 1990), Platnick and Dondale (1992), Dondale et al., (2003), Ubick et al. (2005), Buckle et al. (2001), Paquin et al. (2010), and Platnick (2010). Vouchers are in the collections of the first and second author. Furthermore, vouchers will be deposited at Field Museum of Natural History, Chicago, Illinois, U.S.A. (FMNH), and the University of Alaska Museum Insect Collection (UAM) by the second author. The pitfall trap by-catch (non spiders) was stored in vials of ethanol for future work.

#### ESTIMATION OF SPECIES DIVERSITY AND COMMUNITY SIMILARITY

Total abundance of spiders and community indices were calculated from pooled data for each site over all sampling dates

for both Moist Acidic Tundra (MAT) and Dry Heath (DH) Tundra. Total abundances were calculated for the years 2004–2006 for mature specimens only. We calculated Shannon's Diversity Index, evenness, richness, and Sørensen's Similarity Index for the years 2004–2006 using data for all of the species found in this study.

The indices used were all based on standard forms. Shannon's Diversity Index ( $H$ ) (Shannon, 1948) was calculated using the following equation:

$$H = - \sum_{i=1}^S p_i * (\ln * p_i) \quad (1)$$

with  $S$  = number of species present in sample (Species Richness) and  $p_i$  = the proportion of individuals made up of the  $i$ th species.

Species Evenness ( $E$ ) was calculated using the following (Brower et al., 1998):

$$E = H/H_{max} \quad (2)$$

with  $H$  = Shannon's Diversity Index and  $H_{max}$  = theoretical  $H$  if all the species in the community had an equal number of individuals.  $H_{max}$  is found by taking the natural log of the species richness.

Sørensen's similarity index ( $C_s$ ) (Magurran, 2004) was used to measure the similarity in species composition for two sites, A and B, given by the following equation:

$$C_s = 2 * c / (a + b) \quad (3)$$

where  $a$  is the number of species found in site A,  $b$  is the number of species in site B, and  $c$  is the number of species shared in common by sites A and B. Several arctic and subarctic sites were chosen based on their analyses being restricted to epigeal spider fauna and similarity of trapping method.

Species rarefaction curves were calculated using PAST software (Hammer and Harper, 2005).

#### STATISTICS

Total overall family abundance differences between MAT and DH were tested using the Chi-Square test for families with >5 total spiders (significant difference when  $\alpha \leq 0.05$ ) (SAS Institute, 2000). For families with <5 total spiders, we used Fisher's Exact Test (significant difference when  $\alpha \leq 0.05$ ). Significant differences between individual families in the MAT and DH were tested using a binomial proportion  $T$ -test with a Bonferroni correction ( $p \leq 0.05$ ) (R Development Core Team, 2005). Pairwise site similarity was analyzed using the ANOVA analysis package in Microsoft Excel (version 2007).

## Results

#### SPECIES LIST AND TOTAL ABUNDANCE

We trapped a total of 6980 spiders (5066 adults, 1914 immatures and juveniles) from moist acidic tussock tundra and dry heath tundra from June to August during 2004–2006. Families and species identified are shown in Table 2. We captured 3404 adult spiders in the MAT (67.2% of the total) and 1662 adult spiders (32.8% of the total) in the DH. Analyses of the species list using Chi Square and Fisher's Exact tests indicates that, overall, there are significant differences between the spider families found in the MAT and DH Tundra (Chi Square  $p < 0.001$ ; Fisher's Exact test  $p = 0.0018$ ). The relative abundance (proportions) of



TABLE 2

List of spider species found in Moist Acidic Tundra (MAT) and Dry Heath (DH) Tundra at Toolik Lake LTER, Alaska, from 2004 to 2006. We captured 51 morphospecies, of which 45 were identified to the species level. Note that an immature spider from the family Tetragnathidae and one from the superfamily Amaurobioidea are not included in this table. Furthermore, immature and juvenile spiders are not included in this table. Bold numbers indicate a cumulative total at the family level and across both sites.

| Family/Species   | # Collected per Site | # Collected per Site |                 |
|--|----------------------|----------------------|-----------------|
| Site   | Moist Acidic Tundra  | Dry Heath Tundra     | Total Collected |
| <b>Araneidae</b>   | <b>11</b>            | <b>0</b>             | <b>11</b>       |
| <i>Hypsosinga groenlandica</i> Simon 1889                | 11                   | 0                    | 11              |
| <b>Clubionidae</b>                                       | <b>11</b>            | <b>0</b>             | <b>11</b>       |
| <i>Clubiona praematura</i> Emerton 1909                  | 11                   | 0                    | 11              |
| <b>Dictynidae</b>  | <b>0</b>             | <b>1</b>             | <b>1</b>        |
| <i>Emblyna borealis</i> (O.P.-Cambridge 1877)            | 0                    | 1                    | 1               |
| <b>Gnaphosidae</b>                                       | <b>97</b>            | <b>32</b>            | <b>129</b>      |
| <i>Gnaphosa borea</i> Kulczyn'ski 1908                   | 15                   | 11                   | 26              |
| <i>Gnaphosa microps</i> Holm 1939                        | 46                   | 7                    | 53              |
| <i>Gnaphosa orites</i> Chamberlin 1922                   | 5                    | 11                   | 16              |
| <i>Haplodrassus hiemalis</i> (Emerton 1909)              | 31                   | 3                    | 34              |
| <b>Linyphiidae</b>                                       | <b>246</b>           | <b>229</b>           | <b>475</b>      |
| <i>Agyneta pseudosaxatilis</i> Tanasevitch 1984          | 43                   | 0                    | 42              |
| <i>Agyneta simplex</i> (Emerton 1926)                    | 1                    | 7                    | 8               |
| <i>Bathypantes simillimus</i> (L. Koch 1879)             | 18                   | 0                    | 18              |
| <i>Gonatium crassipalpus</i> Bryant 1933                 | 18                   | 1                    | 19              |
| <i>Hilaira herniosa</i> (Thorell 1875)                   | 72                   | 0                    | 72              |
| <i>Hilaira leviceps</i> (L. Koch 1879)                   | 12                   | 0                    | 12              |
| <i>Horcotes strandi</i> (Sytschevskaja 1935)             | 13                   | 1                    | 14              |
| <i>Hybauchenidium aquilonare</i> (L. Koch 1879)          | 1                    | 0                    | 1               |
| <i>Islandiana cristata</i> Eskov 1987                    | 8                    | 0                    | 8               |
| <i>Mecynargus monticola</i> (Holm 1943)                  | 11                   | 2                    | 13              |
| <i>Mecynargus tungusicus</i> (Eskov 1981)                | 22                   | 0                    | 22              |
| <i>Metopobactrus prominulus</i> (O.P.-Cambridge 1872)    | 0                    | 210                  | 210             |
| <i>Microlinyphia m. mandibulata</i> (Emerton 1882)       | 2                    | 0                    | 2               |
| <i>Microlinyphia pusilla</i> (Sundevall 1829)            | 1                    | 0                    | 1               |
| <i>Poecilometes theridiformis</i> (Emerton 1911)         | 2                    | 0                    | 2               |
| <i>Poecilometes vakkhanka</i> (Tanasevitch 1989)         | 1                    | 0                    | 1               |
| <i>Poecilometes variegata</i> (Blackwall 1841)           | 1                    | 0                    | 1               |
| <i>Scotinotylus alienus</i> (Kulczyn'ski 1885)           | 1                    | 0                    | 1               |
| <i>Walckenaeria clavicornis</i> (Emerton 1882)           | 9                    | 6                    | 15              |
| <i>Walckenaeria fraudatrix</i> Millidge 1983             | 6                    | 0                    | 6               |
| <i>Walckenaeria karpinskii</i> (O.P.-Cambridge 1873)     | 1                    | 0                    | 1               |
| <i>Walckenaeria</i> sp. cf. <i>arctica</i> Millidge 1983 | 1                    | 0                    | 1               |
| <i>Linyphiidae</i> sp. 4                                 | 0                    | 1                    | 1               |
| <i>Linyphiidae</i> sp. 6                                 | 0                    | 1                    | 1               |
| <i>Linyphiidae</i> sp. 7                                 | 1                    | 0                    | 1               |
| <i>Linyphiidae</i> sp. 8                                 | 1                    | 0                    | 1               |
| <b>Lycosidae</b>   | <b>2859</b>          | <b>1272</b>          | <b>4131</b>     |
| <i>Alopecosa aculeata</i> (Clerck 1758)                  | 2                    | 0                    | 2               |
| <i>Alopecosa hirtipes</i> (Kulczyn'ski 1907)             | 2                    | 5                    | 7               |
| <i>Alopecosa pictilis</i> (Emerton 1885)                 | 17                   | 215                  | 232             |
| <i>Arctosa alpigena</i> Doleschall 1852                  | 125                  | 63                   | 188             |
| <i>Arctosa insignata</i> (Thorell 1872)                  | 555                  | 307                  | 862             |
| <i>Pardosa concinna</i> (Thorell 1877)                   | 118                  | 34                   | 152             |
| <i>Pardosa furcifera</i> (Thorell 1875)                  | 0                    | 2                    | 2               |
| <i>Pardosa lapponica</i> (Thorell 1872)                  | 1289                 | 282                  | 1571            |
| <i>Pardosa moesta</i> Banks 1892                         | 1                    | 0                    | 1               |
| <i>Pardosa podhorskii</i> (Kulczyn'ski 1907)             | 379                  | 363                  | 742             |
| <i>Pardosa sodalis</i> Holm 1970                         | 371                  | 1                    | 372             |
| <b>Philodromidae</b>                                     | <b>41</b>            | <b>22</b>            | <b>63</b>       |
| <i>Apollophanes margareta</i> Lowrie & Gertsch 1955      | 1                    | 0                    | 1               |
| <i>Thanatus arcticus</i> Thorell 1872                    | 40                   | 22                   | 62              |
| <b>Salticidae</b>  | <b>0</b>             | <b>1</b>             | <b>1</b>        |
| <i>Chalcoscirtus glacialis</i> Caporiacco 1935           | 0                    | 1                    | 1               |

TABLE 2

Continued.

| Family/Species                           | # Collected per Site | # Collected per Site |                 |
|--|----------------------|----------------------|-----------------|
| Site                                     | Moist Acidic Tundra  | Dry Heath Tundra     | Total Collected |
| <b>Theridiidae</b>                       | <b>10</b>            | <b>0</b>             | <b>10</b>       |
| <i>Robertus</i> sp. 1                    | 10                   | 0                    | 10              |
| <b>Thomisidae</b>                        | <b>129</b>           | <b>105</b>           | <b>234</b>      |
| <i>Ozyptila arctica</i> Kulczyn'ski 1908 | 38                   | 12                   | 50              |
| <i>Xysticus britcheri</i> Gertsch 1934   | 86                   | 3                    | 89              |
| <i>Xysticus deichmanni</i> Sørensen 1898 | 5                    | 89                   | 94              |
| <b>Total Spiders</b>                     | <b>3404</b>          | <b>1662</b>          | <b>5066</b>     |

Araneidae, Clubionidae, Lycosidae, Linyphiidae, Theridiidae, and Thomisidae within the spider communities differed significantly between the MAT and DH Tundra (Table 3). Araneidae, Clubionidae, Lycosidae, and Theridiidae are significantly more abundant in MAT, whereas Linyphiidae and Thomisidae are significantly more abundant in DH Tundra. The proportions of the families Dictynidae, Gnaphosidae, Philodromidae, and Salticidae did not differ significantly between the MAT and DH Tundra.

#### RANGE EXTENSIONS AND STATE RECORDS

We present eight new state records and five extensions of previously reported species ranges (not implying any movement due to climate change), representing 25.5% of the total species captured. *Agyneta pseudosaxatilis* Tanasevitch 1984 has only been described in the Old World in Kazakhstan (Marusik et al., 2000) and is represented here, for the first time, as occurring in the Nearctic (Table 4). The following species are new state records for Alaska: *Emblyna borealis* (O.P.-Cambridge 1877), *Horcotes strandi* (Sytschevskaja 1935), *Mecynargus monticola* (Holm 1943), *Mecynargus tungusicus* (Eskov 1981), *Metopobactrus prominulus* (O.P.-Cambridge 1872), *Poecilometes theridiformis* Emerton 1911, and *Poecilometes vakkhanka* (Tanasevitch 1989). Interestingly, while the previously listed spider species have not been found in Alaska, they have been found across Canada, Greenland, and into the Palearctic zone (e.g., Finland, Siberia, etc.). Thus, these records represent westward extensions of known occurrences. The following five species have been reported previously in Alaska, but not in the North Slope, and so are northern extensions of known occurrences: *Hypsosinga groenlandica* Simon 1889, *Gna-*

*phosa borea* Kulczyn'ski 1908, *Gnaphosa microps* Holm 1939, *Haplodrassus hiemalis* (Emerton 1909), and *Islandiana cristata* Eskov, 1987. Of the records described here, spiders belonging to the family Linyphiidae made up 61.5% of the total, followed by Gnaphosidae (23.1%), Dictynidae (7.7%), and Araneidae (7.7%).

#### FAMILY AND SPECIES RICHNESS, EVENNESS, AND DIVERSITY

We captured 51 putative species, of which 45 were identified to the species level. Six species, belonging to Linyphiidae and Theridiidae, did not match published descriptions and may represent species that are undescribed. We feel that these unconfirmed species are important to include in our richness, evenness, diversity, and similarity calculations because it reflects the actual richness of our samples more accurately than if we only included taxonomically well-known species. These undescribed species should be noted to show the extent of how poorly known spider diversity is in the Toolik Lake region.

We captured 10 spider families in our traps within the MAT and DH Tundra sites, and 12 spider families if we include the immature Tetragnathidae and Amaurobioidea specimens that were captured and identified as immature by a lack of sclerotized genitalia. When the study sites were considered separately, the MAT site yielded 8 spider families and the DH Tundra site yielded 7 (Table 5). When the study sites were considered separately, the MAT had 45 spider species and the DH had 27. The most speciose family, the linyphiids, constituted ~51% of the 51 species recorded (Table 6). Families Lycosidae, Gnaphosidae, Thomisidae, and Philodromidae composed 21.6%, 7.8%, 5.9%, and 3.9% of the species recorded, respectively. The remaining families (Araneidae,

TABLE 3

Total abundances as a percentage of total ground spiders collected per family from 2004 to 2006 in Moist Acidic Tundra (MAT) and Dry Heath (DH) Tundra at Toolik Lake, Alaska, for Whole Site (MAT + DH), MAT only, DH only.

|               | MAT + DH (%) | With MAT Total Only (%) | Within DH Total Only (%) | Point Estimate for Proportion Difference | Confidence Interval Lower and Upper Range |        |
|---------------|--------------|-------------------------|--------------------------|--|---|--------|
| Araneidae *   | 0.22         | 0.32                    | —                        | −0.344                                   | −0.579                                    | −0.108 |
| Clubionidae * | 0.22         | 0.32                    | —                        | −0.344                                   | −0.579                                    | −0.108 |
| Dictynidae    | 0.02         | —                       | 0.06                     | —  | —   | —      |
| Gnaphosidae   | 2.55         | 2.85                    | 1.93                     | —  | —   | —      |
| Linyphiidae * | 9.38         | 7.23                    | 13.78                    | 0.065                                    | 0.038                                     | 0.092  |
| Lycosidae *   | 81.54        | 83.99                   | 76.53                    | −0.082                                   | −0.116                                    | −0.048 |
| Philodromidae | 1.24         | 1.20                    | 1.32                     | —  | —   | —      |
| Salticidae    | 0.02         | —                       | 0.06                     | —  | —   | —      |
| Theridiidae * | 0.20         | 0.29                    | —                        | −0.313                                   | −0.542                                    | −0.082 |
| Thomisidae *  | 4.62         | 3.79                    | 6.32                     | 0.025                                    | 0.006                                     | 0.044  |

\* Indicates a statistically significant difference in familial proportion between the MAT and DH for a binomial proportion *T*-test. A dash indicates non-significance.

TABLE 4

Reported range extensions and Alaska state records for species collected at Toolik Lake, Alaska. Abbreviations used to indicate range change are NE (new to the Nearctic Region), AK (new to the state of Alaska but it has been found elsewhere in the Nearctic), and N (species has been found in southern portions of Alaska).

| Family/Species  | Range Change | Reported Range and References   |
|---|--------------|---|
| <b>Araneidae</b>                                      |              |   |
| <i>Hypsosinga groenlandica</i> Simon 1889             | N            | Yukon Territory and western Northwest Territories, southern British Columbia and Quebec, Greenland (Dondale et al., 2003); southern Alaska (Slowik, 2006); U.S.A., Canada, Greenland (Platnick, 2010)               |
| <b>Dictynidae</b>                                     |              |   |
| <i>Emblyna borealis</i> (O.P.-Cambridge 1877)         | AK           | Nunavut (Jackson, 1933; Davis, 1936); Greenland (Leech, 1966); Russia (Platnick, 2010)  |
| <b>Gnaphosidae</b>                                    |              |   |
| <i>Gnaphosa borea</i> Kulczynski 1908                 | N            | Southern Alaska to Labrador, south to New Mexico and to northern New England; Siberia (Platnick and Dondale, 1992); Yukon Territory (Dondale et al., 1997); western Alaska (Holm, 1970); Holarctic (Platnick, 2010) |
| <i>Gnaphosa microps</i> Holm 1939                     | N            | Southern Alaska to Newfoundland, south to Colorado and northern New England; Europe; U.S.S.R. (Platnick and Dondale, 1992; Holm, 1970); Yukon Territory (Dondale et al., 1997); Holarctic (Platnick, 2010)          |
| <i>Haplodrassus hiemalis</i> (Emerton 1909)           | N            | Middle to southern Alaska to Newfoundland, south to Colorado and New Jersey; U.S.S.R. (Platnick and Dondale, 1992); Yukon Territory (Dondale et al., 1997); Holarctic (Platnick, 2010)                              |
| <b>Linyphiidae</b>                                    |              |   |
| <i>Agyneta pseudosaxatilis</i> Tanasevitch 1984       | NE           | Northwest Kazakhstan (Marusik et al., 2000)   |
| <i>Horcotes strandi</i> (Sytshevskaja 1935)           | AK           | Yukon Territory, Palearctic (Buckle et al., 2001); Finland, Russia, Canada (Platnick, 2010)   |
| <i>Islandiana cristata</i> Eskov 1987                 | N            | Yukon Territory, Siberia. (Buckle et al., 2001); Russia, Alaska, Canada (Platnick, 2010)  |
| <i>Mecynargus monticola</i> (Holm 1943)               | AK           | British Columbia, Northwest Territory, Yukon Territory, Palearctic (Buckle et al., 2001); Sweden, Finland, Russia, Mongolia, Canada (Platnick, 2010)  |
| <i>Mecynargus tungusicus</i> (Eskov 1981)             | AK           | Yukon Territory, eastern Palearctic (Buckle et al., 2001); Russia, Kyrgyzstan, China, Canada (Platnick, 2010)   |
| <i>Metopobactrus prominulus</i> (O.P.-Cambridge 1872) | AK           | Maine, Massachusetts, Michigan, New Hampshire, Alberta, New Brunswick, Nova Scotia, Northwest Territories, Ontario, Quebec, Saskatchewan, Greenland, Palearctic (Buckle et al., 1998); Holarctic (Platnick, 2010)   |
| <i>Poecilometes theridiformis</i> (Emerton 1911)      | AK           | New Hampshire, New Brunswick, Quebec, Saskatchewan, eastern Palearctic (Buckle et al., 2001); Russia, North America (Platnick, 2010)  |
| <i>Poecilometes vakhanka</i> (Tanasevitch 1989)       | AK           | Northwest Territories, eastern Siberia (Buckle et al., 2001); Russia (Platnick, 2010)   |

Clubionidae, Dictynidae, Salticidae, and Theridiidae) were rare and each contributed one species to the list.

Estimates of family level and species evenness ( $E$ ) for the MAT and DH Tundra are presented in Table 5. Overall, the study area (the combined MAT and DH sites) had a family level evenness of  $E = 0.313$ . At the species level, the pooled data had an evenness calculation of  $E = 0.598$ . When the sites are considered separately, the family evenness for the MAT and DH was  $E = 0.322$  and  $E = 0.408$ , respectively. The species level evenness for the MAT and DH was  $E = 0.572$  and  $E = 0.651$ , respectively. Shannon's Diversity Index ( $H$ ) was calculated for spiders captured in MAT and DH (Table 5). Overall, (MAT + DH), the familial diversity for the whole study area was estimated at  $H = 0.724$ . The diversity of epigeal spider community at the species level, for both sites combined, measured  $H = 2.352$ . When the MAT and DH spider communities are considered separately, the familial diversity for both sites study sites are  $H = 0.669$  and  $H = 0.794$  for MAT and DH, respectively (Table 5). The diversity index is  $H = 2.17$  for the MAT and  $H = 2.14$  for the DH at the species level.

#### SITE SIMILARITY

Familial and species similarity based on the Sørensen's similarity index ( $C_s$ ) was calculated for the MAT and DH Tundra. The MAT

and DH had a family level similarity measurement of  $C_s = 0.655$  and a species-level similarity of  $C_s = 0.453$  (Table 5). The MAT and DH shared the following families: Gnaphosidae, Linyphiidae, Lycosidae, Philodromidae, and Thomisidae. The similarity index for the species within each of these shared families are  $C_s = 0.40$ ,  $C_s = 0.046$ ,  $C_s = 0.52$ ,  $C_s = 0.73$ , and  $C_s = 0.17$ , respectively. The families Araneidae, Clubionidae, Dictynidae, Salticidae, and Theridiidae were unique to only one site (MAT or DH).

The pooled Toolik Lake, Alaska, spider fauna, identified here from within the MAT and DH Tundra, was compared with previous studies performed across arctic and subarctic sites in the Nearctic ranging from the Canadian Arctic Archipelago to southern Alaska by pairwise species-level similarity comparisons (Table 7). We included a wide latitudinal range for our similarity study because we wanted to address the similarity of spider fauna using similar methods (e.g., pan traps and pitfall traps). Furthermore, we wanted to investigate the similarity of studies across a poorly known region. For all sites, pairwise similarity indices ( $C_s$ ) measurements ranged from 0.00 to 0.944 with an average measurement of  $C_s = 0.127 \pm 0.121$  (Table 8). The results of an ANOVA test did not return any significant differences in similarity between all sites. The similarity between arctic sites (collection locales north of the 10 °C July isotherm) is  $C_s = 0.162 \pm 0.168$  and not significant in an ANOVA test ( $p = 0.14$ ). These sites

TABLE 5

Diversity metrics arranged by site, Moist Acidic (MAT) and Dry Heath (DH) tundra and across the whole site (MAT + DH).

| Diversity Metrics – By Site                | MAT      | DH    |
|--|----------|-------|
| Family Richness                            | 8        | 7     |
| Family Evenness                            | 0.322    | 0.408 |
| Shannon's Index – Family Level             | 0.669    | 0.794 |
| Sørensen's Index – Family Level            | 0.655    |       |
| Species Richness                           | 45       | 27    |
| Species Evenness                           | 0.572    | 0.651 |
| Shannon's Index – Species Level            | 2.17     | 2.14  |
| Sørensen's Index – Species Level           | 0.453    |       |
| Diversity Metrics – Total                  | MAT + DH |       |
| Whole Site Family Richness                 | 10       |       |
| Whole Site Family Evenness                 | 0.313    |       |
| Whole Site Shannon's Index – Family Level  | 0.721    |       |
| Whole Site Species Richness                | 51       |       |
| Whole Site Species Evenness                | 0.598    |       |
| Whole Site Shannon's Index – Species Level | 2.352    |       |

include Southhampton Island, Baffin Island, Toolik Lake, the arctic portion of the Yukon Territory, Hazen Camp, Devon Island, and the two Akpatok Island studies.

## Discussion

Over the course of 2004–2006, we found 10 spider families and 45 confirmed species and 6 incompletely identified species. The majority of species found during the course of this study have been reported for the arctic Nearctic region (*see* Dondale and Redner, 1978, 1990; Platnick and Dondale, 1992; Dondale et al., 2003; Paquin et al., 2010) except for one species previously unreported to the Nearctic region, *Agyneta pseudosaxatilis* (Tanasevitch 1984). The number of total spiders captured here was dominated by the family Lycosidae, particularly in the genera *Pardosa*, *Arctosa*, and *Alopecosa*. Spiders belonging to the family Linyphiidae made up the bulk of the species in the collection. Spiders belonging to the families Lycosidae and Linyphiidae are the most abundant families reported in descriptions of northern epigeal spider communities (Danks, 1981; Marusik and Koponen, 2002). Linyphiids tend to be the most speciose spider family across the Nearctic (*see* Leech, 1966; Cotton, 1979; Cutler and Saltmarsh, 1986; Koponen, 1992; Dondale et al., 1997; Pickavance, 2006; Paquin et al., 2010). This was the case for samples taken in this study. Other families were all much less abundant. This fits the general pattern that most families, other than Lycosidae and Linyphiidae, tend to decline in the Nearctic regions (Dondale et al., 1997).

The spider species richness reported here is higher than numbers reported by other studies of Alaskan and Canadian Nearctic spider faunas collected in a singular location. For example, Hillyard (1979) described 2 families with 10 species from his work on Baffin Island, Canada. On the North Slope, Weber (1949) described a small collection taken near Barrow, Alaska, that included six families and 18 putative species. However, it remains unclear as to whether Toolik Lake, Alaska, has a larger number of epigeal spider species than other arctic areas or if this pattern is a result of historic undersampling in the region.

Large collections made in geographically widespread regions (e.g., Yukon Territory from Dondale et al., 1997) or studies in more temperate regions (e.g., Chichagof Island, Alaska, from

TABLE 6

Species richness of spiders collected, arranged by family, from 2004 to 2006 in Moist Acidic Tundra (MAT) and Dry Heath (DH) Tundra at Toolik Lake, Alaska, for Whole Site (MAT + DH), MAT only, and DH only.

|               | MAT + DH | MAT Only | DH Only |
|---------------|----------|----------|---------|
| Araneidae     | 1        | 1        | –       |
| Clubionidae   | 1        | 1        | –       |
| Dictynidae    | 1        | –        | 1       |
| Gnaphosidae   | 4        | 4        | 4       |
| Linyphiidae   | 26       | 23       | 8       |
| Lycosidae     | 11       | 10       | 9       |
| Philodromidae | 2        | 2        | 1       |
| Salticidae    | 1        | –        | 1       |
| Theridiidae   | 1        | 1        | –       |
| Thomisidae    | 3        | 3        | 3       |

Slowik, 2006) reported higher family and species richness compared to the spider fauna captured at Toolik Lake (*see* references in Table 9). This is not surprising given the limited collection area included in this study when compared to a survey of arctic fauna occurring in Yukon Territory. Furthermore, Slowik (2006) reported 95 spider species on Chichagof Island, Alaska. However, this island is located in a temperate rainforest in SE Alaska and a direct comparison of spider fauna might not be appropriate due to differences in rainfall pattern, temperature regimes, etc. We feel that these studies are worth mentioning because of the dearth of large collections made in the northern Nearctic.

We are aware that our method of capture, pitfall trapping, does have drawbacks and is heavily biased towards fauna that are actively moving on the soil surface. We made an effort to supplement pitfall trapping with a second collection method, as recommended by Uetz and Unzicker (1976), to address criticisms in using pitfall trap methods. Sweep netting of tundra vegetation around the pitfall traps in moist acidic and dry heath research plots was conducted in the summer of 2007 within a 0.5 m diameter of each pitfall trap for one minute. We did not capture any arboreal or stem/trunk dwelling spiders (Araneidae, Dictynidae, Salticidae, Tetragnathidae, and Theridiidae) via sweep netting. However, we found them in our pitfall traps, albeit in very low numbers, as less than 0.5% of the total number of specimens collected over the three growing seasons belonged to these families. Lack of sweep netting success could be attributed to a variety of factors, including disturbance of plant canopy from vegetation measurements, cold temperatures, etc.

Despite these criticisms, we did capture a large number of species compared to other studies made in the region. We attribute our high species richness to persistence in trapping effort over time (3 years) and efforts made to trap over the growing season (June–August) when spiders are most active. Our collection efforts yielded a species richness estimate in the upper 95% confidence limit of a rarefaction curve, which leads us to conclude that our collection of fauna captured by pitfall trap is fairly complete and our sampling methods are justified (Fig. 2).

Newcomers to the Arctic are initially presented with what appears a uniform biome in which species are widely distributed (Pickavance, 2006). However, patchiness in distribution of arctic Nearctic spider species has been identified (discussed in Pickavance, 2006). For example, of a collective 55 spider species in six arctic locations, over 50% of species were found in only one location and 90% were found in half or fewer of the sites studied (Pickavance, 2006). In his discussion, Pickavance argues that this



TABLE 7

Pairwise Sørensen's similarity index calculated for Toolik Lake LTER, Alaska, and 13 arctic and subarctic study sites in Alaska (AK) and Canada (CAN). Similarity Indices > 0.25 are shaded.

|   | Belcher Island,<br>CAN | Chichagof<br>Island, AK | AK – VEGA<br>Expedition,<br>1878 | AK – Lindroth<br>Expedition,<br>1958 | Akpatok<br>Island, CAN,<br>Part I, 1933 | Akpatok<br>Island, CAN,<br>Part II, 1936 | Mt. Wrangell,<br>AK (Alpine<br>tundra) | East Bay,<br>South-hampton<br>Island, CAN | Yukon<br>Territory, CAN | Baffin Island,<br>CAN | Toolik Lake,<br>AK, U.S.A.,<br>2004–2006 | Devon Island,<br>CAN | Hazen Camp,<br>CAN |
|---|------------------------|-------------------------|----------------------------------|--------------------------------------|---|--|--|---|-------------------------|-----------------------|--|----------------------|--------------------|
| Spiders of Alaska                         | 0.057                  | 0.181                   | 0.076                            | 0.201                                | 0.023                                   | 0.023                                    | 0.061                                  | 0.030                                     | 0.449                   | 0.008                 | 0.101                                    | 0.016                | 0.015              |
| Belcher Island, CAN                       |                        | 0.047                   | 0.187                            | 0.211                                | 0.308                                   | 0.280                                    | 0.125                                  | 0.118                                     | 0.127                   | 0.047                 | 0.119                                    | 0.000                | 0.043              |
| Chichagof Island,<br>AK                   |                        |                         | 0.000                            | 0.068                                | 0.000                                   | 0.000                                    | 0.036                                  | 0.018                                     | 0.179                   | 0.000                 | 0.041                                    | 0.000                | 0.000              |
| AK – VEGA<br>Expedition, 1878             |                        |                         |                                  | 0.211                                | 0.262                                   | 0.271                                    | 0.070                                  | 0.267                                     | 0.136                   | 0.077                 | 0.237                                    | 0.200                | 0.182              |
| AK – Lindroth<br>Expedition, 1958         |                        |                         |                                  |                                      | 0.180                                   | 0.163                                    | 0.104                                  | 0.121                                     | 0.206                   | 0.066                 | 0.152                                    | 0.045                | 0.085              |
| Akpatok Island,<br>CAN, Part I,<br>1933   |                        |                         |                                  |                                      |   | 0.944                                    | 0.000                                  | 0.000                                     | 0.000                   | 0.069                 | 0.086                                    | 0.000                | 0.188              |
| Akpatok Island,<br>CAN, Part II,<br>1936  |                        |                         |                                  |                                      |   |  | 0.063                                  | 0.171                                     | 0.083                   | 0.074                 | 0.088                                    | 0.000                | 0.200              |
| Mt. Wrangell, AK<br>(Alpine tundra)       |                        |                         |                                  |                                      |   |  |  | 0.000                                     | 0.000                   | 0.08                  | 0.121                                    | 0.000                | 0.071              |
| East Bay,<br>Southampton<br>Island, CAN   |                        |                         |                                  |                                      |   |  |  |   | 0.070                   | 0.286                 | 0.058                                    | 0.462                | 0.452              |
| Yukon Territory,<br>CAN                   |                        |                         |                                  |                                      |   |  |  |   |                         | 0.026                 | 0.201                                    | 0.033                | 0.039              |
| Baffin Island, CAN                        |                        |                         |                                  |                                      |   |  |  |   |                         |                       | 0.066                                    | 0.111                | 0.348              |
| Toolik Lake, AK,<br>U.S.A., 2004–<br>2006 |                        |                         |                                  |                                      |   |  |  |   |                         |                       |  | 0.000                | 0.125              |
| Devon Island, CAN                         |                        |                         |                                  |                                      |   |  |  |   |                         |                       |  |                      | 0.667              |

Study sites referenced, from left to right, Chamberlin and Ivie (1947); Koponen (1992); Slowik (2006); Holm (1970); Holm (1960); Jackson (1933); Davis (1936); Cutler and Saltmarch (1986); Pickavance (2006); Dondale et al. (1997); Hillyard (1979); Leech and Ryan (1972); Leech (1966). Toolik Lake, Alaska, data are taken from current study.

TABLE 8

Mean Sørensen's pairwise similarity index calculated for Toolik Lake LTER, Alaska, and 13 arctic and subarctic study sites in Alaska (AK) and Canada (CAN). *P*-value for the ANOVA test was  $p = 0.25$  and  $f = 1.245$ , indicating no statistically significant differences between sites.

| Site and Reference                                      | Mean Sørensen's |            |
|---|-----------------|------------|
|   | Index           | S.D. $\pm$ |
| Spiders of Alaska – Chamberlin and Ivie (1947)          | 0.160           | 0.123      |
| Belcher Island, CAN – Koponen (1992)                    | 0.128           | 0.095      |
| Chichagof Island, AK – Slowik (2006)                    | 0.044           | 0.065      |
| AK – VEGA Expedition 1878 – Holm (1970)                 | 0.167           | 0.088      |
| AK – Lindroth Expedition 1958 – Holm (1960)             | 0.139           | 0.061      |
| Akpatok Island, CAN, Part I, 1933 – Jackson (1933)      | 0.182           | 0.249      |
| Akpatok Island, CAN, Part II, 1936 – Davis (1936)       | 0.109           | 0.099      |
| Mt. Wrangell, AK – Cutler and Saltmarch (1986)          | 0.063           | 0.043      |
| East Bay, South Hampton Island, CAN – Pickavance (2006) | 0.159           | 0.162      |
| Yukon Territory, CAN – Dondale et al. (1997)            | 0.119           | 0.123      |
| Baffin Island, CAN – Hillyard (1979)                    | 0.104           | 0.113      |
| Toolik Lake, AK, USA, 2004–2006 – current study         | 0.107           | 0.064      |
| Devon Island, CAN – Leech and Ryan (1972)               | 0.118           | 0.211      |
| Hazen Camp, CAN – Leech (1966)                          | 0.186           | 0.196      |

patchiness is a result of dispersion difficulties and localized habitat and microclimate preference, not sampling artifact.

Our results show that this pattern can, again, be seen across 14 separate studies examining the spider fauna of arctic and subarctic sites. The mean similarity index for shared spider species was quite low ( $C_s = 0.127$ ) for all sites. This result suggests that, on average, the epigeal spider species for the sites were 0.873 dissimilar. Mean similarity by site ranged from  $C_s = 0.044$  (Chichagof Island, Alaska; Slowik, 2006) to  $C_s = 0.186$  (Hazen Camp, Canada; Leech, 1966). Interestingly, collection sites located on island features or “islands” of habitat (e.g., alpine tundra habitat) returned lower similarity measures compared to their mainland counterparts. This supports the notion that island areas support fewer species and also have less species diversity. However, our data shows that even large-scale, mainland studies (e.g., Chamberlin and Ivie, 1947; Dondale et al., 1997) have low similarity values ( $C_s = 0.16$  and  $C_s = 0.119$ , respectively). However, these similarity comparisons should be interpreted with caution as the spiders faunas, compared here, were derived from studies taken from a wide range of habitats and latitude, and are therefore not standardized to a similar Nearctic locale.

However, spider faunas in the same collection locality show a strong trend of dissimilarity. While the MAT and DH Tundra sites are only ~3.2 km apart in distance, they have different spider communities. For instance, the species level similarity between the two sites is less than half ( $C_s = 0.453$ ). That is, the two sites share ~45% of the total spider species collected. For this study, the habitats (MAT and DH) each represent a distinct, unique microclimate with differences in plant community, primary production, canopy height, litter layer, and insulation from extreme weather events. The two habitats share only about half of the Toolik Lake species of the most abundant family, Lycosidae ( $C_s = 0.518$ ). Interestingly, the family Linyphiidae was the most speciose (26 species out of 51 collected). However, when the similarity between the linyphiids collected at the MAT and DH is compared, they share less than 5% in common ( $C_s = 0.046$ ). Our

interpretation is that the Linyphiidae are specialists to their given habitat. For example, the spider *Metopobactrus prominulus* was collected in over 200 instances in the DH but none were sampled in the MAT. This species makes up most (91.7%) of all linyphiids collected at the dry heath.

Spider species that are unique to only one site make up more than half (~59%) of the collection. Spiders belonging to family Linyphiidae make up the bulk of the unique, one-site-only species found at Toolik Lake. However, it is important to note that we collected 15 singleton species, 11 belonging to the family Linyphiidae, and clearly, these can only be represented from one habitat or the other, regardless of their true habitat distribution. Therefore, this unique status may be premature and requires more investigation. However, the fact remains that, when we compared our species list across the arctic and subarctic areas of the Nearctic zone, and even locally at Toolik Lake, a strong trend of dissimilarity emerges. At the local scale, the MAT and DH Tundra are statistically different in terms of family abundance and distribution and this claim is further supported by the differences in species richness, evenness, and diversity indices. Pickavance (2006) argued that this patchiness and community divergence is indeed the norm with northern spider communities. This is in contrast to the idea that spider faunas are relatively similar across the northern arctic region (Marusik and Koponen, 2000). The question remains as to why these spider communities are so habitat and locale specific.

Numerous studies have detailed the relationship between vegetative structure and the composition of spider communities, and it is often argued that this is the most important parameter involved in retreat and web site selection (Wise, 1993). Habitat requirements vary according to foraging strategy. Epigeal spiders, especially active hunters belonging to the family Lycosidae, are generally most numerous in warmer, open habitats which do not have a closed canopy (Pajunen et al., 1995). Furthermore, an open canopy is important for the Lycosidae because females maintain optimum temperature for their egg sacs by sunning (Humphreys, 1974). These requirements for a thermally favorable environment, especially given the Arctic's limited growing season, must reasonably extend to the other families mentioned in this study.

The influence of variation in litter on ground-spider communities must also be considered. In one experiment, both litter depth and structural complexity were manipulated on a temperate forest floor (Uetz, 1979). More species of epigeal spiders were present in areas of greater depth and/or litter complexity in all sites studied (Uetz, 1979). Relative familial abundance in the study by Uetz (1979) changed over a gradient of increasing litter depth and complexity, with Lycosidae decreasing and Clubionidae, Thomisidae, and Gnaphosidae increasing. Thus, heterogeneity of the plant community will influence the ground-spider community by the input of different amounts and types of plant litter. Arctic spider communities also change when litter shape, structure, and depth is manipulated, and the community responds differentially in terms of abundance and diversity when the plant community is altered by long-term fertilization experiments at both the MAT and DH Tundra sites (Wyant, 2008).

## Conclusion

Long-term, intensive sampling efforts, as used here, can reveal information concerning undescribed species, range extensions of species, and community level information (e.g., species evenness (*E*), species richness (*S*), and species diversity (*H*). Interestingly, the epigeal spider community at Toolik Lake is more

**TABLE 9**  
**Number of spider families and species in 14 arctic and subarctic localities in Alaska (AK) and Canada (CAN).**

| Place                  | Belcher Island, CAN, 1992             | Chichagof Island, AK, 2006                                 | Akpatok Island, CAN, Part 1, 1931 | Akpatok Island, CAN, Part 2, 1936 | Matanuska Valley, AK, 1947 (Spiders of AK) | Mt. Wrangell, AK (Alpine tundra), 1986 | Vega Expedition, AK, 1878–1880              |
|------------------------|---------------------------------------|--|-----------------------------------|-----------------------------------|--|--|---|
| <b>Latitude North</b>  | 56                                    | 58.1   | 60                                | 60                                | 62   | 62                                     | Collections made in Norway, Siberia, and AK |
| <b>Total Specimens</b> | 1065                                  | 1239   | 125                               | Not Reported                      | Not Reported                               | Not Reported                           | 677   |
| <b>Total Families</b>  | 8                                     | 16   | 3                                 | 3                                 | 16   | 8                                      | 8   |
| <b>Total Species</b>   | 33                                    | 95   | 19                                | 17                                | 247  | 16                                     | 42  |
| <b>Sample Method</b>   | Hand Picking, Litter Seiving, Pitfall | Beat Sheets, Head Lamps, Pitfalls Traps, casual collection | Not Reported                      | Not Reported                      | Not Reported                               | Hand Picking, Sweep Netting            | Not Reported                                |

| Place                  | Lindroth Expedition of 1958 | East Bay, South Hampton Island, CAN, 2006 | Yukon Territory, CAN (Arctic Zone), 1997 | Baffin Island, CAN, 1979 | Toolik Lake, AK, 2004–2006 | Devon Island, CAN, 1972 | Hazen Camp, CAN, 1966 |
|------------------------|-----------------------------|---|--|--------------------------|----------------------------|-------------------------|-----------------------|
| <b>Latitude North</b>  | Throughout AK               | 64  | Throughout Yukon Territory               | 66.71                    | 68.38                      | 76                      | 81.5                  |
| <b>Total Specimens</b> | Not Reported                | 22,816                                    | Not Reported                             | 65                       | 5066                       | 2375                    | 20,600                |
| <b>Total Families</b>  | 9                           | 2   | 9  | 2                        | 10                         | 2                       | 4                     |
| <b>Total Species</b>   | 81                          | 18  | 297                                      | 10                       | 51                         | 8                       | 13                    |
| <b>Sample Method</b>   | Not Reported                | Pitfall traps                             | Not Reported                             | Pitfall Trap             | Pitfall Traps              | Pitfall Traps           | Pan Traps             |

Study sites referenced, from left to right, Koponen (1992); Slowik (2006); Jackson (1933); Davis (1936); Chamberlin and Ivie (1947); Cutler and Saltmarch (1986); Holm (1970); Holm (1960); Pickavance (2006); Dondale et al. (1997); Hillyard (1979); current study; Leech and Ryan (1972); Leech (1966).

Note: Devon Island, Southampton Island, Hazen Camp, Akpatok Island: total numbers include immature specimens; for Baffin Island, Chichagof Island, Toolik Lake, and the Belcher Islands, numbers are adults only. Table modified from Pickavance (2006).




FIGURE 2. Spider species richness estimates for number of samples taken from 2004 to 2006 in Moist Acidic Tundra (MAT) and Dry Heath (DH) Tundra at Toolik Lake, Alaska. 95% confidence intervals (LCI=43; UCI=60) are shown on top and bottom and the mean estimate is in the middle. The ground spider species richness for this study is 51 which approaches the upper 95% confidence limit calculated by rarefaction for 2114 total pitfall traps (PAST Software; Hammer and Harper, 2005).

species rich than generally reported for the Arctic. Moreover, habitat specific spider communities have also been documented for Moist Acidic Tundra and Dry Heath Tundra. Our report of the northward range extension of *Hypsosinga groenlandica*, *Gnaphosa borea*, *Gnaphosa microps*, and *Haplodrassus hiemalis* is intriguing and also confounding, given the recent warming and lengthening of the growing season in the Arctic (Oechel et al., 2000; Serreze et al., 2000; Chapin et al., 2005). Whether these records indeed reflect a northward extension of ranges in response to trends in warming or is due to the fact that the region had not been surveyed extensively, remains unclear, but warrants further study. Arctic areas may have higher spider richness than generally reported but intensive sampling efforts must be used to detect the signal of this trend. Natural history studies that document faunal composition and fundamental community patterns will contribute to baseline data with regards to biotic response to future climate change.

Research over the past three decades indicates that the arctic Nearctic region is fragile and highly susceptible to disturbance such as warming (Hinzman et al., 2005). This region is experiencing dramatic environmental changes that are likely to have profound impacts on the local flora and fauna. Documenting the response of the biotic community to changes in temperature is an important part of monitoring and characterizing possible ecosystem level responses to warming. However, much of the arctic fauna remains poorly studied and much natural history work remains, especially for the invertebrate community. Future efforts should focus on documenting the northern Alaska spider community with active searching (e.g., overturning stones and collecting spiders), hand sorting, quadrat sampling, sweep netting, and the use of beat sheets to address potential gaps in describing the spider community.

### Acknowledgments

We thank Howard Horton, Rodney Simpson, Greg Selby, Andreas David Garcia, Carol Moulton, and Skylar Bayer for their help in the field and laboratory. We also thank Drs. Gerald Callahan, Paula Cushing, and Boris Kondratieff for their quality

contributions towards the development of the present work. We also thank Elizabeth M. Hagen for her constructive comments on earlier drafts of this manuscript. We appreciate the constructive criticism made by two anonymous reviewers which greatly improved the manuscript. Additionally, we thank Don Buckle for helping with the identification of spiders in the family Linyphiidae, and Andrew Karl at Arizona State University for his statistical advice. Finally, we thank the National Science Foundation Office of Polar Programs (OPP 0701295) and the National Science Foundation GK-12 Graduate Student Fellowship program for funding this research (DGE0649876).

### References Cited

- Brower, J. E., Zar, J. H., and von Ende, C. N., 1998: *Field and Laboratory Methods for General Ecology*. Boston: WCB McGraw-Hill, 288 pp.
- Buckle, D. J., Carroll, D., Crawford, R. L., and Roth, V. D., 2001: Linyphiidae and Pimoidae of America north of Mexico: checklist, synonymy, and literature. Part 2. In Paquin, P., and Buckle, D. J. (eds.), *Contributions à la connaissance des Araignées (Araneae) d'Amérique du Nord, Fabriques, Supplément 10*, 89–191.
- Chamberlin, R. V., and Ivie, W., 1947: The spiders of Alaska. *Bulletin of the University of Utah Biological Service*, 37: 1–103.
- Chapin, F. S., III, Shaver, G. R., Giblin, A. E., Nadelhoffer, K. J., and Laundre, J. A., 1995: Responses of arctic tundra to experimental and observed changes in climate. *Ecology*, 76: 694–711.
- Chapin, F. S., III, Sturm, M., Serreze, M. C., McFadden, J. P., Key, J. R., Lloyd, A. H., McGuire, A. D., Rupp, T. S., Lynch, A. H., Schimel, J. P., Beringer, J., Chapman, W. L., Epstein, H. E., Euskirchen, E. S., Hinzman, L. D., Jia, G., Ping, C.-L., Tape, K. D., Thompson, C. D. C., Walker, D. A., and Welker, J. M., 2005: Role of land-surface changes in arctic summer warming. *Science*, 310: 657.
- Cheng, W., Virginia, R. A., Oberbauer, S. F., Gillespie, C. T., Reynolds, J. F., and Tenhunen, J. D., 1998: Soil nitrogen, microbial biomass, and respiration along an Arctic trophosequence. *Soil Science Society of America*, 62: 654–662.
- Cotton, M. J., 1979: A collection of spiders from North-East Greenland. *Arctic*, 32: 71–75.
- Cutler, B., and Saltmarch, R., 1986: Spiders (Araneae) from the alpine zone of the south and west flanks of Mt. Wrangell, Alaska. *Arctic*, 39: 82–84.
- Danks, H. V., 1981: *Arctic Arthropods: a Review of Systematics and Ecology with Particular Reference to the North American Fauna*. Ottawa: Entomological Society of Canada, 608 pp.
- Davis, D. H. S., 1936: A reconnaissance of the fauna of Akpatok Island, Ungava Bay. *Journal of Animal Ecology*, 5: 319–332.
- Doles, J., 2000: *A Survey of Soil Biota in the Arctic Tundra and Their Role in Mediating Terrestrial Nutrient Cycling*. M.S. thesis. Department of Biological Sciences, University of Northern Colorado, Greeley.
- Dondale, C. D., and Redner, J. H., 1978: *The Insects and Arachnids of Canada. Part 5. The Crab Spiders of Canada and Alaska: Araneae: Philodromidae and Thomisidae*. Ottawa: NRC Research Press, 255 pp.
- Dondale, C. D., and Redner, J. H., 1990: *The Insects and Arachnids of Canada. Part 17. The Wolf Spiders, Nursery Web Spiders, and Lynx Spiders of Canada and Alaska: Araneae: Lycosidae, Pisauridae, and Oxyopidae*. Ottawa: NRC Research Press, 383 pp.
- Dondale, C. D., Redner, J. H., and Marusik, Y. M., 1997: Spiders (Araneae) of the Yukon. In Danks, H. V., and Downes, J. A. (eds.), *Insects of the Yukon, Ottawa: Biological Survey of Canada (Terrestrial Arthropods)*. Ottawa: Biological Survey of Canada, 73–113.
- Dondale, C. D., Redner, J. H., Paquin, P., and Levi, H. W., 2003: *The Insects and Arachnids of Canada, Part 23. The Orb Weaving*



- Spiders of Canada and Alaska (Araneae: Uloboridae, Tetragnathidae, Araneidae, Theridiosomatidae)*. Ottawa: NRC Research Press, 371 pp.
- Gough, L., Ramsey, E. A., and Johnson, D. R., 2007: Plant-herbivore interactions in Alaskan arctic tundra change with soil nutrient availability. *Oikos*, 116: 407–418.
- Hammer, O., and Harper, D. A. T., 2005: Palaeontological Statistics, version 1.34. <http://folk.uio.no/ohammer/past/>.
- Hillyard, P. D., 1979: Notes on two collections of spiders from Baffin Island and Greenland. *Bulletin of the British Arachnological Society*, 4: 296–297.
- Hinzman, L. D., Bettez, N. D., Bolton, W. R., Chapin, F. S., III, Dyurgerov, M. B., Fastie, C. L., Griffith, B., Hollister, R. D., Hope, A., Huntington, H. P., Jensen, A. M., Jia, G. J., Jorgenson, T., Kane, D. L., Klein, D. R., Kofinas, G., Lynch, A. H., Lloyd, A. H., McGuire, A. D., Nelson, F. E., Oechel, W. C., Osterkamp, T. E., Racine, C. H., Romanovsky, V. E., Stone, R. S., Stow, D. A., Sturm, M., Tweedie, C. E., Vourlitis, G. L., Walker, M. D., Walker, D. A., Webber, P. J., Welker, J. M., Winer, K. S., and Yoshikawa, K., 2005: Evidence and implications of recent climate change in northern Alaska and other Arctic regions. *Climatic Change*, 72: 251–298.
- Hobbie, S. E., and Chapin, F. S., III, 1998: The response of tundra plant biomass, aboveground production, nitrogen, and CO<sub>2</sub> flux to experimental warming. *Ecology*, 79: 1526–1543.
- Holm, A., 1960: On a collection of spiders from Alaska. *Zoological Bidr. Uppsala*, 33: 109–134.
- Holm, A., 1970: Notes on spiders collected by the “Vega” expedition 1878–1880. *Entomologica Scandinavia*, 1: 188–208.
- Humphreys, W. F., 1974: Behavioral thermoregulation in a wolf spider. *Nature*, 251: 502–503.
- Jackson, A. R., 1933: Results of the Oxford University Expedition to Akpatok in 1931: Araneae. *Proceedings of the Zoological Society*, 102: 145–159.
- Koponen, S., 1992: Spider fauna (Araneae) of the Low Arctic Belcher Islands, Hudson Bay. *Arctic*, 45: 358–362.
- Leech, R. E., 1966: The spiders of Hazen Camp 81°49'N, 71°18'W. *Quaestiones Entomologicae*, 2: 153–212.
- Leech, R. E., and Ryan, J. K., 1972: Notes on Canadian Arctic spiders (Araneida), mainly from Devon Island, NWT. *Canadian Entomologist*, 104: 1787–1791.
- Magurran, A., 2004: *Measuring Biological Diversity*. Oxford: Blackwell Publishing, 260 pp.
- Marusik, Y. M., and Koponen, S., 2000: Circumpolar diversity of spiders: implications for research, conservation and management. *Annales Zoologici Fennici*, 37: 265–269.
- Marusik, Y. M., and Koponen, S., 2002: Diversity of spiders in boreal and Arctic zones. *The Journal of Arachnology*, 30: 205–210.
- Marusik, Y. M., Logunov, D. V., and Koponen, S., 2000: *Spiders of Tuva, South Siberia*. Magadan: Russian Academy of Science, 252 pp.
- Nadelhoffer, K. J., Giblin, A. E., Shaver, G. R., and Laundre, J. A., 1991: Effects of temperature and substrate quality on element mineralization in six Arctic soils. *Ecology*, 72: 242–253.
- Oechel, W. C., Vourlitis, G. L., Hastings, S. J., Zulueta, R. C., Hinzman, L., and Kane, D., 2000: Acclimation of CO<sub>2</sub> exchange in the Alaskan Arctic in response to decadal climate warming. *Nature*, 406: 978–981.
- Pajunen, T., Haila, Y., Halme, E., Niemela, J., and Punttila, P., 1995: Ground dwelling spiders (Arachnida, Araneae) in fragmented old forests and surround managed forests in southern Finland. *Ecography*, 18: 62–72.
- Paquin, P., Buckle, D. J., Duperre, N., and Dondale, C. D., 2010: Checklist of the spiders (Araneae) of Canada and Alaska. *Zootaxa*, 2461: 1–170.
- Pickavance, J. R., 2006: The spiders of East Bay, Southampton Island, Nunavut, Canada. *Arctic*, 59: 276–282.
- Platnick, N. I., 2010: The World Spider Catalog, version 10.5. American Museum of Natural History, online at <http://research.amnh.org/entomology/spiders/catalog/index.html>.
- Platnick, N. I., and Dondale, C. D., 1992: *The Insects and Arachnids of Canada. Part 19. The Ground Spiders of Canada and Alaska: Araneae: Gnaphosidae*. Ottawa: NRC Research, 297 pp.
- R Development Core Team, 2005, *R: A Language and Environment for Statistical Computing, Reference Index Version 2.x.x*. Vienna, Austria: R Foundation for Statistical Computing, ISBN 3-900051-07-0, URL <http://www.R-project.org>.
- SAS Institute Inc, 2000, *SAS Online Doc 9.1.2. JMP Statistics and Graphics Guide, version 5*. Cary, North Carolina: SAS Institute Inc.
- Serreze, M. C., Walsh, J. E., Chapin, F. S., III, Osterkamp, T., Dyurgerov, M., Romanovsky, V., Oechel, W. C., Morison, J., Zhang, T., and Barry, R. G., 2000: Observational evidence of recent change in the northern high-latitude environment. *Climatic Change*, 46: 159–207.
- Shannon, C. E., 1948: A mathematical theory of communication. *Bell System Technical Journal*, 27: 379–423, 623–656.
- Shaver, G. R., and Chapin, F. S., III, 1991: Production: biomass relationships and element cycling in contrasting arctic vegetation types. *Ecological Monographs*, 61: 1–31.
- Slowik, J., 2006: A survey if the spiders (Arachnida, Araneae) of Chichagof Island, Alaska, USA. *Journal of the Entomological Society of British Colombia*, 103: 61–70.
- Ubick, D., Paquin, P., Cushing, P. E., and Roth, V., 2005: *Spiders of North America: an Identification Manual*. Poughkeepsie, New York: American Arachnological Society, 377 pp.
- Uetz, G. W., 1979: The influence of variation in litter habitats on spider communities. *Oecologia*, 40: 29–42.
- Uetz, G. W., and Unzicker, J. D., 1976: Pitfall trapping in ecological studies of wandering spiders. *Journal of Arachnology*, 3: 101–111.
- Walker, D. A., 1998: The geobotanical atlas of the Toolik Lake and Kuparuk River region, Alaska. <http://geobotany.uaf.edu/arcticgeobot/>.
- Weber, N. A., 1949: Late summer invertebrates, mostly insects, of the Alaska Arctic Slope. *Entomological News*, 60: 118–128.
- Weber, N. A., 1950: A survey of the insects and related arthropods of Arctic Alaska. Part 1. *Transactions of the American Entomological Society*, 76: 147–206.
- Wise, D. H., 1993: *Spiders in Ecological Webs*. Cambridge: Cambridge University Press, 328 pp.
- Wyant, K. A., 2008: *Effects of Soil Fertilization and Herbivore Exlosures on the Ground Spider Community at Toolik Lake, Alaska*. M.S. thesis. Colorado State University–Fort Collins.