

Molecular Cloning and Analysis of the Tryptophan oxygenase Gene in the Silkworm, *Bombyx mori*

Authors: Yan, Liu, Zhi-Qi, Meng, Bao-Long, Niu, Li-Hua, He, Hong-Biao, Weng, et al.

Source: Journal of Insect Science, 8(54) : 1-7

Published By: Entomological Society of America

URL: <https://doi.org/10.1673/031.008.5401>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.



Molecular cloning and analysis of the *tryptophan oxygenase* gene in the silkworm, *Bombyx mori*

Liu Yan^{1,a}, Meng Zhi-Qi^{1,b}, Niu Bao-Long^{1,c}, He Li-Hua^{1,d}, Weng Hong-Biao^{1,e}, and Shen Wei-Feng^{1,f}

¹Sericulture Research institute, Zhejiang Academy of Agricultural Sciences, Hangzhou, 310021, China

Abstract

A *Bombyx mori* L. (Lepidoptera: Bombycidae) gene encoding tryptophan oxygenase has been molecularly cloned and analyzed. The tryptophan oxygenase cDNA had 1374 nucleotides that encoded a 401 amino acid protein with an estimated molecular mass of 46.47 kDa and a PI of 5.88. RT-PCR analysis showed that the *B. mori tryptophan oxygenase* gene was transcribed in all examined stages. Tryptophan oxygenase proteins are relatively well conserved among different orders of arthropods.

Keywords: eye pigmentation, Lepidoptera, *vermillion* gene

Abbreviations: BMTO - *Bombyx mori* tryptophan oxygenase

Correspondence: ^aliuyanmyg@yahoo.com.cn, ^bmengzq2001@sina.com, ^cniubaolong@126.com, ^dhelh1968@163.com, ^ewenghb2006@126.com, ^fweifsh@126.com

Received: 4 June 2007 | **Accepted:** 11 November 2007 | **Published:** 23 September 2008

Copyright: This is an open access paper. We use the Creative Commons Attribution 3.0 license that permits unrestricted use, provided that the paper is properly attributed.

ISSN: 1536-2442 | Volume 8, Number 54

Cite this paper as:

Yan L, Zhi-Qi M, Bao-Long N, Li-Hua H, Hong-Biao W, Wei-Feng S. 2008. Molecular cloning and analysis of the *tryptophan oxygenase* gene in the silkworm, *Bombyx mori*. 7 pp. *Journal of Insect Science* 8:54, available online: insectscience.org/8.54

Introduction

The silkworm, *Bombyx mori* L. (Lepidoptera: Bombycidae), is an important agriculturally insect that has been domesticated and used in silk production for about 5000 years. As a key model insect for the Lepidoptera, which include many destructive agricultural pests (Tomita et al. 2003), analysis of the genome of the silkworm is important.

The pigments present in the insect eye, usually both pteridine and ommochrome, determine its color (Summers et al. 1982). In *Drosophila*, a large number of mutations are known to affect the pigmentation of the compound eye, including those that affect the biosynthesis or transport of ommochrome (brown) and pteridine (red) pigments (Beadle and Ephrussi 1937). Ommochrome has been recently demonstrated to contribute to eye coloration in mosquitoes and beetles (Beard et al. 1995; Lorenzen et al. 2002; Fabrick et al. 2004). One of the genes involved in the ommochrome biosynthetic pathway is tryptophan oxygenase (TO), an enzyme that converts tryptophan to N-formylkynurenine which is then converted to kynurenine by kynurenine formidase (Ferre et al. 1985; Takikawa et al. 1986). Tryptophan catalyzes the first step in the synthesis of the brown eye pigment of the fly (Linzen et al. 1974; Walker et al. 1986) and is an essential amino acid that is required in several physiological processes in addition to protein synthesis. Early work on *Drosophila* eye color mutants revealed that *vermilion* (*v*) and *cinnabar* (*cn*) are involved in ommochrome production (Beadle and Ephrussi 1937). The *vermilion* gene has been shown to encode tryptophan oxygenase and has been found to be a potentially useful germline transformation marker (Lorenzen et al. 2002). The *tryptophan oxygenase* gene has also been identified from *Anopheles gambiae* (Mukabayire et al. 1996), *Aedes aegypti* (Fang and Li 2001), *Tribolium castaneum* (Lorenzen et al. 2002), *Plodia interpunctella* (Fabrick et al. 2004), and *Shistocerca americana* (Dong and Friedrich 2005).

Abraham et al. (2000) examined the *ABC transporter* genes in *B. mori* that are among the eye- and egg-color mutations affecting the synthesis and accumulation of ommochrome pigments in *B. mori*. They are homologous to the *Drosophila white* gene, and are involved in transporting pigment precursor (Abraham et al. 2000). The protein sequence of *B. mori* kynurenine 3-monooxygenase showed high identity with *cinnabar* (*cn*) in *Drosophila* (Quan et al. 2001; Lorenzen et al. 2002). Yet little is known about the *vermilion* homologous gene in *B. mori*. Here, the *B. mori tryptophan oxygenase* gene was cloned and analyzed.

Materials and Methods

Insects feeding and sample collection

Larvae of the normal strain (*Daizo*) were reared routinely on mulberry leaves. The temperature ranged from 25 to 27 °C and humidity varied between 70 and 80%.

Preparation of RNA and RT-PCR

Total RNA was separately extracted from samples using Trizol. RNA was reverse-transcribed by using oligo (dT)₁₈ adaptor primer (Sangon Bio, www.sangon.com) and avian myeloblastosis virus reverse transcriptase (Promega, www.promega.com) at 42 °C for 60 min. cDNA fragments were amplified using Ex Taq (Takara Bio, www.takara-bio.com) with the following degenerate primers 5' TAY GAR YTN TGG TTY AAR CA 3' (Fto1, sense) and 5' CAT TKC KTT GCA CCA TSA WMA CRT GAT T 3' (Rto1, antisense). Thermal cycling conditions were as follows: 94 °C for 5 min; 30 cycles at 95 °C for 30 s, 50 °C for 30 s, and 72 °C for 1 min. The last cycle was followed by an extension at 72 °C for 10 min. Amplified products were separated by gel electrophoresis on 1.2% agarose gels at 100 v for approximately 1 h using a 1 x TAE buffer (40 mM Tris acetate and 2 mM EDTA in water). After electrophoresis, the gel was stained for 30 min in 0.01% SYBRTM Green I nucleic acid gel stain (FMC Bio, www.fmc.com). Following purification of PCR products by Biospin Gel Extraction Kit (BioFlux Bio, www.fluxionbio.com), the purified PCR fragments were TA-cloned into pMD18-T (Takara Bio.). The insertions were identified by PCR amplification using M13 forward and reverse primers before sequencing. No PCR products were produced when cDNA template was excluded during reverse transcription polymerase chain reaction.

RACE method

Rapid amplification of cDNA ends (RACE) method was applied to obtain full-length cDNAs. cDNA was synthesized from 2 µg of total RNA using BD SMART RACE cDNA Amplification Kit (BD Biosciences, www.bdbiosciences.com). The 5' and 3' cDNA ends were obtained by touchdown PCR with LA Taq (Takara Bio) using a universal primer mixture (UPM) and the following gene-specific primers (GSPs): 5' GTA ATG CCC TCA CAG AAT CCA CTT 3' (Bmtoa, 5'RACE) and 5' GTC TCA TCA CGA AAT GGC GTT ACA 3' (Bmtob, 3'RACE) according to manufacture's instructions. RT-PCR products were gel purified, cloned into pMD18-T (Takara Bio) and sequenced.

To acquire the integral cDNA of *Bmto*, primer 5' CAT TGA AAT GGC GTG TCC TAT GAG 3' (Fto2, Sense) and 5' GTT ATA AAG AAG CTT CAA GGC CGT 3' (Rto2, Antisense) for the gene encoding the *B. mori* homolog of the tryptophan oxygenase were used in amplification. The PCR product was subcloned into pMD18-T and sequenced.

Analysis of the expression at different developmental stages

For developmental analysis, total RNA was extracted from various stages of development (egg, third instars, fifth instars, pupae and adult). RT-PCR was performed as described above. Real-time PCR was performed using the SYBR premix EX Taq™ kit (Takara Bio). The primers designed for *Bmto* were 5' ACA CGC ACG GGT TCA ACT TCT 3' (FBmto-real, Forward) and 5' ATG TGA CAG CCT CCT TTC TCC T 3' (RBmto-real, Reverse), and for the *B. mori* Actin 1 that was used as internal control were 5' ACC CAT CTA CGA AGG TTA CGC 3' (FBmActin, Forward) and 5' ACG AAC GAT TTC CCT CTC AGC 3' (RBmActin, Reverse), yielded 212 and 142 bp bands, respectively. PCR amplification and fluorescence detection were performed using the DNA Engine Option 2 under the following thermal cycle conditions: 95 °C for 1 min, 45 cycles of 95 °C for 10 s, 60 °C for 20 s. To reach reproducibility, each sample was performed three times. Ct values of *Bmto* were calculated to the actual concentrations based on the standard curve. *BmActin* transcripts were used to standardize the different *Bmto* cDNA samples.

Sequence analysis

Predictions of isoelectric point and molecular weight were carried out at <http://cn.expasy.org>. The amino acid sequence of BmTO was submitted to predict secondary structure at <http://npsa-pbil.ibcp.fr> and conserved protein domain at <http://www.ncbi.nlm.nih.gov/>. Alignment of deduced amino acids from cDNA clones was made using DNAMAN software. A phylogenetic tree based on deduced amino acid difference was constructed by NJ (Neighbor-joining) method using PHYLIP (<http://bioweb.pasteur.fr/seqanal/interfaces/protpars.html>). Reliability of the NJ tree was assessed by interior branch test, using 1000 replications.

Results

Identification of cDNA sequences encoding BmTO

Using the degenerate primers, a partial *Bmto* cDNA was obtained. RACE was used to complete the missing 5' and 3' ends of the cDNA. The cDNA consisted of 1374 bp long with an open reading frame of 401 amino acids (Figure 1). BmTO (*B. mori* tryptophan oxygenase) was assigned its name because of its similarity to the known tryptophan oxygenase protein.

Bmto expression profile in B.mori larvae

To examine the expression of *Bmto* gene in various life stages, real time RT-PCR using total RNA obtained from different instars was performed. *Bm actin* was used as an internal control. The level of *Bmto*'s expression was quantified by calculating the ratio of *Bmto*/*BmActin* of the

same sample. As shown in Figure 2, *Bmto* was expressed in all samples examined. Transcript levels were low in larvae and adults, and much greater in embryos and pupae.

Comparison of BmTO to other tryptophan oxygenase proteins

The predicted molecular weight and pI of *B. mori* tryptophan oxygenase were 46.47 kDa and 5.88, respectively. BmTO was not predicted to be a secreted protein as determined by SignalP. There was 60.85% alpha helix, 30.92% random coil and 8.23% extended strand in the secondary structure of BmTO (Figure 3).

To assess the relatedness of the BmTO to tryptophan oxygenase proteins from other species, identities were calculated based on a Clustal alignment including nine tryptophan oxygenase protein sequences (Figure 4). Amino acid comparisons revealed 87.2% identity between the deduced protein of BmTO and previously known tryptophan oxygenases from the moth, *Plodia interpunctella* (Figure 4, 5). Identity of tryptophan oxygenase proteins was high within the arthropods (at least 65.1%), which verified that *tryptophan oxygenase* gene was relatively conserved among the insects.

Discussion

Eye pigments of *D. melanogaster* were determined by specific enzymes of biosynthesis, ATP binding cassette transporters of precursors of pigments and pigment granules (Lloyd et al. 1998). Tryptophan oxygenase is an essential amino acid of insects as its absence or deficiency results in eye color defects in *Drosophila* and other insect species (Summers et al. 1982; Miyashita et al. 1994; White et al. 1996; Lorenzen et al. 2002; Fabrick et al. 2004). In the present study, the *Bombyx mori* tryptophan oxygenases homolog (*Bmto*) was successfully cloned and analyzed. *Bmto* was expressed at a wide variety of developmental stages, which was consistent with that of ommochromes (Sawada et al. 2000). BmTO is highly similar to other tryptophane oxygenases, including those regions of the protein that are constrained by functional requirements among diverse species.

Dominant phenotypic markers such as eye color genes can be used in the development of transgenic organisms. These genes are not limited by cell autonomy and generate an easily scored visible phenotype when introduced into the appropriate mutant background. The use of such genes eliminates the need for specialized detection systems, thus making transformation-based protocols more widely accessible. Fridell and Searles (1991) constructed a germ line transformation vector that utilizes the *vermillion* as the selectable marker gene, which encodes the enzyme tryptophan oxygenase in *Drosophila*. Loukeris et al. (1995) developed a transformation system for the medfly *Ceratitis*

1	GCACGTATCTGTGTCAAGTCTTGTGCTACATTTAATCAATAATTATAATTAGTGAAC
61	AAAATAAAGTCAAAACAAATAGTGCATTGAA ATG GGCGTGTCTATGAGATCAGCGATCGA
1	M A C P M R S A I D
121	CGAAATAAATGGCCAAGAAGGCGCGCACCTTGGTAACGAGGCCGGCATGCTATACGGGGA
11	E I N G Q E G A H L G N E A G M L Y G E
181	ATATTTGATGCTGGATAAACTGCTGTCTGCGCAGAGGATGCTCAGCGCTGAGTCCTCTAA
31	Y L M L D K L L S A Q R M L S A E S S K
241	ACCTGTACACGATGAGCATCTCTTCATCATCACACCAAGCCTACGAACTCTGGTTCAA
51	P V H D E H L F I I T H Q A Y E L W F K
301	GCAGATAATATTTGAAGTGGATTCTGTGAGGGCATTACTTAACGTGGAAGGCTTGGACGA
71	Q I I F E V D S V R A L L N V E G L D E
361	AAGCCACACCATGGAGATATTGAAGAGACTCAACAGGATTGTGCTCATCTTGAAACTGCT
91	S H T M E I L K R L N R I V L I L K L L
421	GGTGGATCAAGTGATGATTTTGGAGACGATGACACCGCTCGACTTCATGGATTTCAGGCA
111	V D Q V M I L E T M T P L D F M D F R H
481	CTATCTCCGCCAGCATCAGGTTTCCAGAGCTTGCAGTTCAGACTACTAGAAAACAAGCT
131	Y L R P A S G F Q S L Q F R L L E N K L
541	TGGTTTGAACAAGCCCTGCGCGTGAAGTACAATCAAATTATCAAACCGTGTGGCGA
151	G L K Q A L R V K Y N Q N Y Q T V F G D
601	CGACCCTGAAGCTATGGACTCCTTACAAAACCTGAACAAGAACCTGCACTGCTAGCGCT
171	D P E A M D S L Q K P E Q E P A L L A L
661	GATCGAGCGTTGGCTGGAGCGTACTCCCGGATTGAACACGCACGGGTTCAACTTCTGGGG
191	I E R W L E R T P G L N T H G F N F W G
721	GAAATTCCAGGCAACAGTCAATAAAATGCTGAAGGAAGATATAGAGACTGCTAATCTTGA
211	K F Q A T V N K M L K E D I E T A N L E
781	GTCCATAGACGCTGTGCGCCGCCATCGGCTTCAAGATGTGGAGAATAGAAGGGAGATTTA
231	S I D A V R R H R L Q D V E N R R E I Y
841	TCGTTTCGATATTTGATCCGACTGTGCATGACGCGCTCAGGTCCAGAGGAGAAAGGAGGCT
251	R S I F D P T V H D A L R S R G E R R L
901	GTCACATAGAGCACTCCAAGGTGCAATCATGATCACGTTTACC CGGACGAGCCTCGGTT
271	S H R A L Q G A I M I T F Y R D E P R F
961	CTCGCAGCCGCATCAGCTCCTCACTCTGCTGATGGACATCGACAGTCTCATCACGAAATG
291	S Q P H Q L L T L L M D I D S L I T K W
1021	GCGTTACAACCACGTTATAATGGTGCAAAGGATGATCGGGTCGCAACAGCTCGGGACCGG
311	R Y N H V I M V Q R M I G S Q Q L G T G
1081	GGGCTCGTCTGGCTATCAGTATCTACGGTCAACATTAAGCGATCGTTACAAAGTCTTTCT
331	G S S G Y Q Y L R S T L S D R Y K V F L
1141	TGATCTGTTCAATTTGTCCACGTTCTGTTGCCTCGTGCCCTAATACCGCCACTGGACGA
351	D L F N L S T F L L P R A L I P P L D E
1201	AGACATGAAACGCAGCCTCAATCTCACATGGGGAGACAACGTCAGGGAAAACGGAGATGA
371	D M K R S L N L T W G D N V R E N G D E
1261	GACGACCCCGCAAACGGCCTTGAAGCTTCTTTA TAAC GAAAAATTATAATGTGTCCTGA
391	T T P Q N G L E A S L *
1321	CAATATTTTAAAGACATATTTTATTACATTTTAAATTCAAAAAAAAAAAAAAAAA

Figure 1. The nucleotide and deduced amino acid sequence of *Bmto* cDNA. Nucleotides are numbered on the left of each line. The deduced amino acid sequence is shown below the nucleotide sequence and numbered from the first methionine. The primer sites are indicated by arrows. The amino acid of conserved tryptophan oxygenase domain is shadowed. Initiation and termination codons are shown in bold face.

capitata with the eye color gene white. White et al. (1996) linked *vermilion* cDNA of *D. melanogaster* to the inducible hsp82 promoter of *D. pseudoobscura*. This marker rescued adult eye color in a strain of *Musca domestica* that was homozygous for a mutant tryptophan oxygenase gene

(White et al. 1996). Besansky et al. (1997) placed *Anopheles gambiae* cDNA encoding tryptophan oxygenase under the control of the constitutive baculovirus promoter, *ie-1*. This chimeric construct, expressed transiently in *vermilion* mutants of *D. melanogaster*, partially rescued adult eye





Figure 3. Secondary structure of the *Bombyx mori* tryptophan oxygenase determined by <http://npsa-pbil.ibcp.fr>. Alpha helices are shown in blue, β -sheets in red and random coils in magenta.

color (Besansky et al. 1997). Genes such as *Bmto* may prove invaluable as transformation markers in the future.

Acknowledgments

We thank Prof. Lu Chang-De for thoughtful advice, critical reading of manuscript. We also thank Wang Shou-Xian and Xe Su-Feng for help with insect feeding. This work was supported by the National Hi-Tech Research and Development Program (863) of China (No. 2006AA10A119).

References

Abraham EG, Sezutsu H, Kanda T, Sugasaki T, Shimada T, Tamura T. 2000. Identification and characterisation of a silkworm ABC transporter gene homologous to *Drosophila* white. *Molecular and General Genetics* 264: 11-19.

Beadle GW, Ephrussi B. 1937. Development of eye colors in *Drosophila*: diffusible substances and their interrelations. *Genetics* 22: 76-86.

Beard CB, Benedict MQ, Primus JP, Finnerty V, Collins FH. 1995. Eye pigments in wild-type and eye-color mutant strains of the African malaria vector *Anopheles gambiae*. *Journal of Heredity* 86: 375-380.

Besansky NJ, Mukabayire O, Benedict MQ, Rafferty CS, Hamm DM, Mcnitt L. 1997. The *Anopheles gambiae* tryptophan oxygenase gene expressed from a baculovirus promoter complements *Drosophila melanogaster* vermilion. *Insect Biochemistry and Molecular Biology* 27: 803-805.

Dong Y, Friedrich M. 2005. Nymphal RNAi: systemic RNAi mediated gene knockdown in juvenile grasshopper. *BMC Biotechnology* 5: 25

Fabrick JA, Kanost MR, Baker JE. 2004. RNAi induced silencing of embryonic tryptophan oxygenase in the pyralid moth, *Plodia interpunctella*. *Journal of Insect Science* 4: 15
Available online at <http://insectscience.org/4.15>




Figure 4. Sequence alignment of tryptophan oxygenase proteins from *Bombyx mori*, *Tribolium castaneum* (Tc, NP_001034499), *Anopheles gambiae* (Ag, XP_312204), *Drosophila melanogaster* (Dm, NP_511113), *Mayetiola destructor* (Md, ABC69733) and *Plodia interpunctella* (Pi, AAR24625). Completely conserved residues are indicated below the alignment.

Fang J, Li J. 2001. Isolation and characterization of tryptophan oxygenase gene of *Aedes aegypti*. *Direct sequence submission to NCBI database* (AF325458)

Ferre J, Real MD, Mensua JL, Jacobson KB. 1985. Xanthurenic acid 8-o- β -D-glucoside, a novel tryptophan metabolite in eye-color mutants of *Drosophila melanogaster*. *The Journal of Biological Chemistry* 260: 7509-7514.

Fridell YW, Searles LL. 1991. *Vermilion* as a small selectable marker gene for *Drosophila* transformation. *Nucleic acids research* 19: 5082

Linzen B. 1974. The tryptophan to ommochrome pathway in insects. *Advances in Insect Physiology* 10: 117-246.


Lloyd V, Ramaswami M, Kramer H. 1998. Not just pretty eyes: *Drosophila* eye colour mutations and lysosomal delivery. *Trends in Cell Biology* 8: 257-259.

Lorenzen MD, Brown SJ, Denell RE, Beeman RW. 2002. Cloning and characterization of the *Tribolium castaneum* eye-color genes encoding tryptophan oxygenase and kynurenine 3-monooxygenase. *Genetics* 160: 225-234.

Loukeris TG, Livadaras I, Arca B, Zabalous S, Savakis C. 1995. Gene transfer into the medfly, *Ceratitis capitata*, with a *Drosophila hydei* transposable elements. *Science* 270: 2002-2005.

Quan G, Kim I, Komoto N, Sezutsu H, Ohet M, Shimada T, Kanda T, Mita K, Kobayashi M, Tamura T. 2001. Characterization of the *kynurenine 3-monooxygenase* gene in the *white egg 1* (w-1) mutant of the silkworm, *Bombyx mori*. *Direct sequence submission to NCBI database* (BAB62419)

Miyashita NT, Langley CH. 1994. Restriction map polymorphism in the forked and vermilion region of *Drosophila melanogaster*. *The Japanese Journal of Genetics* 69: 297-305.



Mukabayire O, Cornel AJ, Dotson EM, Collins FH, Basansky NJ. 1996. The *tryptophan oxygenase* gene of *Anopheles gambiae*. *Insect Biochemistry and Molecular Biology* 26: 525-528.

Sawada H, Nakagoshi M, Mase K, Yamamoto T. 2000. Occurrence of ommochrome-containing pigment granules in the central nervous system of the silkworm, *Bombyx mori*. *Comparative Biochemistry and Physiology* 125: 421-428.

Summers KM, Howells AJ, Pylotis NA. 1982. Biology of eye pigmentation in insects. 29>Advances in Insect Physiology 29: 374-375.

Takikawa O, Yoshida R, Kido R, Hayaishi O. 1986. Tryptophan degradation in mice initiated by indoleamine 2,3-dioxygenase. *The Journal of Biological Chemistry* 261: 3648-3653.

Walker AR, Howells AJ, Tearle RG. 1986. Cloning and characterization of the *vermillion* gene of *Drosophila melanogaster*. *Molecular and General Genetics* 202: 102-107.

White LD, Coates CJ, Atkinson PW, O'Brochta DA. 1996. An eye color gene for the detection of transgenic non-*Drosophilid* insects. *Insect Biochemistry and Molecular Biology* 26: 641-644.