

**Pathogenicity of the Fungus, *Aspergillus clavatus*,
Isolated from the Locust, *Oedaleus senegalensis*,
Against Larvae of the Mosquitoes *Aedes aegypti*,
Anopheles gambiae and *Culex quinquefasciatus***

Authors: Seye, Fawrou, Faye, Oumar, Ndiaye, Mady, Njie, Ebrima, and Marie Afoutou, José

Source: Journal of Insect Science, 9(53) : 1-7

Published By: Entomological Society of America

URL: <https://doi.org/10.1673/031.009.5301>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.



Pathogenicity of the fungus, *Aspergillus clavatus*, isolated from the locust, *Oedaleus senegalensis*, against larvae of the mosquitoes *Aedes aegypti*, *Anopheles gambiae* and *Culex quinquefasciatus*

Fawrou Seye^{1,a}, Oumar Faye^{2,b}, Mady Ndiaye^{1,c}, Ebrima Njie^{3,d} and José Marie Afoutou^{2,e}

¹ U.E.R.B.V., Laboratory of Reproduction Biology, Animal Biology Department, Faculty of Sciences and Technics, Cheikh Anta Diop University of Dakar, Senegal. P.O. Box: 5005, Dakar Fann, Senegal

² Laboratoire d'Histologie, Embryologie et Cytogénétique, Faculté de Médecine, Pharmacie et d'Odontostomatologie, Université Cheikh Anta Diop de Dakar

³ Department of Agriculture and Biological Sciences P.O. Box: 3530 Gambia, University of the Gambia

Abstract

The use of insect pathogenic fungi is a promising alternative to chemical control against mosquitoes. Among the Hyphomycetes isolated from insects for mosquito control, the genus *Aspergillus* remains the least studied. In September 2005, four fungi were isolated from the Senegalese locust, *Oedaleus senegalensis* Kraus (Orthoptera: Acrididae), collected in Dakar, Senegal. One of these fungi, identified as *Aspergillus clavatus*, Desmazières (Eurotiales: Trichocomaceae) was highly pathogenic against larvae of the mosquitoes *Aedes aegypti* L., *Anopheles gambiae* s.l. Giles and *Culex quinquefasciatus* Say (Diptera: Culicidae). An application of 1.2 mg/ml dry conidia yielded 100% mortality after 24 hours against both *Ae. aegypti* and *Cx. quinquefasciatus* while with *An. gambiae* it was 95%. With unidentified species in the genus *Aspergillus*, mortality after 24 h was <5% against all the larval species. Application of *A. clavatus* produced in a wheat powder medium using doses ranging between 4.3 to 21x10⁷ spores/ml, caused 11 to 68% mortality against *Cx. quinquefasciatus* at 24h, and 37 to 100% against *Ae. aegypti*. Microscopic observations showed fungal germination on both *Ae. aegypti* and *Cx. quinquefasciatus* larvae. Histological studies revealed that *A. clavatus* penetrated the cuticle, invaded the gut and disintegrated its cells. Some *Cx. quinquefasciatus* larvae, treated with *A. clavatus* reached the pupal stage and produced infected adults. However, the infection was mainly located on the extremity of their abdomen. These results suggest that *A. clavatus* could be an effective tool to manage mosquito proliferation.

Keywords: biological control, entomopathogenic fungi

Correspondence: ^afawrou@yahoo.fr, ^bofaye@ucad.sn, ^cmydya2001@yahoo.fr, ^debrimanjie@hotmail.com, ^ejmafoutou@hotmail.com

Received: 25 December 2007 | **Accepted:** 3 March 2008 | **Published:** 13 July 2009

Associate Editor: Fernando Vega was editor of this paper

Copyright: This is an open access paper. We use the Creative Commons Attribution 3.0 license that permits unrestricted use, provided that the paper is properly attributed.

ISSN: 1536-2442 | Vol. 9, Number 53

Cite this paper as:

Seye F, Faye O, Ndiaye M, Njie E, Afoutou JM. 2009. Pathogenicity of the fungus, *Aspergillus clavatus*, isolated from the locust, *Oedaleus senegalensis*, against larvae of the mosquitoes *Aedes aegypti*, *Anopheles gambiae* and *Culex quinquefasciatus*. 7pp. *Journal of Insect Science* 9:53, available online: insectscience.org/9.53

Introduction

Mosquito-borne diseases currently represent a great health threat in tropical and subtropical climates. As an alternative to chemical insecticides, natural products (Koua, 1998; Batra et al., 1998; Ravindra et al., 2002), predatory fish (Legner 1995; Karch and Coz 1983; Fillingner et al. 2003) and entomopathogenic fungi (Nnakumusana 1985; Su et al. 2001; Scholte et al. 2003) are frequently used in mosquito control. Fungi such as *Metarhizium anisopliae* (Metchnikoff) Sorokin (Moniliales) and *Beauveria bassiana* (Balsamo) Vuillemin (Clavicipitaceae), commonly found on terrestrial insects can also kill mosquito larvae (Alves et al. 2002; Silva et al. 2004a). Studies have shown that mosquito larvae are susceptible to infections by fungi such as *Leptolegna chapmanii* (Lord and Fukuda, 1988), *M. anisopliae* (Riba et al. 1986; Lacey et al. 1988; Alves et al. 2002; Silva et al. 2004a; Wright et al. 2005), *B. bassiana* (Alves et al. 2002), *Aspergillus parasiticus* (Hati and Ghosh 1965), *Aspergillus* spp. (de Moraes et al. 2001), *Aspergillus flavus*, *A. parasiticus*, *Penicillium falicum*, *Fusarium vasinfectum* and *Trichoderma viride* (Govindarajan et al. 2005). Among the various *Aspergillus* species known to infect mosquitoes, *A. clavatus* Desmazières (Eurotiales: Trichocomaceae) has not been examined as a possible biological control agent. This study assessed the pathogenicity of an *A. clavatus* strain against larval stages of various species of mosquitoes.

Materials and Methods

Isolation

In September 2005, locusts of the species *Oedaleus senegalensis* Kraus (Orthoptera: Acrididae) were collected from plants growing near the department of Animal Biology (University CHEIKH A. DIOP, Dakar- Senegal). They were killed and placed for 24 hours on soil collected from the botanical garden to allow saprophytic fungi attack. Afterward, they were placed in Petri dishes containing 20 mg of wheat flour (locally purchased and sterilized in the autoclave for about 15 min at 120°C) mixed with 15 ml of sterile distilled water. One Petri dish containing the same medium only was used as a control. They were maintained at 26°C average temperature and in the range of 80– 85% ambient RH. Four days later, 4 fungal isolates appeared in the plates containing the insects, but not on the control. These fungi were separately cultivated in Petri dishes on the same medium. Dry conidia were harvested from the surface of the medium directly by scraping and conserved in Pyrex bottles sterilized at 120°C.

Fungi were identified according to Rapper and Fennel (1965), Samson (1979) and Guarro et al. (1999). The fungus *A. clavatus* was easily identified by his long phototrophic conidiophores (Yaguchi et al., 1993) on microscopic examination during germination.

Preliminary fungal tests on the mosquito larvae (Bioassay 1)

In September 2005, larvae of *Aedes aegypti* L., *Anopheles gambiae* s.l. Giles and *Culex quinquefasciatus* Say (Diptera: Culicidae) were collected from various vats containing rainwater. For each fungal isolate, 1.2 mg/ml dry conidia were applied to 25 larvae (3rd and 4th instar) in 9 X 1.5 cm Petri dishes sterilized at 120°C and containing sterile 25 ml of tap water. There were four replicates for each treatment. Four non-treated Petri dishes served as control. The more virulent *A. clavatus* fungus was selected for production and application.

Microscopic observations

Larvae of *Ae. aegypti* and *An. gambiae* treated with *A. clavatus* were fixed after dying, sectioned, mounted, and then observed under light microscopy. The larvae of *Cx. quinquefasciatus* treated with *A. clavatus* at 1.2 mg/ml were used for cuticle observations under light microscopy. Larvae treated with three drops of the aqueous spore solution (1.2 mg/ml), were incubated on wheat powder medium (for *Ae. aegypti*) and on wet filter paper (for *Cx. quinquefasciatus*) at 85% R.H. and 26°C followed by observation of fungal germination.

Fungal production

A. clavatus was grown in Petri dishes containing 20 g of wheat powder (sterilized for 15 min at 120°C) and 15 ml of sterile distilled water. After four days incubation at 26°C, the substrate and dry conidia content were mixed to obtain a powder (conidia - wheat powder mixture). The number of conidia was determined using a haemocytometer.

Application of fungus (Bioassay 2)

Ae. aegypti and *Cx. quinquefasciatus* mosquito larvae were collected from vats containing rainwater in late November 2005. *An. gambiae* larvae were scarce at this time and were not used for the bioassay. Six plastic bottles (10 x 10 x 7 cm), each containing 500 ml of sterile tap water and 50 larvae (L3 and L4), were used for each mosquito species. Larvae were then treated with *A. clavatus* conidial mixture at 4, 8, 12, 16 and 20 g/l. The corresponding concentrations were 4.3; 8.5; 13; 17 and 21 x 10⁷ spores/ml respectively. Culture medium (wheat powder only) at 0.02 mg/ml served as control for each treatment. Total mortality was recorded for all replicates of each treatment at 24 hours post-inoculation. The surviving larvae were reared in plastic boxes (10 x 10 x 7 cm) containing 500 ml of tap water for 7 days. The emerging adults were incubated for 24 h at 26°C and fungal germination was observed after microscopic examination.

Data analysis

Data on mortality were corrected with Abbott's formula (Abbott, 1925). Student's t-test was used to compare mortality for *Ae. aegypti* and *Cx. quinquefasciatus*.




Figure 1. Larvae of *Culex quinquefasciatus* infected by *Aspergillus clavatus* via cuticle (1a and 1b). x 400. c = cuticle, ger sp = germinating spore, myc = mycelium, sp = spore.

Results

Four *Aspergillus* species were isolated from the dead locusts. Preliminary tests (Bioassay 1) with conidia suspensions of each isolate revealed that *A. clavatus* was highly pathogenic against larvae of *Ae. aegypti*, *Cx. quinquefasciatus* and *An. gambiae*. Mortality rates were 100% against both *Ae. aegypti* and *Cx. quinquefasciatus*, while against *An. gambiae* it was 95%. All rates were in comparison to the control mortality (< 5%) after 24 hours (Table 1). With the other

isolates (S1, S2 and S4), also identified as species in the genus *Aspergillus*, infection against the larvae was less than 5%.

A. clavatus induced significant mortality against mosquito larvae (Table 2) when applied (Bioassay 2 in relative humidity ranging from 65 to 80%, and temperature ranging from 24 to 26°C). The mortality varied from 10.6 to 68% for *Cx. quinquefasciatus* and 36.7 to 100% for *Ae. aegypti*. Larval mortality was significantly higher against

Ae. aegypti than *Cx. quinquefasciatus* ($P = 0.0001$) (Table 2). *A. clavatus* infection was observed under the microscope. On dead larvae, spores were found attached to the cuticle of *Cx. quinquefasciatus* (Figure 1a). Germinating spores (Figure 1a) and mycelia (Figure 1b) were found growing on *C. quinquefasciatus* larva.

Table 1. Larval mortality 24 h after fungal test on the mosquito larvae *Aedes aegypti*, *Anopheles gambiae* and *Culex quinquefasciatus* (average of four replicates).

Species	Fungal stocks				Control
	S1	S2	S3	S4	
<i>Aedes aegypti</i>	0	0	100	0	0
<i>Anopheles gambiae</i>	0	0	95	0	0
<i>Culex quinquefasciatus</i>	0	4	100	0	0

Histological studies revealed that the gastric caeca of some *An. gambiae* was invaded by *A. clavatus* spores (Figure 2a). Gut invasion by conidia and initial stages of germinating conidia were observed on *Ae. aegypti* larvae (Figure 2d). The fungus penetrated the cuticle of *Ae. aegypti* larvae. Conidial germination was also observed on *Cx. quinquefasciatus* larvae incubated on wet filter paper and larvae of *Ae. aegypti* incubated on sterile wheat flour. Fungal growth was observed on all treated and incubated larvae.

Occasionally, a low percentage of *Cx. quinquefasciatus* larvae treated with *A. clavatus* conidia were able to pupate and produce adults. Germinating conidia was observed on the tip of adult abdomen 24h after incubation.

Discussion

A. clavatus was more virulent to the mosquito larvae than the other three fungal isolates. Laboratory results showed that, *A. clavatus* was highly pathogenic against larvae of *Ae. aegypti*, *An. gambiae* and *Cx. quinquefasciatus*. However,

these mosquito larvae do not have the same susceptibility to the fungus. With the same dry conidial dose, (1.2 mg/ml), death rate was 100% against both *Ae. aegypti* and *Cx. quinquefasciatus* larvae and 95% against *A. gambiae* larvae. Referring to larval species susceptibility, the effect of this fungus is similar to that of *M. anisopliae* (Moniliales) and *Tolytocladium cylindrosporum* (Hypocreales) against *An. stephensi*, *Cx. pipiens* and *Ae. aegypti* larvae (Riba et al., 1986).

When treated with *A. clavatus* spores, mortality was 68 % against *Cx. quinquefasciatus* and 100% against *Ae. aegypti* for 21×10^7 spores/ml. Bisht et al. (1996) found that the fungus *Leptolenia caudata* (Oomycetes) yielded a LD_{100} of 7.10^3 spores/ml against *An. culicifacies* after 7 days. Riba et al. (1986) obtained a LD_{100} in the order of 10^7 spores/ml with a stock of *M. anisopliae* against *Ae. aegypti* larvae within 26 hours. From our results the LD_{100} against *Ae. aegypti* was closer to that of *M. anisopliae* against *Ae. aegypti*. Observations of larvae treated with *A. clavatus* revealed that just after adhesion of conidia on the cuticle, some germinated. The conidial proliferation on the cuticle became more obvious after 48 hours that is similar to previous studies on insects (Brett et al. 2004). Silva et al. (2004a) showed that larvae of mosquito treated with *M. anisopliae* had high amounts of conidia adhering to the colloid chitin with at least 90 % germination after 24 hrs incubation.

High levels of germination occurred on *A. clavatus* on dead *Ae. aegypti* larvae incubated on medium for 48 to 72 hours. Fungal germination was also observed on *Cx. quinquefasciatus* larvae in contact with aqueous solution of *A. clavatus* spores and incubated on wet filter paper, which is in agreement with Silva et al. (2004b).

However, the cuticle does not represent the only way for fungal infection. Other possible routes of invasion for *M. anisopliae* have been identified in mosquitoes via the respiratory siphon or the alimentary canal (Al-Aidroos and Roberts, 1978; Lacey et al., 1988; Silva et al., 2004b). Our histological results revealed a high gastric caeca

Table 2. Percent mortality rates for *Culex quinquefasciatus* and *Aedes aegypti* larvae treated with mixture of *Aspergillus clavatus* (average of four replicates).

When grown in wheat flour, after conidial production by *A. clavatus* the concentration of the homogenous mixture was 10.7×10^9 spores/g.

Species	Dose (10^7 spores/ml)					control
	4.3	8.5	13	17	21	
<i>Culex quinquefasciatus</i>	10.6 ± 0.5	46.3± 0.8	42.4 ± 0.4	65 ± 0.2	68 ± 0.6	0.5 ±0.0
<i>Aedes aegypti</i>	36.7 ±0.9	75.5± 0.4	95.9 ± 0.1	100 ± 0.0	100 ± 0.1	1 ± 0.1

Each value represent mean of four replicates and ± SE.
Statistical analysis with t test shows that the difference in mortality between *Ae. aegypti* and *Cx. quinquefasciatus* larvae was highly significant ($p = 0.001$).




Figure 2. *Anopheles gambiae* gastric caecae (2a) and *Aedes aegypti* gut (2b) after infection by *Aspergillus clavatus*. x 400. c cg = cell of gastric caecum, d cell = disintegrating cells, g cel d = gut cell in disintegration, t m = tissue of muscle.

invasion by *A. clavatus* spores in *An. gambiae* larvae. In the digestive tract for *Ae. aegypti* larvae, it was observed germinating conidia, rupture and disintegrating cells of gut. This has also been reported in previous histological studies (Lacey et al. 1988; Lord and Fukuda 1988; Silva et al. 2004b). According to Crisan (1971) and Lacey et al. (1988), a partial digestion of fungal conidia in the gut may induce a release of toxic substances. Silva et al.

(2004b) revealed that rupture and disintegration of cells in the gut on dead larva might be due to the chitinolytic enzymes or others substances produced by the spores. According to Hajek and St. Leger (1994), aggressiveness of entomopathogenic fungus is related to proteolytic, lipolytic and chitinolytic mechanisms that can act after conidial adhesion on the larval cuticle or after invasion of the gut (Crisan 1971; Lacey et al. 1988; Domnas and

Warner 1991; Silva et al. 2004b). *A. clavatus* produces a number of secondary metabolites as tryptoquivaline and tryptoquivalone (Clardy et al. 1975; Buchi et al. 1977); cytochalasin (Demain et al. 1976; Steyn et al. 1982; Lopez-Diaz and Flannigan 1997) and patulin (Varga et al. 2003). The pathological effects noted on the larvae treated with *A. clavatus* in our experiments might be due to these substances.

Light microscopy observations showed that *A. clavatus* conidia produce germ tubes on *Cx. quinquefasciatus* larval cuticle and germinate. Sweeney (1978) showed that with temperatures higher than 30°C, spores of *Culicinomyces* sp. could adhere to the cuticle and invade the gut of *An. amictus* or that of *Cx. fatigans* larvae. This would explain the speed of *A. clavatus* germination on larvae incubated at temperatures ranging between 24 and 26°C.

Cx. quinquefasciatus larvae treated with *A. clavatus* could pupate and produce adults. The resulting adults were collected seven days later and incubated and displayed fungal germination on their abdominal extremities. There was no fungal germination on adults that resulted from untreated larvae (control). This suggests that adult mosquitoes that result from treated larvae are likely contaminated at a pre-imaginal stage. Such an observation was also reported in previous studies on adult mosquitoes. Indeed *Ae. albopictus* larvae (Laird et al. 1992) and *Ae. aegypti* larvae in contact with fungus such as *Coelomomyces* could pupate and produce infected adults (Lucarotti and Shoulkamy 2000). According to Laird et al. (1992), infection of adult *Ae. albopictus* by the fungus *Coelomomyces stegomyiae* var *stegomyiae* could be mortal.

However, in our study, no mortality was recorded for adults reared during 7 days. Lucarotti (1992) found that, on adult mosquitoes, infection by *C. stegomyiae* targets mainly the ovaries, which may explain the germination of *A. clavatus* on the tip of the abdomen.

The laboratory results show that *A. clavatus* isolated from *O. senegalensis*, is virulent against *Ae. aegypti*, *An. gambiae* and *Cx. quinquefasciatus* larvae and could be developed as a biological control agent against mosquitoes. However, further studies are needed for *A. clavatus* strain optimization, and development of better substrates for mass production and practical use. Characterization and application of toxins on mosquito larvae are needed to better understand their rapid killing effects.

References

Abbott WS. 1925. A method of computing the effectiveness of an insecticide. *Journal of Economic Entomology* 18: 265-267.

- Al-Aidroos K, Roberts DW. 1978. Mutants of *Metarhizium anisopliae* with increased virulence toward mosquito larvae. *Canadian Journal of Genetics and Cytology* 20: 211-219.
- Alves SB, Alves LFA, Lopes RB, Pereira RM, Vieira SA. 2002. Potential of some *Metarhizium anisopliae* isolates for control of *Culex quinquefasciatus* (Dipt. Culicidae). *Journal of Applied Entomology* 126: 504-509.
- Batra CP, Mittal PK, Adak T, Sharma VP. 1998. Efficacy of neem oil-water emulsion against mosquito immatures. *Indian Journal of Malariology* 35: 15-21.
- Bisht GS, Joshi C, Khulbe RD. 1996. Water molds: Potential biological control agents of malaria vector *Anopheles culicifacies*. *Current Science* 70: 393-395.
- Buchi G, Luk KC, Kobbe B, Townsend JM. 1977. Four new mycotoxins from *Aspergillus clavatus* related to tryptoquivaline. *Journal of Organic Chemistry* 42: 244-246.
- Clardy J, Springer JP, Buchi G, Matsuo K, Wrightman R. 1975. Tryptoquivaline and Tryptoquivalone, Two new tremorgenic metabolites of *Aspergillus clavatus*. *Journal of the American Chemical Society* 97: 663-665.
- Crisan AV. 1971. Mechanism responsible for release of toxin by *Metarhizium* spores in mosquito larvae. *Journal of Invertebrate Pathology* 17: 260-264.
- Demain AL, Hunt HA, Malik V, Kobbe B, Hawkins H, Matsuo K, Wogan GN. 1976. Improved procedure for production of cytochalasin E by *Aspergillus clavatus*. *Applied and Environmental Microbiology* 31: 138-140.
- Domnas AJ, Warner SA. 1991. Biochemical activities of entomophagous fungi. *Critical Reviews in Microbiology* 18: 1-13.
- Fillinger U, Knols BG, Becker N. 2003. Efficacy and efficiency of new *Bacillus thuringiensis* var. *Israelensis* and *Bacillus sphaericus* formulations against Afrotropical anophelines in Western Kenya. *Tropical Medicine And International Health* 8: 37- 47.
- Guarro J, Gene J, Stchigel AM. 1999. Developments in Fungal Taxonomy. *Clinical Microbiology Reviews* 12: 454-500.
- Govindarajan M, Jebanesan A, Reetha D. 2005. Larvicidal effect of extracellular secondary metabolites of different fungi against the mosquito, *Culex quinquefasciatus* Say. *Tropical Biomedicine* 22: 1-3.
- Hajek A, St Leger RJ. 1994. Interaction between fungal pathogens and insect hosts. *Annual Review of Entomology* 39: 293-322.
- Hati , Ghosh . 1965. *Aspergillus parasiticus* infection in adult mosquitoes. *Bulletin of the Calcutta School of Tropical Medicine* 13: 18-19.
- Karch S, Coz J. 1983. Histopathologie de *Culex pipiens* Linné (Diptera, Culicidae) soumis à l'activité larvicide de *Bacillus sphaericus* 1593-4. *Entomologie Médicale et Parasitologie* 21: 225-230.
- Koua HK, Han SH, d'Almeida MA. 1998. Histopathology of *Anopheles gambiae* s.l. Giles, 1902 (Diptera, Culicidae) subjected to the larvicidal activity of the aqueous extract of *Persea americana* Miller, 1768 (Lauraceae). *Bulletin of the Exotic Pathology Society* 91: 252-256.

- Lacey CM, Lacey LA, Roberts DR. 1988. Route of invasion and histopathology of *Metarhizium anisopliae* in *Culex quinquefasciatus*. *Journal of Invertebrate Pathology* 52: 108-118.
- Laird M, Mogi M, Sota T. 1992. Northernmost occurrences of the protistan pathogen, *Coelomomyces stegomyiae* var. *Stegomyiae*. *Journal of the American Mosquito Control Association* 8: 430-432.
- Legner EF. 1995. Biological control of Diptera of medical and veterinary importance. *Journal of Vector Ecology* 20: 59-120.
- Lord JC, Fukuda T. 1988. An ultrastructural study of the invasion of *Culex quinquefasciatus* larvae by *Leptoglossa chapmani* (Oomycetes: Saprolegniales). *Mycopathologia* 104: 67-73.
- Lopez-Diaz TM, Flannigan B. 1997. Production of patulin and cytochalasin E by *Aspergillus clavatus* during malting of barley and wheat. *International Journal of Food Microbiology* 35: 129-136.
- Lucarotti CJ. 1992. Invasion of *Aedes aegypti* ovaries by *Coelomomyces stegomyiae*. *Journal of Invertebrate Pathology* 60: 176-184.
- Lucarotti CJ, Shoulkamy MA. 2000. *Coelomomyces stegomyiae* infection in adult female *Aedes aegypti* following the first, second and third host blood meals. *Journal of Invertebrate Pathology* 75: 292-295.
- de Moraes AML, Costa GL, Camargo Barcellos MZ, Oliveira RL, Oliveira PC. 2001. The entomopathogenic potential of *Aspergillus* spp. in mosquitoes vectors of tropical diseases. *Journal of Basic Microbiology* 41: 45-49.
- Nnakumusana ES. 1985. Laboratory infection of mosquito larvae by entomopathogenic fungi with particular reference to *Aspergillus parasiticus* and its effects on fecundity and longevity of mosquitoes exposed to sporadic infections in larval stages. *Current Science* 54: 1221-1228.
- Rapper, Fennel. 1965. *The genus Aspergillus*. Williams & Wilkins.
- Ravindra J, Eapen A, Kar I. 2002. Evaluation of repellent action of neem oil against the filarial vector, *Culex quinquefasciatus* (Diptera: Culicidae). *Indian Journal of Malariology* 39: 13-17.
- Riba G, Keita A, Soares GGJ, Ferron P. 1986. Comparative studies of *Metarhizium anisopliae* and *Tolypocladium cylindrosporium* as pathogens of mosquito larvae. *Journal of American Mosquito Control Association* 2: 469-473.
- Samson R. 1979. A compilation of the aspergilli described since 1965. *Studies in Mycology* 18: 1-38.
- Scholte E-J, Njiru BN, Smallegange RC, Takken W, Knols GJ. 2003. Infection of malaria (*Anopheles gambiae* s.s.) and filariasis (*Culex quinquefasciatus*) vectors with the entomopathogenic fungus *Metarhizium anisopliae*. *Malaria Journal* 2: 29.
- Silva RO, Silva HHG, Luz C. 2004a. Effect of *Metarhizium anisopliae* isolates originating from soil samples of the Central Brazilian Cerrado against *Aedes aegypti* larvae under laboratory conditions. *Revista de Patologia Tropical* 33: 207-216.
- Silva RO, Silva HHG, Ulhoa CJ, Luz C. 2004b. Is there a relationship between N-acetyl-d-glucosamidase activity of *Metarhizium anisopliae* (Metschn.) Sorokin (Hyphomycetes) isolates from peridomestic areas in Central Brazil and larvicidal effect on *Aedes aegypti* (L.) (Diptera, Culicidae)? *Journal of Applied Entomology* 129: 158-164.
- Steyn PS, Van Heerden FR, Rabie CJ. 1982. Cytochalasins E and K, toxic metabolites from *Aspergillus clavatus*. *Journal of the Chemical Society Perkin Transactions I* 1982: 541-544.
- Su X, Zou F, Guo Q, Huang J, Chen TX. 2001. A report on a mosquito-killing fungus, *Pythium carolinianum*. *Fungal Diversity* 7: 129-133.
- Sweeney AW. 1978. The effects of temperature on the mosquito pathogenic fungus *Culicinomyces*. *Australian Journal of Zoology* 26: 47-53.
- Varga J, Rigo K, Molnar J, Toth B, Szencz S, Teren J, Kozakiewicz Z. 2003. Mycotoxin production and evolutionary relationships among species of *Aspergillus* section *Clavati*. *Antonie van Leeuwenhoek* 83: 191-200.
- Wright MS, Raina AK, Lax AR. 2005. A strain of the fungus *Metarhizium anisopliae* for controlling subterranean termites. *Journal of Economic Entomology* 98: 1451-1458.
- Yaguchi T, Someya A, Miyadoh S, Udagawa S. 1993. *Aspergillus ingratus*, a new species in *Aspergillus* section *Clavati*. *Transactions of the Mycological Society of Japan* 34: 305-310.