



Imaging Live Bee Brains using Minimally-Invasive Diagnostic Radioentomology

Authors: Greco, Mark K, Tong, Jenna, Soleimani, Manucher, Bell, Duncan, and Schäfer, Marc O

Source: Journal of Insect Science, 12(89) : 1-7

Published By: Entomological Society of America

URL: <https://doi.org/10.1673/031.012.8901>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.



Imaging live bee brains using minimally-invasive diagnostic radioentomology

Mark K Greco^{1a*}, Jenna Tong^{2b}, Manucher Soleimani^{2c}, Duncan Bell^{4d}, Marc O Schäfer^{3e}

¹Department of Biology and Biochemistry, University of Bath, BA2 7AY, United Kingdom

²INVERT Centre, Department of Electrical and Electronic Engineering, University of Bath, BA2 7AY, United Kingdom

³National Reference Laboratory for Bee Diseases, Friedrich-Loeffler-Institute, Federal Research Institute for Animal Health, Südufer 10, 17493, Greifswald – Insel Riems, Germany

⁴East Anglian Radiography Research, modelling and 3-D printing Group, School of Science, Technology and Health, University Campus Suffolk, Ipswich IP4

Abstract

The sensitivity of the honey bee, *Apis mellifera* L. (Hymenoptera: Apidae), brain volume and density to behavior (plasticity) makes it a great model for exploring the interactions between experience, behavior, and brain structure. Plasticity in the adult bee brain has been demonstrated in previous experiments. This experiment was conducted to identify the potentials and limitations of MicroCT (micro computed tomography) scanning “live” bees as a more comprehensive, non-invasive method for brain morphology and physiology. Bench-top and synchrotron MicroCT were used to scan live bees. For improved tissue differentiation, bees were fed and injected with radiographic contrast. Images of optic lobes, ocelli, antennal lobes, and mushroom bodies were visualized in 2D and 3D rendering modes. Scanning of live bees (for the first time) enabled minimally-invasive imaging of physiological processes such as passage of contrast from gut to haemolymph, and preliminary brain perfusion studies. The use of microCT scanning for studying insects (collectively termed ‘diagnostic radioentomology’, or DR) is increasing. Our results indicate that it is feasible to observe plasticity of the honey bee brain in vivo using diagnostic radioentomology, and that progressive, real-time observations of these changes can be followed in individual live bees. Limitations of live bee scanning, such as movement errors and poor tissue differentiation, were identified; however, there is great potential for in-vivo, non-invasive diagnostic radioentomology imaging of the honey bee for brain morphology and physiology.

Keywords: *Apis mellifera*, microCT, X-ray

Abbreviations: DR, diagnostic radioentomology; **MicroCT**, micro computed tomography

Correspondence: ^a m.k.greco@bath.ac.uk, ^b j.r.tong@bath.ac.uk, ^c M.Soleimani@bath.ac.uk, ^d d.bell@ucs.ac.uk,

^e marc.schaefer@fli.bund.de, * Corresponding author

Editor: Carla Penz was Editor of this paper.

Received: 29 June 2011, **Accepted:** 3 November 2011

Copyright : This is an open access paper. We use the Creative Commons Attribution 3.0 license that permits unrestricted use, provided that the paper is properly attributed.

ISSN: 1536-2442 | Vol. 12, Number 89

Cite this paper as:

Greco MK, Tong J, Soleimani M, Bell D, Schäfer MO. 2012. Imaging live bee brains using minimally-invasive diagnostic radioentomology. *Journal of Insect Science* 12:89. Available online: <http://www.insectscience.org/12.89>

Introduction

The European honey bee, *Apis mellifera* L. (Hymenoptera: Apidae), workers weigh approximately 0.1g. Their brain weighs approximately 0.001 g, has a volume of approximately 1 mm³, and has approximately one million neurons (Ribi et al. 2008). The main parts of the brain are the optic lobes, the antennal lobes, the mushroom bodies, and the central complex. The optic and antennal lobes are responsible for processing vision and olfaction, respectively. The mushroom bodies and the central complex constitute the most important centers for behavior, instinct, and memory (Hourcade et al. 2010). Other parts of the brain include the suboesophageal ganglion, tritocerebrum, and ventral cord. It is thought that complex behavior is based on overarching brain networks superimposed on smaller local networks controlling individual responses. Since simple environmental manipulations can both accelerate and delay brain growth in young bees, and since brain volume is sensitive to behavior throughout life, the honey bee has great potential as a model for exploring the interactions between environment, behavior, and brain structure. Experience related changes in brain structure are believed to be an important part of the memory engram (Kolb and Whishaw 1998; Kim and Diamond 2002; Mohammed et al. 2002; Gerber et al. 2004; Kim et al. 2006; Liston et al. 2006), and understanding the relationships between experience and brain structure is key to understanding the relationships between brain and behavior (Kolb and Whishaw 1998). A worker honey bee's natural behavioral change is associated with conspicuous growth of the mushroom bodies in the brain (Withers et al. 1993; Farris et al. 2001; Ismail et al. 2006). The mushroom body calyx is larger in forager bees than

same-aged nurse bees that have not left the hive (Withers et al. 1993; Farris et al. 2001). This structural change may be part of the memory engram for the many foraging-related and navigational tasks learned by a forager bee (Farris et al. 2001; Fahrbach et al. 2003).

Phenotypic plasticity in the adult bee brain has been demonstrated in previous experiments using various techniques, such as the Cavalieri or computer volume segmentation methods (Gundersen & Jenson 1987; Michel & Cruz-Orive 1988; Withers et al. 1993; Brown et al. 2000; Ribi et al. 2008; Maleszka et al. 2009). In all cases, dead bees were used to collect data, which invariably led to differences among individuals.

Our experiment was conducted to identify limitations and potentials for micro computed tomography (MicroCT) scanning of live bees to be used as a comprehensive, non-invasive method for studying brain plasticity, and for teaching morphology and physiology of the brain.

Materials and Methods

The SYRMEP beamline facilities at the ELETTRA synchrotron in Trieste, and a SCANCO μ CT40 bench-top scanner at the University of Bern, were used to scan the bees. At the beamline, newly emerged, adult bees were scanned once daily over five days to observe differential brain plasticity as a result of asymmetric environmental stimuli. Scans on live bees at the beamline facility were performed using phase contrast with the following parameters: 15keV X-ray energy, 20 cm sample to detector distance, a number of projection (over 180°) of 1800, 9 μ m isotropic voxel size, 0.9 seconds exposure time, 1 hour 48 minutes measurement time.




Figure 1. To enhance brain tissue differentiation, bolus injections of radiographic contrast media were delivered via a 30G needle (a) directly into the haemolymph, between the dorsal abdominal terga, of live bees that were previously secured for scanning (b and c). The 3D rendered brain (d) showed that contrast had perfused into tissue to enable improved structural differentiation. High quality figures are available online.

To enhance tissue differentiation, bolus injections of radiographic contrast media were delivered directly into the haemolymph, between the dorsal abdominal terga, via a 30G needle (Figure 1). For visual comparisons of gross anatomical features, MicroCT scans of an ancient bee trapped in amber were also performed on the benchtop scanner, using absorption techniques. The tube operating conditions consisted of an HV peak set at 45kV, and a 177 μ A current. The rest of the parameters were: high resolution mode (1000 Projections/180°), 2048 \times 2048 pixels image matrix, 10 μ m size isotropic voxel, 3 seconds integration time, 610 total slices, 2 hours and 30 minutes measurement time.

Images and brain volume data (Figure 2) were measured using BeeView volume rendering software (DISECT Systems Ltd).

Results

Gross brain morphology, such as the optic




Figure 2. A 3D volume rendered image of a live honey bee's head capsule showing gross morphological structures such as the optic lobes (OL), antennal lobes (AL), aorta (AO), mushroom body calyxes (MBc), and median ocellus (MO). The compound eyes (CE) are visualized immediately adjacent and lateral to the optic lobes. High quality figures are available online.

lobes, antennal lobes, aorta, mushroom body calyxes, and median ocellus, were visualized in 2D and 3D projections. Brain volume measurements (Figure 2) enabled estimates of plasticity. Scanning of live bees enabled minimally-invasive imaging of physiological processes (for the first time), such as passage of contrast from gut to haemolymph (Figure 3), as well as preliminary brain perfusion and plasticity studies (Figure 4a). The image in Figure 4b shows a similar view to Figure 4a, which was produced by Rybak et al. (2010) using data from two-channel confocal microscopy scans. Comparisons of brain images from live extant bees and the 20 million year old bee *Proplebeia abdita* showed little variation in gross morphological features (Figure 4c).

Discussion

The use of MacroCT and MicroCT imaging for the non-invasive study of insects, collectively termed 'diagnostic radioentomology' (DR), is increasing (Hornschemeyer et al. 2002; Johnson et al. 2004; Hönnickea et al. 2005; Greco et al.




Figure 3. A 3D volume rendered image with BeeView software of a live honey bee showing the three body segments (a) and orthogonal, 2D images (b, c, and d) showing the passage of radiographic contrast from the ventriculus (true stomach) to the haemolymph in the coelom. Images were rendered 1.5 hours after ingestion of contrast. High quality figures are available online.

2005; Greco et al. 2006; Greco et al. 2008; Greco et al. 2009, Greco et al. 2011). Results from this study indicate that it is feasible to observe plasticity of the honey bee brain ‘in vivo’ using DR, and that progressive, real-time observations of these changes can be followed in individual live bees in association with environmental stimuli. Plasticity in the adult bee brain has been demonstrated in previous experiments using various techniques, such as the Cavalieri or computer volume segmentation methods. In all cases previous to this study, dead bees were used. However, the use of ex-vivo samples increases the chances of fundamental errors in correlation data analyses due to inherent differences among individuals. Movement errors were not a major limitation of this study, because it was possible to completely immobilize the head. However, haemolymph flow continued, which caused exposure variations between tomographic slices. The exposure variations were easily corrected by using the intensity averaging function during image reconstruction. The greatest challenge for this study was achieving adequate brain tissue differentiation, and it was clear that although radiographic contrast showed




Figure 4. (a) A 2D axial view of a live honey bee brain showing perfusion of contrast medium (C) into peripheral regions. Arrows indicate areas of higher concentration. At 30 minutes post bolus injection into the haemolymph, the lateral ocelli (LO) and aorta (AO) contained more contrast than the sub oesophageal ganglion (SOG). (b) A comparative 2D axial view from the bee brain atlas (<http://www.neurobiologie.fu-berlin.de/beebrain/Default.html>) that was reconstructed from imaging data from two-channel confocal microscopy scans. (c) An axial view of the head capsule of an ancient stingless bee *Proplebeia abdita* (Greco et al. 2011) trapped in amber. The brain of this 20 million years old bee was particularly well preserved, as evidenced by the optic lobes including the medullae (Me) and lobulae (Lo), antennal lobes (AL), protocerebral lobes (P), and the mushroom bodies (MB). The retinal zone (RT) was also well preserved. High quality figures are available online.

promise for improving tissue visualization, further improvements on reconstruction

algorithms are required to better separate brain structures. Bee brain imaging studies from Ribi et al. 2008 and Rybak et al. 2010 are still of superior quality; however, the results in this experiment demonstrate great potential for in-vivo, non-invasive DR imaging of the honey bee for future research in brain plasticity, and for teaching brain morphology and physiology.

Acknowledgements

The authors would like to thank Giuliana Tromba, Lucia Mancini, and Nicola Sodini for their contribution to data rendering. We are grateful to DISECT Systems Ltd for donating their 3D rendering and telelinking software for this study.

References

- Brown SM, Napper RM, Mercer AR. 2000. Analysis of Structural Plasticity in the Honey Bee Brain using the Cavalieri Estimator of Volume and the Disector Method. *Image Analysis and Stereology* 19: 139-144.
- Fahrback SE, Farris SM, Sullivan JP, Robinson GE. 2003. Limits on volume changes in the mushroom bodies of the honey bee brain. *Journal of Neurobiology* 57:141–151.
- Farris SM, Robinson GE, Fahrback SE. 2001. Experience-and age related outgrowth of the intrinsic neurons in the mushroom bodies of the adult worker honey bee. *Journal of Neuroscience* 21(16):6395–6404.
- Gerber B, Tanimoto H, Heisenberg M. 2004. An engram found?—evaluating the evidence from fruit Xies. *Current Opinion in Neurobiology* 14:737–744.
- Greco M, Spooner-Hart R, Holford P. 2005. A new technique for monitoring *Trigona carbonaria* nest contents, brood and activity using X-ray computerized tomography. *Journal of Apicultural Research* 44(3): 97–100.
- Greco M, Bell M, Spooner-Hart R, Holford P. 2006. X-ray computerized tomography as a new method for monitoring *Amegilla holmesi* nest structures, nesting behaviour, and adult female activity. *Entomologia Experimentalis et Applicata* 120: 71–76.
- Greco M, Jones A, Spooner-Hart R, Holford P. 2008. X-ray computerised microtomography (MicroCT): a new technique for assessing external and internal morphology of bees. *Journal of Apicultural Research and Bee World* 47(4): 286–291.
- Greco MK, Hoffmann D, Dollin A, Duncan M, Spooner-Hart R, Neumann P. 2009. The alternative Pharaoh Approach: Stingless bees encapsulate beetle parasites alive. *Naturwissenschaften* 97(3): 319-323.
- Greco MK, Spooner-Hart RN, Beattie GAC, Barchia I, Holford P. 2011. Australian stingless bees improve greenhouse Capsicum production. *Journal of Apicultural Research* 50(2): 102-115.
- Greco MK, Welz PM, Siegrist M, Ferguson SJ, Gallmann P, Roubik DW, Engel MS. 2011. Describing an ancient bee trapped in amber using Diagnostic Radioentomology. *Insectes Sociaux* 58(4): 487-494.
- Gundersen HJG, Jensen EB. 1987. The efficiency of systematic sampling in stereology and its prediction. *Journal of Microscopy* 147: 229–263.

- Hönnicke MG, Foersterb LA, Navarro-Silvab MA, Menke R-H, Rigond L, Cusatisa C. 2005. Preliminary studies of enhanced contrast radiography in anatomy and embryology of insects with Elettra synchrotron light. *Nuclear Instruments and Methods in Physics Research* 548: 207–212.
- Hörschemeyer T, Beutel RG, Pasop F. 2002. Head Structures of *Priacma serrata* Leconte (Coleoptera, Archostemata) Inferred From X-ray Tomography. *Journal of Morphology* 252: 298–314.
- Hourcade B, Muenz TS, Sandoz J-C, Rössler W, Devaud J-M. 2010. The Long-Term Memory Leads to Synaptic Reorganization in the Mushroom Bodies: A Memory Trace in the Insect Brain? *Journal of Neuroscience* 30(18): 6465–6461.
- Ismail N, Robinson GE, Fahrbach SE. 2006. Stimulation of muscarinic receptors mimics experience-dependent plasticity in the honey bee brain. *Proceedings of the National Academy of Sciences of the United States of America* 103:207–211.
- Kim JJ, Diamond DM. 2002. The stressed hippocampus, synaptic plasticity and lost memories. *Nature Reviews Neuroscience* 3:453–462.
- Kim JJ, Song EY, Kosten TA. 2006. Stress effects in the hippocampus: synaptic plasticity and memory. *Stress* 9:1–11.
- Kolb B, Whishaw IQ. 1998. Brain plasticity and behaviour. *Annual Review of Psychology* 49:43–64.
- Johnson SN, Read DB, Gregory PJ. 2004. Tracking larval insect movement within soil using high resolution X-ray microtomography. *Ecological Entomology* 29(1): 117–122.
- Liston C, Miller MM, Goldwater DS, Radley JJ, Rocher AB, Hof PR, Morrison JH, McEwen BS. 2006. Stress-induced alterations in prefrontal cortical dendritic morphology predict selective impairments in perceptual attentional set-shifting. *Journal of Neuroscience* 26(30):7870–4.
- Maleszka J, Barron AB, Helliwell PG, Maleszka R. 2009. Effect of age, behaviour and social environment on honey bee brain plasticity. *Journal of Comparative Physiology A Neuroethology Sensory Neural And Behavioral Physiology* 195:733–740.
- Michel RP, Cruz-Orive LM. 1988. Application of Cavalieri principle and vertical sections method to lung: estimation of volume and pleural surface area. *Journal of Microscopy* 150:117–136.
- Mohammed AH, Zhu SW, Darmopil S, Hjerling-LeZer J, Ernfors P, Winblad B, Diamond MC, Eriksson PS, Bogdanovic N. 2002. Environmental enrichment and the brain. *Progress in Brain Research* 138:109–133.
- Ribi W, Sendenb TJ, Sakellariou A, Limayec A, Zhang S. 2008. Imaging honey bee brain anatomy with micro-X-ray-computed tomography. *Journal of Neuroscience Methods* 171: 93–97.
- Rybak J, Kuß A, Lamecker H, Zachow S, Hege H-C, Lienhard M, Singer J, Neubert K, Menzel R. 2010. The digital bee brain: integrating and managing neurons in a common 3D reference system. *Frontiers in Systems Neuroscience* 4:30.

Withers GS, Fahrbach SE, Robinson GE.
1993. Selective neuroanatomical plasticity and
division of labour in the honey bee. *Nature*
364:238–240.