

## **Molecular Identification and Expressive Characterization of an Olfactory Co-Receptor Gene in the Asian Honeybee, *Apis cerana cerana***

Authors: Zhao, Huiting, Gao, Pengfei, Zhang, Chunxiang, Ma, Weihua, and Jiang, Yusuo

Source: Journal of Insect Science, 13(80) : 1-14

Published By: Entomological Society of America

URL: <https://doi.org/10.1673/031.013.8001>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.



## Molecular identification and expressive characterization of an olfactory co-receptor gene in the Asian honeybee, *Apis cerana cerana*

Huiting Zhao<sup>1a</sup>, Pengfei Gao<sup>1b</sup>, Chunxiang Zhang<sup>1c</sup>, Weihua Ma<sup>1,2d</sup>, Yusuo Jiang<sup>1e\*</sup>

<sup>1</sup>College of Animal Science and Technology, Shanxi Agricultural University, Taigu Shanxi, 030801, China

<sup>2</sup>Institute of Horticulture, Shanxi Academy of Agricultural Sciences, Taiyuan Shanxi, 030031, China

### Abstract

Olfaction recognition process is extraordinarily complex in insects, and the olfactory receptors play an important function in the process. In this paper, a highly conserved olfactory co-receptor gene, *AcerOr2* (ortholog to the *Drosophila melanogaster Or83b*), cloned from the antennae of the Asian honeybee, *Apis cerana cerana* Fabricius (Hymenoptera: Apidae), using reverse transcriptase PCR and rapid amplification of cDNA ends. The full-length sequence of the gene was 1763 bp long, and the cDNA open reading frame encoded 478 amino acid residues, including 7 putative transmembrane domains. Alignment analysis revealed that *AcerOr2* shares high homology (> 74%) with similar olfactory receptors found in other Hymenoptera species. The amino acid identity with the closely related species *Apis mellifera* reached 99.8%. The developmental expression analysis using quantitative real-time reverse transcriptase PCR suggested that the *AcerOr2* transcript was expressed at a relatively low level in the larval stage, whereas it was expressed broadly in the pupal and adult stages, with a significantly high level on the days just before and after eclosion. *In situ* hybridization showed that *AcerOr2* mRNA was expressed in sensilla placodea and on the basal region of the worker antennal cuticle, in accordance with the previous conclusions that the conserved genes are expressed in most olfactory receptor neurons.

**Keywords:** gene expression, location analysis, molecular cloning, olfactory receptor

**Abbreviations:** GSP, gene specific primer; OR, olfactory receptor; RACE, rapid amplification of cDNA ends; qRT-PCR, real time quantitative reverse transcriptase PCR; RT-PCR, reverse transcriptase PCR; TM, transmembrane

**Correspondence:** <sup>a</sup> [zhaohuiting@yahoo.com.cn](mailto:zhaohuiting@yahoo.com.cn), <sup>b</sup> [gpf800411@126.com](mailto:gpf800411@126.com), <sup>c</sup> [zhchx66@126.com](mailto:zhchx66@126.com), <sup>d</sup> [mwhuachina@yahoo.com.cn](mailto:mwhuachina@yahoo.com.cn), <sup>e</sup> [jiangys-001@163.com](mailto:jiangys-001@163.com), \*Corresponding author

**Editor:** Yong Zhang was editor of this paper.

**Received:** 20 August 2012 **Accepted:** 22 January 2013

**Copyright:** This is an open access paper. We use the Creative Commons Attribution 3.0 license that permits unrestricted use, provided that the paper is properly attributed.

**ISSN:** 1536-2442 | Vol. 13, Number 80

#### Cite this paper as:

Zhao H, Gao P, Zhang C, Ma W, Jiang Y. 2013. Molecular identification and expressive characterization of an olfactory co-receptor gene in the Asian honeybee, *Apis cerana cerana*. *Journal of Insect Science* 13:80. Available online: <http://www.insectscience.org/13.80>

## Introduction

Insect olfaction plays a vital role in various life processes, such as mating, ovipositing, avoiding predators, searching for food, and communicating with other members of the species. In insect antennae, the olfactory receptors (ORs) are expressed in many dendritic membranes of the olfactory sensory neurons, participating in the olfaction detection. Seven transmembrane (TM) domain receptor proteins transform the chemical signal to stimulate the neurons. Insect ORs adopt a reverse membrane of G protein-coupled receptors different from the topology structure in mammals (Benton et al. 2006; Ludin et al. 2007).

Olfactory receptors in insects are highly diverse except for the Or83b family. The family shares a highly conserved gene sequence with different insects and plays an important role in regulating insect behavior (Gao and Chess 1999; Larsson et al. 2004; Jones et al. 2005; Wicher et al. 2008). In *Drosophila*, Or83b forms a heterodimer with other ORs and tends to act as a co-receptor (Neuhaus et al. 2005; Benton et al. 2006). *Or83b* mutant flies did not show odorant-evoked action potential with slight spontaneous activity (Larsson et al. 2004; Tsitoura et al. 2010). Hence, the Or83b family plays a crucial role in detecting odorants or pheromones. Recently, *Or83b* was renamed *Orco*, the acronym for olfactory receptor co-receptor (Vosshall and Hansson 2011). The conservative genes from species of Diptera, Lepidoptera, Hymenoptera, and Coleoptera have been identified based on their homology (Hill et al. 2002; Krieger et al. 2003; Melo et al. 2004; Xia and Zwiebel 2006; Malpel et al. 2008; Jordan et al. 2009; Shen et al. 2011).

*AmelR2*, the *DOr83b* orthologue gene in *Apis mellifera*, has been cloned and located (Krieger et al. 2003); however, the expression pattern of *AmelR2* has not yet been fully described in detail. The Asian honeybee, *Apis cerana cerana* Fabricius (Hymenoptera: Apidae), is an indigenous honeybee species in China. In our study, we report the molecular and expressive characterization of *AcerOr2*, a new *Orco* gene in *A. cerana*, to provide a basis for further research on the function of *AcerOr2* in *A. c. cerana*.

## Materials and Methods

### Samples

Bees were reared at the apiary of Shanxi Agriculture University, Shanxi, China. The thorax and the antennae of adult worker bees were collected for separate PCR analyses of the corresponding DNA and RNA sequences. Tissues obtained from different stages, larval (6 whole bodies), pupal (6 heads), and adult (30 pairs of antennae), were dissected and used directly in isolating the total RNA for the expression profiling of worker bees at different ages.

### DNA and RNA extraction

Total DNA was extracted from the thorax of each individual using a modified method described by Smith and Hagen (1996). Total RNA was isolated from 50 antennae of adult worker bees using Trizol reagent (Invitrogen, [www.invitrogen.com](http://www.invitrogen.com)), and stored at -70° C until they were used.

### Genome internal amplification

cDNA was synthesized from the total RNA isolated from the antennae using a Primer-Script RT reagent Kit (TaKaRa, [www.takara-bio.com](http://www.takara-bio.com)). Specific primers were designed based on the sequences closely related to the gene *AmelOr2* (DQ449670) for *AcerOr2*

DNA and cDNA amplification. DNA primers were: Forward1: 5' - TCACCATGCTCTTCTTCACG- 3', Reverse1: 5' - CGCTGAATTCCATCAAAGGC- 3'; and F2: 5' - TGCTCGTGGCTCCTGTTCGC- 3', R2: 5' - AGCAGTTGGCCGGAAGGTGG- 3'. Samples were subjected to pre-denaturation for 4 min at 94° C first, then 30 cycles of amplification were performed at 94° C for 30 sec, 55° C for 32 sec, and 72° C for 1 min, followed by incubation at 72° C for 8 min. The cDNA primers were: F: 5' - AAGACGTGGACGATCTCACC- 3' and R: 5' -GCTACACCATAGGCGTCTCC- 3'). PCR reactions were performed as follows: after 3 min at 94° C, 35 cycles for 30 sec at 94° C, 1 min at 56° C and 1 min at 72° C, then an 8 min elongation at 72° C. PCR products were analyzed by 1.5% agarose gel electrophoresis.

### Rapid amplification of cDNA ends (RACE) amplification

cDNAs were synthesized using the SMART RACE cDNA amplification kit (Clontech, [www.clontech.com](http://www.clontech.com)) for the 3' and 5' RACE. The 3' and 5' regions of the cDNAs were obtained using 3' and 5' RACE-PCRs following the manufacturer's instructions. Gene-specific primers (GSPs) were designed based on the sequence of internal PCR products. 5' 'RACE GSP: 5' - CCCTCCAGCTCCTCGCAGAGTCATGCC G- 3'; 3' RACE GSP: 5' - GCCAATGGTATGATGGCTCCGAGGAAG CC- 3'. Amplification reactions were carried out as follows: 94° C for 3 min; then 33 cycles at 94° C for 30 sec, 58~60° C for 40 sec, and 72° C for 1 min, and 72° C for 6 min.

### Cloning and sequencing

The PCR products were cloned and sequenced in Huada Gene Research Center (Beijing,

China). The overlapping sequences obtained from two internal amplification fragments of DNA were combined to produce the *AcerOr2* DNA; similarly, the internal amplification segment of cDNA and the 3', 5' RACE products were combined to produce the *AcerOr2* mRNA. Homology searches were conducted on the NCBI platform ([www.ncbi.nlm.nih.gov](http://www.ncbi.nlm.nih.gov)). Alignments of multiple sequences were carried out using ClustalW (Thompson et al. 1994). The gene phylogenetic tree was built on Mega 4.0 software ([www.megasoftware.net](http://www.megasoftware.net)) according to neighbor-joining algorithms (Tamura et al. 2007).

### Developmental expression by real-time quantitative PCR (qRT-PCR)

The total RNA of worker bees, larvae (2, 4, and 6 days old), pupae (5 and 10 days old), and adults (1, 5, 10, 15, 20, 25, 30, 35, and 40 days old) for expression profiling were extracted. The corresponding partial cDNA sequences were amplified with the primers designed on the complete sequence of *AcerOr2* (F: 5' - GGATCAGAGGAGGCCAAAC- 3', R: 5' - CCAACACCGAAGCAAAGAGA- 3'). *RPS18* was used as a housekeeping gene based on *A. mellifera RPS18* (Scharlaken et al. 2008) (F: 5' - GATTCCCGATTGGTTTTTGA- 3'; R: 5' - CCCAATAATGACGCAAACCT- 3'). qRT-PCR was run on Mx3000P real-time PCR system (Stratagene, [www.stratagene.com](http://www.stratagene.com)) using the SYBR Premix Ex Taq kit (Takara). All samples were tested in triplicate. The reactions were incubated in a 96-well plate at 95° C for 20 sec, followed by 45 cycles at 95° C for 15 sec and at 60° C for 20 sec. A dissociation curve was created with the thermal profile at 95° C for 30 sec, 60° C for 30 sec, and 95° C for 30 sec. The  $2^{-\Delta\Delta C_t}$  method was used to

measure the relative expression levels of the samples (Livak and Schmittgen 2001).

### mRNA in situ hybridization

Three digoxigenin-labelled RNA antisense oligonucleotide probes were synthesized by Boster ([www.bosterbio.com](http://www.bosterbio.com)) according to the cDNA full-length of *AcerOr2*, which was acquired in the study. The sequence of the antisense probes was as follows:

1. 5' - TTGCGCGCTATGGTAGTCTATCGGCAA  
TTCTTCTA - 3'
2. 5' - AAGCGAACGCTCGTCTAGGTCGTAGAC  
TTCTTGTA - 3'
3. 5' - ATGGTTCGTTGTTTCTATGTGCGTCATC  
TGTGCAT - 3'

Antennae from 1- to 3-day-old adult workers were dissected, embedded in Tissue-Tec OCT Compound (Sakura, [www.sakura.eu](http://www.sakura.eu)). Cryosections (8  $\mu$ m) were prepared at -22° C and mounted on poly-L-lysine treated slides. Samples were air-dried for 5 min before being fixed in 4% paraformaldehyde/0.1M phosphate-buffered saline (PBS) with 0.1% diethyl pyrocarbonate. Then, the slides were washed 3 times with double distilled water and air-dried for about 30 min. The hybridization protocol was performed according to the RISH kit instructions (Boster). In the process, slides were prehybridized for 3 hr. Then, hybridization solution was added to the tissue section, and each slide was covered with a coverslip. Slides were hybridized for 16 hr with the digoxigenin-labeled antisense probes at 42° C in a humid box with 20% glycerol (double distilled water was substituted for the antisense probes as a negative control). After hybridization, the coverslip was removed, and slides were washed twice in 2×SSC for 5 min

at 37° C, and once in 0.5×SSC and 0.2×SSC for 15 min at 37° C respectively. Then, they were treated with blocking solution for 30 min at 37° C and washed 4 times in 1×PBS for 5 min. Slides were incubated in biotinylation mouse anti-digoxigenin for 1 hr at 37° C and then in Strept Avidin-Biotin Complex-alkaline phosphatase for 30 min at room temperature. After being washed 4 times in 1×PBS for 5 min, the hybridization signals were visualized using nitroblue tetrazolium and 5-brom-4-chlor-3-indolyl. Finally, the sections were viewed by using a light microscope (Olympus BX53, [www.olympus-global.com](http://www.olympus-global.com)) and were imaged using Image-Pro Plus 7.0 software (Media Cybernetics, [www.mediacy.com](http://www.mediacy.com)).

## Results

### Characterization and analysis of the partial DNA fragments

*AcerOr2*, a fragment of 1138 bp, was obtained by splicing the 2 internal specific overlapping PCR sequences. The fragment contained 3 exons and 2 introns, with 58.9% A+T nucleotides and 41.1% G+C nucleotides.

Alignment with the other 2 homologous DNA sequences (BLAST in NCBI) revealed that *AcerOr2* exhibits higher identity with *AmeIOr2* (80.4%) than with *BterOr2* (72.6%). Respectively, *AmelOr2* and *BterOr2* exhibit 96%/64.8% and 83%/62.2% similarity with the exons/introns of *AcerOr2*. The divergence was mainly attributed to the difference in length of the introns and the nucleotide variation. The alignment sequences and the ideograph of the 3 genes are shown in Figure 1.

### Characterization and analysis of the full-length cDNA



Using RT-PCR and RACE methods, identified the full-length cDNA sequence of *AcerOR2* was identified (GenBank accession number: JN792581) (Figure 2). Sequence analysis showed that the open reading frame was 1437 bp long and encodes a protein comprising 478 amino acids with a molecular weight of 53.81 kDa. The 5' and 3' non-coding regions contained 116 bp and 210 bp, respectively. *AcerOR2* was predicted to contain 7 TM topology structures of ORs (determined by TMHMM 2.0, Krogh et al. 2001).

The alignment of amino acid sequence of *AcerOr2* with *DmelOr83b* and other orthologues of Hymenoptera (Figure 3) indicated that *AcerOr2* shared high levels of conservation with other Hymenoptera *Orco* genes (> 74% identity). It showed a considerable similarity to *AmelOr2* (99.8% identity) and a relatively low similarity to *DmelOr83b* (64.9%). Isogeny in Hymenoptera was similar with that in Lepidoptera (Xiu et al. 2010). The alignment also revealed a highly conserved region in the C-terminus, particularly between TMs VI and VII, which is remarkably identical in all of the species. To investigate and confirm the evolutionary relationships among these alignment species, a phylogenetic tree was constructed using the neighbor-joining method (Figure 4). The tree showed that the selected Hymenoptera insects were grouped into 4 main clusters or families, namely Apidae, Formicidae, Braconidae, and Chalcididae.

### Developmental expression pattern in worker bees

The expression pattern of *AcerOr2* transcripts was determined by qRT-PCR on different developmental stages in worker bees (Figure 5). qRT-PCR results showed that *AcerOr2* transcripts were expressed broadly in adults. The expressions were expressed only slightly in

larvae, and as the developmental stages progressed, the expression level increased. Notably, the expression reached the maximum level on the day prior to eclosion, revealing that the ability for olfactory recognition enhanced sharply. Furthermore, there was a relatively high level of expression on the first day after eclosion.

### Location analysis in antennae of worker bees

There are several types of chemosensilla in the antennae of honeybee (Esslen and Kaissling 1976), and the sensilla placodea and sensilla trichoidea may be associated with olfactory sensation (Sliber and Sekhon 1961; Kaissling and Renner 1968). In our study, the expression pattern of *AcerOr2* was investigated in workers' antennae. The positive signals were distributed in most sensilla placodea and on the basal region of the antennal cuticle (Figure 6). The expression location of the mRNA transcripts was similar to that found in *AmelOr2* (Krieger et al. 2003). No signal was detected in the negative control section.

### Discussion

The ORs of insects are highly diverse and share almost no identity with other vertebrates and nematodes (Gao and Chess 1999; Smith 1999). In OR superfamily, the *Orco* family is unique. The family contains the highly conserved OR proteins with 60% to 80% amino acid identity shared among the divergent insect species (Larsson et al. 2004; Patch et al. 2009).

In the current study, *AcerOr2*, a putative OR gene, was identified from the *A. c. cerana*. The *AcerOr2* protein exhibited similar characteristics with other *Or83b* orthologues. For instance, it comprised a predicted 7-TM domain structure and shared a high sequence identity with other *Orco* genes. Moreover, *Ac-*

*erOr2* contained the remarkably conserved motif domain spans TMs VI and VII, which has been demonstrated to be at least part of the interaction domain between the conventional ORs and the co-receptor in *D. melanogaster* (Benton et al. 2006). Thus, we deduce that *AcerOr2* belongs to the *Orco* sub-family.

DOr83b and the corresponding orthologs represent the highly conserved protein family, as observed in our study. However, different species exhibit a distinct variation in the introns, which may be primarily attributed to the insertion or the deletion of a base. Although the insects belong to the same order, comparing the similarity among the insects is difficult because of the variation in their introns. The result indicates that the *Or2* gene in *A. cerana*, perhaps including other members of *Orco* family, is evolving quickly, as showed in the study of the *Or2* gene in *Ceratosolen solmsi* (Lu et al. 2009).

The qRT-PCR results revealed that *AcerOr2* transcripts were expressed in all developmental stages, especially in pupae and adults. The faint expression level was tested in the larvae, as observed in *D. melanogaster* (Kreher et al. 2005), *Culex quinquefasciatus* (Xia and Zwiebel 2006), *Spodoptera littoralis* (Malpel et al. 2008), and *Bactrocera cucurbitae* (Shen et al. 2011), indicating that the ability of olfactory identification may be very weak. Two expression peaks appeared on the day just before and after eclosion. This result might be attributed to the response mechanism to chemical stimuli for the environmental change during eclosion. In the *in situ* experiment, labeling of DIG-labeled antisense probes to *AcerOr2* was observed in the sensilla placodea and numerous cells located near the basement membrane of the antennae. The numerous cells were likely housed in the sensilla

placodea and sensilla trichoidea. The expression pattern of *AcerOr2* was similar to other studies about the *Orco* members, such as *DOr83b*, *AmelR2*, *AgOr7*, and *MsexOr2* (Vosshall et al. 1999; Krieger et al. 2003; Pitts et al. 2004; Malpel et al. 2008; Patch et al. 2009). The broad expression profile of *Orco* family members suggests that these olfactory receptors may play an extensive role in olfactory recognition, such as receiving kinds of pheromones and foraging for nectar, pollen, or other odorants. The labeled sensilla, sensilla placodea and sensilla trichoidea, have been described to be associated with olfactory sensation (Sliber and Sekhon 1961; Kaissling and Renner 1968). So, *AcerOr2* might play a role in odorant perception of *A. cerana cerana*.

Several studies have verified that *Or83b* is necessary olfactory responses in *Drosophila* (Larsson et al. 2004; Benton et al. 2006). It can also be co-expressed with a conventional OR to form a ligand-gated cation channel (Sato et al. 2008; Wicher et al. 2008; Jones et al. 2011). The new signal transduction model of insects induced more interest in the study of the relationship between the co-expressed ORs. As a native Chinese bee species, *A. cerana cerana* appears to exhibit highly sensitive olfactory receptors. Based on the identification of the new Hymenoptera *Orco* genes described in this study, further research on ORs have been developed to elucidate the interacting mechanism of the co-receptors and the superior ability of *A. cerana cerana* in searching for nectar and pollen.

## Acknowledgements

This work was supported by grants from the National Natural Science Foundation of China (31040078 and 31272513).

## References

- Benton R, Sachse S, Michnick SW, Vosshall LB. 2006. A typical membrane topology and heteromeric function of *Drosophila* odorant receptors in vivo. *PLoS Biology* 4(2): 240–257.
- Esslen J, Kaissling KE. 1976. Number and distribution of the sensilla on the antennal flagellum of the honeybees (*Apis mellifera*). *Zoomorphology* 83(3): 227–251.
- Gao Q, Chess A. 1999. Identification of candidate *Drosophila* olfactory receptors from genomic DNA sequence. *Genomics* 60(1): 31–39.
- Hill CA, Fox AN, Pitts RJ, Kent LB, Tan PL, Chrystal MA, Ravchik A, Collins FH, Robertson HM, Zwiebel LJ. 2002. G protein-coupled receptors in *Anopheles gambiae*. *Science* 298(5591): 176–178.
- Jones PL, Pask GM, Rinker DC, Zwiebel LJ. 2011. Functional agonism of insect odorant receptor ion channels. *Proceedings National Academy of Sciences of the USA* 108(21): 8821–8825.
- Jones WD, Nguyen TA, Kloss B, Lee KJ, Vosshall LB. 2005. Functional conservation of an insect odorant receptor gene across 250 million years of evolution. *Current Biology* 15(4): 119–121.
- Jordan MD, Anderson A, Begum D, Carraher C, Authier A, Marshall SDG, Kiely A, Gatehouse LN, Greenwood DR, Christie DL, Kralicer AV, Trowell SC, Newcomb RD. 2009. Odorant receptors from the light brown apple moth (*Epiphyas postvittana*) recognize important volatile compounds produced by plants. *Chemical Senses* 34(5): 383–394.
- Kaissling KE, Renner M. 1968. Antennae Rezeptoren für Queen Substance und Sterozelduft bei der Honigbiene. *Zverg Physiol* 59(4): 357–361.
- Krieger J, Klink O, Mohl C, Raming K, Breer H. 2003. A candidate olfactory receptor subtype highly conserved across different insect orders. *Journal of Comparative Physiology (A)* 189(7): 519–526.
- Krieger J, Raming K, Dewer YME, Bette S, Conzelmann S, Breer H. 2002. A divergent gene family encoding candidate olfactory receptors of the moth *Heliothis virescens*. *European Journal Neuroscience* 16(4): 619–628.
- Krogh A, Larsson B, von Heijne G, Sonnhammer EL. 2001. Predicting transmembrane protein topology with a hidden Markov model: application to complete genomes. *Journal of Molecular Biology* 305(3): 567–580.
- Lacher V. 1964. Elektrophysiologische Untersuchungen an einzelnen Rezeptoren für Geruch, Kohlendioxyd, Luftfeuchtigkeit und Temperatur auf den Antennen der Arbeitsbiene und der Drohne (*Apis mellifica* L.). *Zeitschrift für Vergleichende Physiologie* 48(6): 587–623.
- Larsson MC, Domingos AI, Jones WD, Chiappe ME, Amrein H, Vosshall LB, Amrein H. 2004. Or83b encodes a broadly expressed odorant receptor essential for *Drosophila* olfaction. *Neuron* 43(5): 703–714.
- Livak KJ, Schmittgen TD. 2001. Analysis of relative gene expression data using real-time quantitative PCR and the 2<sup>-ΔΔC<sub>T</sub></sup> Method. *Methods* 25(4): 402–408.



- Lu B, Wang N, Xiao J, Xu Y, Murphy RW, Huang D. 2009. Expression and evolutionary divergence of the non-conventional olfactory receptor in four species of fig wasp associated with one species of fig. *BMC Evolutionary Biology* 9(20): 43.
- Lundin C, Käll L, Kreher SA, Kapp K, Sonnhammer EL, Carlson JR, Heijne G, Nilsson I. 2007. Membrane topology of the *Drosophila* OR83b odorant receptor. *FEBS Letters* 581(29): 5601–5604.
- Malpel S, Merlin C, Francois MC, Jacquin-Joly E. 2008. Molecular identification and characterization of two new Lepidoptera chemoreceptors belonging to the *Drosophila melanogaster* OR83b family. *Insect Molecular Biology* 17(5): 587–596.
- Melo ACA, Rützler M, Pitts RJ, Zwiebel LJ. 2004. Identification of a chemosensory receptor from the yellow fever mosquito, *Aedes aegypti*, that is highly conserved and expressed in olfactory and gustatory organs. *Chemical Senses* 29(5): 403–410.
- Neuhaus EM, Gisselmann G, Zhang W, Dooley R, Stortkuhl K, Hatt H. 2005. Odorant receptor heterodimerization in the olfactory system of *Drosophila melanogaster*. *Nature Neuroscience* 8(1): 15–17.
- Patch HM, Velarde RA, Walden KKO, Robertson HM. 2009. A Candidate Pheromone Receptor and Two Odorant Receptors of the Hawkmoth *Manduca sexta*. *Chemical Senses* 34(4): 305–316.
- Pitts RJ, Fox AN, Zwiebel LJ. 2004. A highly conserved candidate chemoreceptor expressed in both olfactory and gustatory tissues in the malaria vector *Anopheles gambiae*. *Proceedings National Academy of Sciences of the USA* 101(14): 5058–5063.
- Sato K, Pellegrino M, Nakagawa T, Vosshall LB, Touhara K. 2008. Insect olfactory receptors are heteromeric ligand-gated ion channels. *Nature* 452(7190): 1002–1006.
- Scharlaken B, Graaf DC, Goossens K, Brunain M, Peelman LJ, Jacobs F. 2008. Reference gene selection for insect expression studies using quantitative real-time PCR: The head of the honeybee, *Apis mellifera*, after a bacterial challenge. *Journal of Insect Science* 8:33. Available online: <http://www.insectscience.org/8.33/>
- Shen JM, Hu LM, Bin SY, Lin JT. 2011. Cloning and expression profiling of an olfactory receptor gene in *Bactrocera cucurbitae* (Coquillett) (Diptera: Tephritidae). *Acta Entomologica Sinica* 54(3): 265–271.
- Slifer EH, Sekhon SS. 1961. Fine structure of the sense organs on the flagellum of the honey bee, *Apis mellifera* Linnaeus. *Journal of Morphology* 109(3) 351–381.
- Smith DP. 1999. *Drosophila* odor receptors revealed. *Neuron* 22(2): 203–204.
- Smith DR, Hagen RH. 1996. The biogeography of *Apis cerana* as revealed by mitochondrial DNA sequence data. *Journal of the Kansas Entomological Society* 69(4): 294–310.
- Tamura K, Dudley J, Nei M, Kumar S. 2007. MEGA4: Molecular Evolutionary Genetics Analysis (MEGA) software version 4.0. *Molecular Biology Evolution* 24(8): 1596–1599.
- Thompson JD, Higgins DG, Gibson TJ. 1994. CLUSTAL W: improving the sensitivity of

progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Research* 22(22): 4673–4680.


Tsitoura P, Andronopoulou E, Tsikou D, Agalou A, Papakonstantinou MP, Kotzia GA, Labropoulou V, Swevers L, Georgoussi Z, Iatrou K. 2010. Expression and Membrane Topology of *Anopheles gambiae* Odorant Receptors in Lepidopteran Insect Cells. *PloS One* 5(11): e15428.

Vosshall LB, Hansson BS. 2011. A unified nomenclature system for the insect olfactory coreceptor. *Chemical Senses* 36(6): 497–498.

Wicher D, Schfer R, Bauernfeind R, Stensmyr MC, Heller R, Heinemann SH, Hansson BS. 2008. *Drosophila* odorant receptors are both ligand-gated and cyclic-nucleotide-activated cation channels. *Nature* 452(7190): 1007–1011.

Xia Y, Zwiebel LJ. 2006. Identification and characterization of an odorant receptor from the West Nile Virus mosquito, *Culex quinquefasciatus*. *Insect Biochemistry and Molecular Biology* 36(3): 169–176.


Xiu W, Zhang Y, Yang D, Dong S, Liu Y. 2010. Molecular Cloning and cDNA Sequence Analysis of Two New Lepidopteran OR83b Orthologue Chemoreceptors. *Agricultural Science in China* 9(8): 1160–1166.




```
1   TCGCGTTGATTTCGCGGAGGAGGATCTCTAAGTTTCTCTGATCTCTGCGGAGGAGTTGCGTTCTTCTGCGGATTTGAAACCGGTTGGOC
100  CACGTAACCGGCTCAAGATGATGAAGTTCAAGCAACAGGGCTAATCGCGAAGTGTGCGCAATATAAACTTGATGAAAGCAACTGCGCACTTCATGT
      M M K F K Q Q G L I A D L M P N I N L M K A T G H F M
199  TCAATTACTACACTGACAGTTCCAGAAACACATACACAAGATCTACTGTATGTCACCTGCTGATAGTATGATGAGTTGCGATTCTGCGGTATCA
      F N Y Y T D S S T K H I H K I Y C I V H L V L I L M Q F G F C G I
298  ATCTAATGATGAGAGCGAGAGTGAAGATCTCAAGGAAACCATCAACATGCTCTTCTCAAGCAGCGTGTCAAGCTGTTTATTTGCGG
      N L M M E S D D V D D L T A N T I T M L F F T H S V V K L V Y F A
397  TCAGGAGTAAATGTTCTACAGAACCTTGCCATATGGAACAATCGAAGCATGCGCTCTTCTGCGAGAACCGCGGATACATCAGATAGCGG
      V R S K L F Y R T L G I V N N P N S H P L F A E S N A R Y H Q I A
496  TTAAGAAGATGAGGATACTGCTGCGGTTATAGGGAACGAGTGTGTCGCGCATTTCTTGAACCATCAAGTTTCTGCGGAGTGTGAAAA-
      V K K M R I L L L A V I G T T V L S A I S W T T I T F I G D S V K
595  AGGTGATGATGCTGCTCAAGAAACAACTAGTGTGAGATACCAAGTTGATGCTGCGTTCTGCTGAGCGCTTACGACCGCAGCGAGGATGCGG
      K V I D P V T N E T T Y V E I P R L M V R S W Y P Y D P S H G M A
694  ATATTTTAACATTGATATTCATTTTACTGCTGATATCTGATGCGAGAGCGAATCTGCTGAGCGTGTCTTCTGCTGCTGCTGCTGCTGCTGCT
      H I L T L I F Q F Y W L I F C M A D A N L L D V L F C S W L L F A
793  GCGAGCAGATGCGATCTGAAGAACATCATCAAGCTTTGATGAGTTGAGCGCGAGTGTGAGAGCGTGTGCGAACAGTGTGAGAACTGTTCAAG
      C E Q I Q H L K N I M K P L M E F S A T L D T V V P N S G E L F K
892  CTGCGAGTGTGAGAGCAAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAG
      A G S A E Q P K E Q E P L P P V T P P Q C E N M L D M D L R C I Y
991  GCAACAGGAGCGATCTGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAG
      S N R T D F T T T F R P T A C M T F N G C V C P N G L T K K Q E M
1090 TGGTACGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAG
      L V R S A I K Y W V E R H K H I V R L V T A I G D A Y G V A L L L
1189 ATATGTTGACTACTACTATTACGTTAAGTTTCTGCTGCTTACCAAGCAACAAAGATACAGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAG
      H M L T T T I T L T L L A Y Q A T K I H A V D T Y A A S V V G Y L
1288 TGTACTCTCTAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAG
      L Y S L G Q V F M L C I F G N R L I E E S S S V M E A A Y S C H W
1387 ATGATGATGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAGGAG
      Y D G S E E A K T F V Q I V C Q Q C Q K A M S I S G A K F F T V S
1486 TCGATCTCTTCTGCTGCTGCTGCTGCTGCTGCTGCTGCTGCTGCTGCTGCTGCTGCTGCTGCTGCTGCTGCTGCTGCTGCTGCTGCTGCTGCT
      L D L F A S V L G A M V T Y F M V L V Q L K *
1585 ATTGCAAGCATGATGATGATGATGATGATGATGATGATGATGATGATGATGATGATGATGATGATGATGATGATGATGATGATGATGATGATGAT
1684 CCTCGAAGCTGCTCAATGATGATGATGATGATGATGATGATGATGATGATGATGATGATGATGATGATGATGATGATGATGATGATGATGATGATGAT
```

**Figure 2.** The nucleotide and deduced amino acid sequences of *AcerOr2*. The boxed codons represent the start and stop codons. Double-red underline represents the polyadenylation signal. The shaded sequences represent the TM domains. High quality figures are available online.











**Figure 4.** Phylogenetic relationships of the *Orco* family in Hymenoptera. The branch labels are bootstrap values. Bootstrap support values (%) are based on 1,000 replicates. Scale bar is 0.05. High quality figures are available online.



**Figure 5.** *AcerOr2* gene expression in *Apis cerana cerana* antennae. The  $2^{-\Delta\Delta CT}$  method was used to measure the relative expression levels of the samples. The expression ratios were first normalized to the control gene *RPS18*, and then normalized on the 30th day in each trial. Lar2 and Lar4 represent the 2- and 4-day-old uncapped larvae, respectively. Lar6 represents the 6-day-old capped larvae. Pup5 and pup10 represent the 5- and 10-day-old pupae, respectively. adu1 to adu40 represent the days after eclosion. Data are shown as mean  $\pm$  SEM. High quality figures are available online.



**Figure 6.** The distribution of *AcerO2* transcripts in worker antennae of *Apis cerana cerana*. A, B are longitudinal and cross section with DIG-labeled probes to *AcerOr2* ( $\times 40$ ). Positive signals are in dark blue. C, D are controls with no labeling ( $\times 40$ ). sp, sensilla placodea; st, sensilla trichoidea. Scale bar: 20  $\mu$ m. High quality figures are available online.