

## **Nutritional Performance of the Tomato Fruit Borer, *Helicoverpa armigera*, on Different Tomato Cultivars**

Authors: Kouhi, Davoud, Naseri, Bahram, and Golizadeh, Ali

Source: Journal of Insect Science, 14(102) : 1-12

Published By: Entomological Society of America

URL: <https://doi.org/10.1673/031.014.102>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.



## Nutritional performance of the tomato fruit borer, *Helicoverpa armigera*, on different tomato cultivars

Davoud Kouhi<sup>a</sup>, Bahram Naseri<sup>b\*</sup>, and Ali Golizadeh<sup>c</sup>

Department of Plant Protection, Faculty of Agriculture, University of Mohaghegh Ardabili, Ardabil, Iran

### Abstract

The development and cultivation of tomato cultivars that are resistant to the tomato fruit borer, *Helicoverpa armigera* (Hübner) (Lepidoptera: Noctuidae), are very limited in Iran and other parts of the world because of the lack of information about resistant tomato cultivars to minimize the use of insecticides. Therefore, the present study was carried out to identify alternative methods to chemical control. Nutritional performance of the larval stages (fourth, fifth, and sixth instars) of *H. armigera* on fruit of eight tomato cultivars, including SUN 6108 fl, Rio grande UG, Korral, Super strain B, CH falat, Hed rio grande, Cal.JN3, and Super crystal, was studied under laboratory conditions. Fourth instars reared on CH falat and SUN 6108 fl respectively showed the highest and lowest values of approximate digestibility. The highest values of efficiency of conversion of ingested food and efficiency of conversion of digested food of fifth instars were on Super strain B. The relative consumption rate and relative growth rate values of the sixth instars were the highest on Korral. The highest and lowest values of consumption index of sixth instars were on Super strain B and Hed rio grande, respectively. The efficiency of conversion of ingested food and efficiency of conversion of digested food values of whole larval instars were the highest on Hed rio grande and lowest on Rio grande UG. The results of nutritional indices indicated that Rio grande UG is an unsuitable host for *H. armigera*.

**Keywords:** nutritional indices, Noctuidae, food consumption, tomato

**Correspondence:** <sup>a</sup> [davud.kuhi@yahoo.com](mailto:davud.kuhi@yahoo.com), <sup>b</sup> [bnaseri@uma.ac.ir](mailto:bnaseri@uma.ac.ir), <sup>c</sup> [golizadeh@gmail.com](mailto:golizadeh@gmail.com), \*Corresponding author

**Editor:** Henry Hagedorn was editor of this paper.

**Received:** 22 July 2012 **Accepted:** 19 June 2014 **Published:** 1 August 2014

**Copyright:** This is an open access paper. We use the Creative Commons Attribution 3.0 license that permits unrestricted use, provided that the paper is properly attributed.

**ISSN:** 1536-2442 | Vol. 14, Number 102

#### Cite this paper as:

Kouhi D, Naseri B, Golizadeh A. 2014. Nutritional performance of the tomato fruit borer, *Helicoverpa armigera*, on different tomato cultivars. *Journal of Insect Science* 14(102). Available online: <http://www.insectscience.org/14.102>

## Introduction

Tomato (*Lycopersicon esculentum* Mill. (Solanales: Solanaceae)) is one of the most important vegetables grown in many parts of the world because it is a good source of vitamins. During 2010–2011, 163,000 ha were under cultivation in Iran, with a total production of 5,887 million tons of tomatoes, and a yield of 36,189 Kg/ha (Iran Ministry of Agriculture 2011). A wide range of insects attack tomato and are a major limiting factor in its successful cultivation and yield (Ashok Kumar et al. 2009). Tomato is more susceptible to the pests' attack than other vegetable crops, mainly because of its tenderness and softness. It is devastated by an array of pests; however, the major damage is caused by the tomato fruit borer, *Helicoverpa armigera* (Hübner) (Lepidoptera: Noctuidae) (Sajjad et al. 2011). This insect is a highly polyphagous and serious pest that infests more than 100 plant species, including vastly planted, economically important crops such as cotton, maize, tobacco, pigeonpea, and chickpea (Talekar et al. 2006). It causes widespread economic damage to tomato farms in Asia (Srinivasan 1959; Singh and Singh 1975; Vattanatumgum and Ruchtapakornchai 1978; Talekar et al. 1984). Controlling the insect pests with insecticides causes serious side effects, including development of insecticide resistance in the insects, pest resurgence, environmental pollution, and health hazards. The development and cultivation of *H. armigera*-resistant tomato cultivars are very limited in Iran and the world because of the lack of information about resistant tomato cultivars. Therefore, the present study was carried out to identify alternative methods to chemical control.

Different nutritive values of host plants can influence the rate of development of *H. armigera* larvae and so affect the population

dynamics of this pest (Ruan and Wu 2001). In addition, food consumption and use link plants attributes with insect performance (Slansky 1990). Host plant resistance is an important tool in integrated pest management (IPM) (Sajjad et al. 2011).

To determine the potential of different tomato cultivars to manage *H. armigera* populations, data on the effects of various cultivars on nutritional indices of the pest are necessary. In spite of the economic importance of *H. armigera* in tomato farms, to our knowledge, no published information exists on the nutritional indices of this pest on different tomato cultivars. Some related studies, however, have been conducted on the effects of host plants, apart from tomato cultivars, on nutritional indices of *H. armigera* (Ashfaq et al. 2003, Naseri et al. 2010, Arghand et al. 2011, Hemati et al. 2012). The objective of this study was to compare nutritional indices of *H. armigera* larvae on tomato cultivars. The results of this research can be used in IPM programs of *H. armigera* on tomato.

## Materials and Methods

### Tomato sources

Seeds of eight tomato (*L. esculentum*) cultivars, SUN 6108 fl, Rio grande UG, Super crystal, CH falat, Super strain B, Hed rio grande, Korral, and Cal.JN3, were obtained from the Plant and Seed Modification Research Institute, Karaj, Iran. The selected tomato cultivars are the most economically important cultivars grown in Iran and many other parts of the world. They were sown in the research field of the University of Mohaghegh Ardabili, located in Ardabil, Iran, in May 2011. For this study, the leaves of different tomato cultivars were used for feeding first and second larval instars, and the fruits were used for feeding third to sixth larval instars.

### Laboratory colony

*Helicoverpa armigera* larvae were originally collected from tomato fields of the Moghan region in Northwest Iran. The stock culture was initiated on different tomato cultivars in a growth chamber at  $25 \pm 1^\circ\text{C}$ ,  $65 \pm 5\%$  RH, with a 16:8 L:D photoperiod.

### Experiments

Neonate larvae were collected from the stock cultures and divided into five replicates (10 larvae in each) and transferred into plastic containers (18.5 cm diam  $\times$  7.5 cm deep) with a hole covered by a mesh net for aeration; the containers held the fresh leaves of each tomato cultivar. The petioles of detached leaves were covered with water-soaked cotton to maintain freshness. Nutritional indices were determined using fourth to sixth instars because they are more easily measured than the primary instars. The first, second, and third instars were reared in groups until they reached the fourth instar, after which they were separated into individual plastic plates (8 cm diam  $\times$  1 cm deep) to avoid cannibalistic behavior. Sixth instars were kept in plastic tubes (1.5 cm diam  $\times$  5 cm deep) for prepupation and pupation.

The gravimetric method (Waldbauer 1968) was used to determine weight gain, food utilization, and feces produced. Nutritional indices were calculated on the dry weight basis. After measuring the weight of the fourth instars, they were introduced on the fruits of different tomato cultivars; the weights of the larvae were recorded daily before and after feeding until they finished feeding and reached the prepupal stage. The initial fresh fruits and the fruits and feces remaining at the end of each experiment were weighed daily. The quantity of food ingested was determined by subtracting the diet remaining at the end of each experiment from the total weight of fresh diet

supplied. The weight of feces produced by the larvae was recorded daily. To obtain the dry weights of the fruits, feces, and larvae, 20 specimens from each cultivar were weighed, oven-dried (48 hr at  $60^\circ\text{C}$ ), and then reweighed.

The formulas described by Waldbauer (1968) were used to determine nutritional indices; CI (consumption index), AD (approximate digestibility), ECI (efficiency of conversion of ingested food), ECD (efficiency of conversion of digested food), RCR (relative consumption rate), and RGR (relative growth rate):

$$CI = \frac{E}{A}$$

$$AD = \frac{E - F}{E}$$

$$ECI = \frac{P}{E}$$

$$ECD = \frac{P}{E - F}$$

$$RCR = \frac{E}{A * T}$$

$$RGR = \frac{P}{A * T}$$

in which, A = mean dry weight of insect over unit time, E = dry weight of food consumed, F = dry weight of feces produced, P = dry weight gain of insect, and T = duration of feeding period.

### Data analysis

Nutritional indices of *H. armigera* reared on different tomato cultivars were analyzed with one-way ANOVA using the statistical software Minitab 16 (Minitab, [www.minitab.com](http://www.minitab.com)) to determine the similarities or significant differences. Statistical differences among the means were evaluated using the LSD test at  $\alpha = 0.05$ . Before analysis, data were tested for normality.

**Table 1.** Nutritional indices of fourth instar larvae of *Helicoverpa armigera* on different tomato cultivars.

Index (mean ± SE)						
Tomato cultivar	CI	AD (%)	ECI (%)	ECD (%)	RCR (mg/mg/day)	RGR (mg/mg/day)
Sun 6108 fl	20.961 ± 2.76b	93.257 ± 0.968f	11.793 ± 1.48a	12.839 ± 1.67a	6.693 ± 0.994a	0.599 ± 0.052a
Cal.JN3	21.180 ± 1.81b	95.324 ± 0.494de	12.261 ± 1.47a	12.943 ± 1.59a	5.112 ± 0.703da	0.551 ± 0.075a
Korral	20.473 ± 1.26b	95.048 ± 0.537e	10.945 ± 0.796a	11.586 ± 0.870ab	6.323 ± 0.472a	0.627 ± 0.051a
Super crystal	26.155 ± 1.94a	96.193 ± 0.320cd	5.857 ± 0.657b	6.095 ± 0.692c	6.236 ± 0.828a	0.358 ± 0.058b
CH falat	26.393 ± 1.65a	98.731 ± 0.106a	7.143 ± 0.479b	7.238 ± 0.488c	5.740 ± 0.476a	0.387 ± 0.025b
Hed rio grande	16.934 ± 0.738c	97.849 ± 0.176ab	10.592 ± 0.657a	10.835 ± 0.700b	4.210 ± 0.274a	0.429 ± 0.028b
Super strain B	19.754 ± 0.698b	97.003 ± 0.256bc	11.422 ± 0.754a	11.787 ± 0.790ab	5.256 ± 0.277a	0.581 ± 0.032a
Rio grande UG	19.917 ± 0.801b	97.704 ± 0.115b	11.65 ± 1.11a	11.95 ± 1.14ab	5.338 ± 0.347a	0.595 ± 0.041a

The means followed by different letters in the same columns are significantly different (LSD,  $P < 0.01$ ). CI, consumption index; AD, approximate digestibility; ECI, efficiency of conversion of ingested food; ECD, efficiency of conversion of digested food; RCR, relative consumption rate; RGR, relative growth rate.

**Table 2.** Nutritional indices of fifth instar larvae of *Helicoverpa armigera* on different tomato cultivars.

Index (mean ± SE)						
Tomato cultivar	CI	AD (%)	ECI (%)	ECD (%)	RCR (mg/mg/day)	RGR (mg/mg/day)
Sun 6108 fl	6.730 ± 0.300cd	86.510 ± 1.11e	13.743 ± 1.78b	16.169 ± 2.16b	2.150 ± 0.180abc	0.285 ± 0.042bc
Cal.JN3	7.221 ± 0.655c	87.473 ± 1.04e	12.960 ± 0.919b	14.972 ± 1.10bc	1.841 ± 0.218cd	0.222 ± 0.017de
Korral	6.816 ± 0.265c	84.645 ± 0.614f	16.432 ± 1.11a	19.490 ± 1.36a	2.470 ± 0.246a	0.344 ± 0.027a
Super crystal	8.805 ± 0.595a	94.523 ± 0.542b	10.257 ± 1.32c	10.880 ± 1.42de	1.597 ± 0.220d	0.157 ± 0.023f
CH falat	8.482 ± 0.270ab	95.690 ± 0.193ab	12.703 ± 1.35b	13.281 ± 1.42cd	1.699 ± 0.106d	0.202 ± 0.016ef
Hed rio grande	6.167 ± 0.255d	90.927 ± 0.443d	17.070 ± 1.08a	18.818 ± 1.21a	1.581 ± 0.103d	0.261 ± 0.024cd
Super strain B	6.753 ± 0.359cd	92.331 ± 0.513c	18.134 ± 1.51a	19.683 ± 1.65a	1.896 ± 0.123bcd	0.323 ± 0.020ab
Rio grande UG	8.185 ± 0.291b	96.305 ± 0.143a	8.878 ± 0.665c	9.213 ± 0.694e	2.183 ± 0.110ab	0.195 ± 0.021ef

The means followed by different letters in the same columns are significantly different (LSD,  $P < 0.01$ ). CI, consumption index; AD, approximate digestibility; ECI, efficiency of conversion of ingested food; ECD, efficiency of conversion of digested food; RCR, relative consumption rate; RGR, relative growth rate.

A dendrogram of different tomato cultivars based on nutritional indices of whole larval instars (fourth, fifth, and sixth instars) of *H. armigera* on different cultivars of tomato was constructed after cluster analysis by Ward’s method using SPSS 16.0 statistical software (IBM, [www.ibm.com](http://www.ibm.com)).

Results

The results of the nutritional indices of fourth, fifth, and sixth, and whole larval instars of *H. armigera* are provided in Tables 1, 2, 3, and 4. Nutritional indices of fourth instars of *H. armigera*, except RCR, were significantly different on tomato cultivars ( $P < 0.01$ ). The larvae reared on cultivar Cal.JN3 showed the highest values of ECI ( $F = 3.41$ ;  $df = 125$ ;  $P < 0.01$ ) and ECD ( $F = 3.50$ ;  $df = 125$ ;  $P < 0.01$ ). The lowest values of ECI and ECD were on Super crystal. The highest values of AD ( $F = 10.02$ ;  $df = 125$ ;  $P < 0.01$ ) and CI ( $F = 2.94$ ;  $df = 125$ ;  $P < 0.01$ ) were observed on cultivar

CH falat. However, the lowest value of AD was on SUN 6108 fl. The larvae that fed on Korral and Super crystal respectively showed the highest and lowest RGR values ( $F = 3.73$ ;  $df = 125$ ;  $P < 0.01$ ). However, the lowest value of CI was on cultivar Hed rio grande (Table 1). The highest values of RCR ( $F = 2.70$ ;  $df = 125$ ;  $P < 0.05$ ) and RGR ( $F = 4.64$ ;  $df = 125$ ;  $P < 0.01$ ) of fifth instars were observed on Korral. The larvae that fed on cultivars Rio grande UG and Korral had respectively the highest and lowest values of AD ( $F = 36.21$ ;  $df = 125$ ;  $P < 0.01$ ) (Table 2).

The highest and lowest CI values ( $F = 17.10$ ;  $df = 125$ ;  $P < 0.01$ ) of sixth instar *H. armigera* were respectively on cultivars Super strain B and Hed rio grande. The ECI ( $F = 6.15$ ;  $df = 125$ ;  $P < 0.01$ ) and ECD ( $F = 4.53$ ;  $df = 125$ ;  $P < 0.01$ ) of larvae reared on Hed rio grande showed the highest values compared to the other cultivars (Table 3).



**Table 3.** Nutritional indices of sixth instar larvae of *Helicoverpa armigera* on different tomato cultivars.

Index (mean ± SE)						
Tomato cultivar	CI	AD (%)	ECI (%)	ECD (%)	RCR (mg/mg/day)	RGR (mg/mg/day)
Sun 6108 fl	3.942 ± 0.092d	86.770 ± 1.14c	13.251 ± 1.11d	15.255 ± 1.27d	0.600 ± 0.036c	0.074 ± 0.006d
Cal.JN3	4.316 ± 0.132c	81.673 ± 1.77d	14.816 ± 0.894bc	18.790 ± 1.31b	0.494 ± 0.034d	0.072 ± 0.006d
Korral	4.455 ± 0.144bc	81.157 ± 0.923d	12.945 ± 0.803d	16.076 ± 1.00cd	0.837 ± 0.047a	0.105 ± 0.007a
Super crystal	3.874 ± 0.128d	82.243 ± 2.77d	18.461 ± 0.972a	21.393 ± 1.70a	0.448 ± 0.026d	0.082 ± 0.006bcd
CH falat	4.254 ± 0.129c	92.942 ± 0.330a	16.300 ± 1.07b	17.554 ± 1.15bc	0.569 ± 0.023c	0.091 ± 0.005b
Hed rio grande	3.389 ± 0.104e	89.289 ± 0.577b	19.022 ± 0.927a	22.750 ± 1.09a	0.435 ± 0.019d	0.081 ± 0.004bcd
Super strain B	5.337 ± 0.158a	85.380 ± 1.05c	12.911 ± 0.770d	15.232 ± 1.00d	0.686 ± 0.029b	0.087 ± 0.005bc
Rio grande UG	4.460 ± 0.092b	78.365 ± 0.421e	13.952 ± 0.679cd	17.822 ± 0.900bc	0.568 ± 0.024c	0.079 ± 0.005cd

The means followed by different letters in the same columns are significantly different (LSD,  $P < 0.01$ ). CI, consumption index; AD, approximate digestibility; ECI, efficiency of conversion of ingested food; ECD, efficiency of conversion of digested food; RCR, relative consumption rate; RGR, relative growth rate.

**Table 4.** Nutritional indices of whole larval instars of *Helicoverpa armigera* on different tomato cultivars.

Index (mean ± SE)						
Tomato cultivar	CI	AD (%)	ECI (%)	ECD (%)	RCR (mg/mg/day)	RGR (mg/mg/day)
Sun 6108 fl	5.906 ± 0.182d	88.682 ± 0.552d	12.569 ± 0.472c	14.187 ± 0.544c	1.338 ± 0.065b	0.166 ± 0.009c
Cal.JN3	6.425 ± 0.230c	86.591 ± 1.29e	13.441 ± 0.395b	15.709 ± 0.668b	1.092 ± 0.062cd	0.146 ± 0.007d
Korral	6.382 ± 0.195c	80.970 ± 1.17f	13.350 ± 0.295b	16.612 ± 0.468a	1.568 ± 0.065a	0.207 ± 0.008a
Super crystal	6.667 ± 0.154bc	90.592 ± 1.34c	12.197 ± 0.501cd	13.276 ± 0.517d	1.051 ± 0.051de	0.127 ± 0.010e
CH falat	6.754 ± 0.119b	95.336 ± 0.142a	12.609 ± 0.383c	13.227 ± 0.406d	1.167 ± 0.030c	0.145 ± 0.004d
Hed rio grande	5.219 ± 0.096e	92.156 ± 0.230b	15.985 ± 0.361a	17.351 ± 0.405a	0.980 ± 0.031e	0.156 ± 0.005cd
Super strain B	7.065 ± 0.176a	90.51 ± 0.499c	13.978 ± 0.529b	15.475 ± 0.632b	1.391 ± 0.049b	0.191 ± 0.005b
Rio grande UG	7.081 ± 0.106a	94.22 ± 0.191a	11.535 ± 0.225d	12.244 ± 0.246e	1.347 ± 0.042b	0.155 ± 0.006cd

The means followed by different letters in the same columns are significantly different (LSD,  $P < 0.01$ ). CI, consumption index; AD, approximate digestibility; ECI, efficiency of conversion of ingested food; ECD, efficiency of conversion of digested food; RCR, relative consumption rate; RGR, relative growth rate.

The results depicted in Table 4 for whole larval instars showed that the ECI ( $F = 9.49$ ;  $df = 125$ ;  $p < 0.01$ ) and ECD ( $F = 8.02$ ;  $df = 125$ ;  $P < 0.01$ ) values were the highest on Hed rio grande and lowest on Rio grande UG. The highest and lowest AD values ( $F = 29.89$ ;  $df = 125$ ;  $P < 0.01$ ) were respectively on cultivars CH falat and Korral. The larvae that fed on cultivar Korral had the highest RCR ( $F = 12.55$ ;  $df = 125$ ;  $P < 0.01$ ) and RGR ( $F = 10.77$ ;  $df = 125$ ;  $P < 0.01$ ) values. However, the highest and lowest values of CI were on Rio grande UG and Hed rio grande, respectively ( $F = 10.46$ ;  $df = 125$ ;  $p < 0.01$ ).

The results presented in Fig. 1 for whole larval instars showed significant difference for larval weight ( $F = 16.84$ ;  $df = 128$ ;  $P < 0.01$ ), feces produced ( $F = 21.46$ ;  $df = 128$ ;  $P < 0.01$ ), and food consumed ( $F = 45.69$ ;  $df = 128$ ;  $P < 0.01$ ). The highest and lowest values of larval weight were observed on SUN 6108 fl ( $18.984 \pm 0.581$  mg) and Super crystal

( $12.484 \pm 0.725$  mg), respectively. The food consumed and feces produced were respectively highest on CH falat and Korral ( $91.910 \pm 1.68$  and  $12.059 \pm 0.708$  mg, respectively) and lowest on Korral and Rio grande UG ( $63.422 \pm 1.07$  and  $4.395 \pm 0.168$  mg, respectively).

A dendrogram based on nutritional indices of whole larval instars of *H. armigera* reared on different tomato cultivars is shown in Fig. 2. The dendrogram shows two separate clusters labeled A (including subclusters A1 and A2) and B (including subclusters B1 and B2). Different cultivars were grouped within each cluster according to the comparison of the nutritional indices of *H. armigera* fed different tomato cultivars. Cluster A included subclusters A1 (CH falat and Rio grande UG) as a partially resistant group and A2 (Cal.JN3, Super crystal and Sun 6108 fl) as an intermediate group. Cluster B consisted of subclusters B1 (Korral and Super strain B)

and B2 (Hed rio grande) as susceptible cultivars.

## Discussion

Study of tomato cultivars for resistance and susceptibility against tomato fruit borer is an important tool to manage the fruit borer with environmentally safe tactics (Sajjad et al. 2011). Differences in allelochemical concentrations, poor nutritional quality of the food, and pericarp thickness between host plant cultivars can affect an insect's performance as larva and survival and development of insects on different cultivars (Sharma et al. 1982; Samraj and David 1988; Martin and Pulin 2004). In this study, the data of nutritional indices, especially ECI and ECD values, of *H. armigera* reared on different tomato cultivars were significantly different, which suggests that the cultivars had different nutritional value. Among nutritional indices, ECI is an indicative index of an insect's ability to utilize the food ingested for growth and development, and ECD is an index of the efficiency of conversion of digested food into growth (Nathan et al. 2005). Also, change in ECD index indicates the overall increase or decrease of the proportion of digested food metabolized for energy (Naseri et al. 2010).

The data generated for the *H. armigera* fifth and sixth instar larvae are inconsistent with each other because the nutritional requirements of the insect change through development, and such differences typically result in changes of food consumption and utilization (Barton Browne 1995). Usually, when the quantity of food ingested is decreased, the duration of development is extended and the insect becomes smaller and lighter. Another reason may be related to increased instar duration, when increased amount of ingested food must be dedicated to

maintaining metabolism. Nutritional requirements would be positively correlated with the mass of the insect (Phillipson 1981; Schroeder 1981).

The results for the fourth instars showed that the larvae that fed on Cal.JN3 had the highest ECI and ECD values. High ECI and ECD values indicate that the larval feeding and their weight have improved. Consequently, larger pupae are produced, which has a direct correlation to adult fertility, which is ecologically very important for the survival of this insect (Daryaei et al. 2007). Foss and Rieske (2003) suggested that a combination of plant characteristics may be responsible for insect preference and performance, and that an optimal combination of plant components serves to maximize host suitability. Also, as fourth instars, larvae that fed on cultivar CH falat showed the highest AD and almost the lowest ECD. Apparently, the increase in AD value could not compensate for the decrease in ECD value, which consequently led to reduced growth rate. Growth reduction is a general response and reaction of phytophagous insects confronted with a new host plant (Grabstein and Scriber 1982; Sheppard and Friedman 1990; Lazarevic and Peric-Mataruga 2003).

Higher CI value of the whole *H. armigera* larval instars on Rio grande UG indicated that the rate of intake relative to the mean larval weight during the feeding period was the highest on this cultivar. The results for AD value of the whole larval instars reared on CH falat and Rio grande UG were different from those reported by Hemati et al. (2012) on tomato var. Meshkin ( $67.470 \pm 0.016\%$ ). This difference is probably related to difference in tomato cultivars, which are different in acidity and secondary compounds.

Among different tomato cultivars, the highest ECI and ECD values of the whole larval instars were observed on cultivars Hed rio grande and Korral, indicating that they were more efficient at the conversion of ingested and digested food to biomass in larval body. Also, the whole larval instars on Rio grande UG showed the lowest ECD and ECI values, maybe because of a shortage of nutritional components and the existence of some secondary chemicals in this cultivar. The results for ECI values of the larvae that fed on Rio grande UG was nearly similar to those reported by Hemati et al. (2012) on chickpea (var. Arman) ( $11.981 \pm 0.006\%$ ). The mean ECD value of whole larval instars reared on different tomato cultivars was lower than that reported by Naseri et al. (2010) on soybean cultivars ( $60.592 \pm 2.012\%$ ), suggesting that the larvae that fed on fruit of tomato cultivars were apparently not efficient in turning digested food into biomass compared with those that fed on pods of soybean cultivars. This finding related to the unsuitability of tomato agrees with the results of Liu et al. (2004), who reported that tomato was an unsuitable host plant compared with other test hosts.

For the whole larval instars, the larval weight, as an important fitness indicator of insect population dynamics (Liu et al. 2004), was highest on SUN 6108 fl and CH falat. This reinforces the suggestion that SUN 6108 fl and CH falat are more suitable cultivars for *H. armigera* larvae than the others. Also, the larvae reared on cultivar Super crystal were lighter than those reared on the other cultivars. We found that the larval weight of *H. armigera* reared on cultivar Korral was similar to that reported by Hemati et al. (2012) on tomato var. Meshkin ( $15.205 \pm 0.272$  mg) and Naseri et al. (2010) on soybean var. 'L17' ( $15.497 \pm 0.911$  mg).

The duration of developmental period of larval stages is a determining factor that shows whether a cultivar is suitable or unsuitable for feeding larvae, as the lepidopteran larvae fed highly-nutritious food increase growth rates and complete development period faster than those larvae that feed on low-nutrient food (Hwang et al. 2008). For the RGR and RCR values, the duration of feeding period is an effective factor. Among different tomato cultivars, the highest RCR and RGR values of the whole larval instars *H. armigera* were on cultivar Korral and the lowest ones were on Hed rio grande and Super crystal. Our results indicate that Hed rio grande and Super crystal have low nutritive value for larvae of *H. armigera*, and on these cultivars a longer development period was needed to complete immature stages. Conversely, the cultivar Korral was highly nutritive for the larvae, and a shorter period of development was necessary to complete immature stages.

Comparing the results of Tables 1–3 shows that the ECI and ECD values, in most cases, increased from fourth to sixth (ultimate) instar. The trend of increase in ECD from early to late instars has been reported by Slansky and Scriber (1985) and Hemati et al. (2012). Nation (2001) noted that the physiological changes in the penultimate and ultimate instars reared on different host plants probably are responsible for the differences in such decreases in ECD values of the two larval instars.

High ECI and ECD values were frequently associated with low CI and vice versa. A similar result was reported in numerous studies on *H. armigera* and other insects (Barbosa and Greenblatt 1979; Naseri et al. 2010; Arghand et al. 2011; Hemati et al. 2012). Also, the results of whole larval instars showed that there is a negative correlation between RCR and



ECI values. Soo Hoo and Fraenkel (1966) noted that the negative correlations between RCR and ECI could have either of two explanations. First, when larvae consume less, the food tends to pass through their digestive system more slowly, and so it can be converted more completely and used by the insects. Second, it may be that insects consume less of a special food simply because they are capable of converting it more efficiently and therefore do not need to eat large quantities of that food to reach appropriate levels of growth.

The results of the cluster analysis revealed that grouping within each cluster might be due to a high level of physiological similarity of different tomato cultivars. The results of the comparative nutritional indices of *H. armigera* on different tomato cultivars indicated that subcluster A1 was the least suitable, subclusters B1 and B2 were the most suitable cultivars for *H. armigera*, and the cultivars in subcluster A2 had an intermediate status.

The results of this study suggest that cultivars Korral and Super strain B are more nutritive, and cultivars CH falat and Rio grande UG are less nutritive for *H. armigera* larvae compared with the other test cultivars.

Analysis of nutritional indices can lead to the understanding of the behavioral and physiological basis of an insect's response to host plants (Lazarevic and Peric-Mataruga 2003). The lesser suitability of some cultivars as host plants of *H. armigera* may be due to the presence of some secondary phytochemicals in these cultivars or the absence of primary nutrients necessary for growth and development of *H. armigera*. These findings will be helpful in understanding the host preference of this particular pest and could help in its management and control, particularly on tomato. Therefore, future studies should focus on test-

ing a wider range of cultivars of tomato for the nutritional indices of *H. armigera*. Finally, assessment of the chemical components of the test cultivars would be helpful to better understand the mechanism of host suitability.

## Acknowledgements

We thank M. Nemati, F. Rahimi Namin, and H. Abbasi (from University of Mohaghegh Ardabili) for their valuable help in this study. This research was supported by grants from the University of Mohaghegh Ardabili, (Ardabil, Iran), which is appreciated.

## References

- Arghand, A., B. Naseri, J. Razmjou, and M. Hassanpour. 2011.** *Feeding indices of the cotton bollworm, *Helicoverpa armigera* (Hübner) (Lepidoptera: Noctuidae) on seed of five different maize hybrids*. Global Conference on Entomology, 5–9 March 2011, Chiang Mai, Thailand.
- Ashfaq, M., K. J. Ahmad, and A. Ali. 2003.** Morphophysical factors affecting consumption and coefficient of utilization of *Helicoverpa armigera* (Hübner). *Pak. J. Appl. Sci.* 3: 225–230.
- Ashok Kumar, C. T., and C. Shivaraju. 2009.** Bioefficacy of newer insecticide molecules against tomato fruit borer, *Helicoverpa armigera* (Hübner). *J. Agric. Sci.* 22: 288–289.
- Barbosa, P., and J. Greenblatt. 1979.** Suitability, digestibility and assimilation of various host plants of the gypsy moth *Lymantria dispar* (L.). *Oecologia* 43: 111–119.
- Barton Browne, L. 1995.** Ontogenetic changes in feeding behavior, pp. 307–342. *In*

R. F. Chapman and G. devBoer (eds.).  
*Regulatory mechanisms in insect feeding*.  
Chapman and Hall, New York.

**Daryaei, G. M., S. Darvishi, K. Etebari, and M. Salehi. 2007.** Host preference and nutrition efficiency of the gypsy moth, *Lymantria dispar* L. (Lymantriidae: Lepidoptera), on different poplar clones. *Turk. J. Agric.* 32: 469–476.

**Foss, L. K., and L. K. Rieske. 2003.** Species-specific differences in oak foliage affect preference and performance of gypsy moth caterpillars. *Entomol. Exp. Appl.* 108: 87–93.

**Grabstein, E. M., and J. M. Scriber. 1982.** Host-plant utilization by *Hyalophora cecropia* as affected by prior feeding experience. *Entomol. Exp. Appl.* 32: 262–268.

**Hemati, S. A., B. Naseri, G. N. Ganbalani, H. R. Dastjerdi, and A. Golizadeh. 2012.** Effect of different host plants on nutritional indices of the pod borer, *Helicoverpa armigera*. *J. Insect Sci.* 12:55. Available online: [www.insectscience.org/12.55](http://www.insectscience.org/12.55)

**Hwang, S. Y., C. H. Liu, and T. C. Shen. 2008.** Effects of plant nutrient availability and host plant species on the performance of two *Pieris* butterflies (Lepidoptera: Pieridae). *Biochem. Syst. Ecol.* 36: 505–513.

**Iran Ministry of Agriculture. 2011.** *Agricultural Statistics of Iran 2010–11*. Iran Ministry of Agriculture.

**Lazarevic, J., and V. Peric-Mataruga. 2003.** Nutritive stress effects on growth and digestive physiology of *Lymantria dispar* larvae. *Yugosl. Med. Biochem.* 22: 53–59.

**Liu, Z. D., D. M. Li, P. Y. Gong, and K. J. Wu. 2004.** Life table studies of the cotton bollworm, *Helicoverpa armigera* (Hübner) (Lepidoptera: Noctuidae), on different host plants. *Environ. Entomol.* 33: 1570–1576.

**Martin, L. A., and A. S. Pulin. 2004.** Host-plant specialization and habitat restriction in an endangered insect, *Lycaena dispar batavus* (Lepidoptera: Lycaenidae) I. Larval feeding and oviposition preferences. *Eur. J. Entomol.* 101: 51–56.

**Naseri, B., Y. Fathipour, S. Moharramipour, and V. Hosseininaveh. 2010.** Nutritional indices of the cotton bollworm, *Helicoverpa armigera*, on 13 soybean varieties. *J. Insect Sci.* 10: 151 available online: [www.insectscience.org/10.151](http://www.insectscience.org/10.151)

**Nathan, S. S., P. G. Chung, and K. Murugan. 2005.** Effect of biopesticides applied separately or together on nutritional indices of the rice leaf folder *Cnaphalocrocis medinalis*. *Phytoparasitica* 33: 187–195.

**Nation, J. L. 2001.** *Insect physiology and biochemistry*. CRC Press, Boca Raton, FL.

**Phillipson, J. 1981.** Bioenergetic options and phylogeny, pp. 20–45. In C. R. Townsend and P. Calow (eds.). *Physiological ecology: an evolutionary approach to resource use*. Blackwell Scientific, Oxford.

**Ruan, Y. M., and K. J. Wu. 2001.** Performances of the cotton bollworm, *Helicoverpa armigera* on different food plants. *Acta Entomol. Bohemoslov.* 44: 205–212.

**Sajjad, M., M. Ashfaq, A. Suhail, and S. Akhtar. 2011.** Screening of tomato genotypes

for resistance to tomato fruit borer, *Helicoverpa armigera* in Pakistan. *Pak. J. Agric. Sci.* 48: 49–52.

**Samraj, D. A., and B. V. David. 1988.** Life table studies on the spotted bollworm, *Earias vittella* (Fabricious) (Lepidoptera: Noctuidae) in cotton ecosystem. *J. Bombay Nat. Hist. Soc.* 85: 637–641.

**Schroeder, L. A. 1981.** Consumer growth efficiencies: Their limits and relationships to ecological energetics. *J. Theor. Biol.* 93: 805–828.

**Sharma, H. C., R. A. Agarwal, and M. Singh M. 1982.** Effect of some antibiotic compounds in cotton on post-embryonic development of spotted bollworm (*Earias vittella* F.) and the mechanism of resistance in *Gossypium arboreum*. *Proc. Ind. Acad. Sci. Anim. Sci.* 91: 67–77.

**Sheppard, C. A., and S. Friedman. 1990.** Influence of host plant, foliar phenology and larval dietary history on *Lymantria dispar* larval nutritional indices. *Entomol. Exp. Appl.* 55: 247–255.

**Singh. H., and G. Singh. 1975.** Biological studies on *Heliothis armigera* (Huebner) in Punjab. *Ind. J. Entomol.* 34: 154–164.

**Slansky, F., Jr. 1990.** Insect nutritional ecology as a basis for studying host plant resistance. *Fla. Entomol.* 73: 359–378.

**Slansky, F. J., and J. M. Scriber. 1985.** Food consumption and utilization, Vol. 4, pp. 87–163. In G. A. Kerkut and L. I. Gilbert. (eds). *Comprehensive insect physiology, biochemistry and pharmacology*. Pergamon Press, New York.

**Soo Hoo, C. F., and G. Fraenkel. 1966.** The consumption, digestion, and utilization of food plants by a polyphagous insect, *Prodenia eridania* (Cramer). *J. Insect Physiol.* 12: 711–730.


**Srinivasan, P. M. 1959.** Control of fruitborer, *H. armigera*(Hb) on tomato. *Ind. J.I-Tort* 16: 187–188.

**Talekar, N. S., Y. F. Chang, and S. T. Lee. 1984.** Tomato insect pests: major management strategies, pp. 53–171. In *Proceedings of the Symposium on the Insect Control of Vegetables in Taiwan*. Department of Agriculture and Forestry, Taiwan Provincial Government, Chung Hsing Village, Taiwan.

**Talekar, N. S., R. T. Opena, and P. Hanson P. 2006.** *Helicoverpa armigera* management: A review of AVRDC's research on host plant resistance in tomato. *Crop Prot.* 25: 461–467.

**Vattanatungum, A., and W. Ruchtapakornchai. 1978.** Principal insects which attack tomato in the tropics and their control, pp. 132–135. In *First International Symposium on Tropical Tomato*. Asian Vegetable Research and Development Center, Shanhua, Taiwan.

**Waldbauer, G. P. 1968.** The consumption and utilization of food by insects. *Adv. Insect Physiol.* 5: 229–288.



**Figure 1** Mean larval weight (A), food consumed (B) and feces produced (C) of whole larval instars of *Helicoverpa armigera* on different tomato cultivars. Bars represent standard error of the means. The means followed by different letters are significantly different (LSD,  $P < 0.01$ ).

Figure 1.



