

Absence of Geographic Population Structure in the Soybean Pod Borer *Leguminivora glycinivorella* (Tortricidae)

Authors: Wang, Hong, Han, Lanlan, Xu, Zhongxin, Ye, Lefu, Wang, Ling, et al.

Source: The Journal of the Lepidopterists' Society, 69(3) : 173-182

Published By: The Lepidopterists' Society

URL: <https://doi.org/10.18473/lepi.69i3.a4>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

ABSENCE OF GEOGRAPHIC POPULATION STRUCTURE IN THE SOYBEAN POD BORER
LEGUMINIVORA GLYCINIVORELLA (TORTRICIDAE)

HONG WANG, LANLAN HAN, ZHONGXIN XU, LEFU YE, LING WANG, KUIJUN ZHAO*

College of Agriculture, Northeast Agricultural University, Harbin 150030, China

AND

KEQIN WANG*

Institute of Plant Protection, Heilongjiang Academy of Agricultural Sciences, Harbin 150086, China

* Corresponding authors: email: kjzhao@163.com (Kuijun Zhao) and Wang.keqin@163.com (Keqin Wang)

ABSTRACT. In order to investigate geographic population structure and genetic diversity in the soybean pod borer *Leguminivora glycinivorella*, partial sequences of the mitochondrial DNA of 337 individuals from northeastern China were sequenced and analyzed. 16 haplotypes were found in CO II, and 14 haplotypes were defined in Cytb, including one haplotype shared by ten populations in each gene. *L. glycinivorella* populations are characterized by medium/low haplotype diversity and nucleotide diversity. The Tajima's *D* and Fu's *F_s* test indicated that there might not have been a recent population expansion. All pairwise gene flow *N_m* parameters were greater than one in the 10 populations. Molecular variance analysis (AMOVA) demonstrated that the observed genetic differentiation occurs primarily within populations, rather than among populations, no large-scale regional differences are detected. Genetic distance is not significantly correlated with geographical distance between populations. Maximum likelihood phylogenetic trees and a haplotype network showed that the haplotypes are distributed in different clades and no obvious geographical structure has been formed. The result suggested that geographic population structure among *L. glycinivorella* are not affected by geographic isolation and recent dispersal (some gene flow) resulting in no significant genetic differentiation occurred among populations.

Additional key words: *Leguminivora glycinivorella*; CO II; Cytb; genetic diversity; gene flow

The soybean pod borer *Leguminivora glycinivorella* (Mats.) is an important agricultural pest, widely distributed in China, Japan, Korea and Far East coast. In China, it is distributed in the northern and central regions, and, especially, the northeastern three provinces suffer seriously outbreaks. *L. glycinivorella* is a relatively monophagous pest, damaging grain by larval drill decay (Wu 2002). Its primary hosts are *Glycine max*, *Glycine ussuriensis* and *Sophora flavescens*. High insect feeding rates occurred in China (10%-30%), sometimes exceeding 50%. Serious impacts have been observed on soybean yield and quality (Wu 2001), and this pest is the primary target of prevention and control in soybean fields.

Genetic markers, in particular sequences of mitochondrial DNA (mtDNA), have been proven to be very informative for assessing genetic diversity and gene flow (Avise et al. 1987, Li et al. 2013, Coates et al. 2004, Meraner et al. 2008). Because of mtDNA's maternal inheritance, absence of intermolecular genetic recombination, fast evolutionary rate relative to nuclear DNA, availability of efficient PCR primers, and wealth of comparative data (Feng et al. 2012), it has been used extensively in studying population diversity, phylogeography and phylogenetic relationships at

various taxonomic levels (Yang et al. 2008). Sequences encoding mitochondrial cytochrome oxidase subunit II (CO II) are shown to be appropriate for intraspecific analysis due to the high degree of polymorphism observed (Li 2010, Liang et al. 2011, Li et al. 2013). Cytb gene has been documented to be sensitive in detecting genetic diversity and population genetic structure and has been used for analysis of populations in other lepidopteran species (Gao et al. 2011). This study aimed at identifying that whether geographic isolation could lead to genetic variation of *L. glycinivorella*, a species that is known to have a weak flight ability. A total of 159 partial mtDNA CO II gene and 178 partial mtDNA Cytb gene sequences were employed to assess geographic population structure, genetic diversity, gene flow, and intraspecific and interspecific population differentiation in ten *L. glycinivorella* populations in the northeastern China. Determining whether the genetic distance of this species was correlated with the geographical distance among populations or not requires the type of molecular phylogenetic study developed here. This baseline information is also critical for genetically-targeted management of *L. glycinivorella* within different crop regions.

MATERIALS AND METHODS

Sample collection. A total of 337 samples of last instar larvae of *L. glycinivorella* were collected from northeastern China in September 2013. The samples comprised ten populations of *L. glycinivorella*: Daqing (DQ), Dehui (DH), Gongzhuling (GZL), Harbin (HRB), Heihe (HH), Mudanjiang (MDJ), Jiamusi (JMS), Qiqihar (QQHR), Shenyang (SY), and Suihua (SH). Geographic locations are given in Fig. 1. All individuals were identified based on morphological characteristics, and samples were preserved in anhydrous ethanol and stored at -20 °C until DNA extraction.

DNA extraction, PCR amplification and sequencing. Genomic DNA was extracted according to a standard phenol-chloroform protocol (Sambrook & Russell 2001). The DNA was diluted to obtain a final concentration of 100 ng/mL. The CO II gene was PCR amplified using a set of universal primers CO II F (5'-TAGTGCAATGGATTAAACC-3') and CO II R: (5'-GTTTAAGAGACCACTACTTG-3') (Folmer et al. 1994). And the following is primers of Cytb gene: CYTB1 (5'-TATGTACTACCATGAGGACAAATATC-3'), CYTB2 (5'-ATTACACCTCCTAATTTATTAGGAAT-3') (Folmer et al. 1994). PCR amplification was performed using a Well Thermal Cycler (EDC-810) starting with 4 min of denaturation at 94°C, followed by 35 cycles of denaturation at 94°C for 50s, annealing at 48.3°C for 50 s, and extension at 72°C for 50 s, with a final extension at 72°C for 5 min. The reaction mixture contained approximately 3 µL of diluted genomic DNA as a template, 5 µL of 10×PCR buffer (100 mmol/L Tris-HCl, pH 8.3, 500 mmol/L KCl, 15 mmol/L MgCl₂), 1 µL of each primer, 4 µL of dNTP, and 0.5 µL of DNA polymerase (Trans GenBitech, 5 U/mL), resulting in a total volume of 50 µL with sterilized water. The PCR products were gel-purified using an agarose gel DNA purification kit (Trans GenBitech) following the manufacturer's instructions. The purified fragments were sequenced by the Beijing Huada Gene Research Center. Sequences were determined in both directions (using the same primers individually as for the PCR), and the electro-pherograms were verified by eye.

Data analysis. Nucleotide composition and variable sites were analyzed in MEGA 5.0 (Tamura et al. 2011). The genetic diversity indices of mtDNA, such as nucleotide diversity (π) (Lynch & Crease 1990) and haplotype diversity (H_d) (Nei 1987) were calculated using DnaSP 5.0 (Librado & Rozas 2009). The demographic history of *L. glycinivorella* was examined with the neutrality statistics of Tajima's D and Fu's F_s test (Kimura 1983, Tajima 1989), which can indicate




FIG.1 Map of northeastern China, showing sampling localities of *Leguminivora glycinivorella*. The ten sampling localities were Daqing (DQ), Dehui (DH), Gongzhuling (GZL), Harbin (HRB), Heihe (HH), Mudanjiang (MDJ), Jiamusi (JMS), Qiqihar (QQHR), Shenyang (SY), and Suihua (SH). A (Heilongjiang Prov.), B (Jilin Prov.), C (Liaoning Prov.).

whether population expansion has occurred. The geographical distance among collection sites was calculated according to latitude and longitude. Genetic relationship among haplotypes was reconstructed using the Maximum likelihood method in MEGA 5.0 (Tamura et al. 2011). A bootstrap analysis with 1000 replicates was used to evaluate phylogenetic relationships. We chose the method of Maximum likelihood (Felsenstein 1985) to construct a phylogenetic tree of different haplotypes according to the Kimura 2-parameter model. To depict phylogenetic and geographical relationships between the haplotype sequences, a haplotype network was created with the median-joining method using Network 4.6 (Bandelt et al. 1999). A hierarchical analysis of molecular variance (AMOVA) was performed to reveal the geographical structure of genetic variation using Arlequin 3.5.1.2 (Excoffier et al. 2005). The significance of the fixation index (F_{st}) was tested with 1000 permutations of the data set. Using the formula $F_{st} = 1/(1+2Nm)$, which is specific to organelle genetic data (Takahata & Palumbi 1985, Goldberg & Ruvolo 1997), we derived the values for Nm .




FIG.2 Neighbor-joining phylogenetic trees of haplotypes of different geographic populations of *Leguminivora glycinivorella*, A was for CO II, and B was for Cytb. *Grapholita molesta* (WQ1) and *Zeiraphera diniana* (WQ2) were used as the outgroup. All haplotypes in the populations were significantly separated from the outgroup.

TABLE 1. Genetic diversity, Tajima's *D* and Fu's *F_s* test in different geographic populations of *Leguminivora glycinivorella* based on CO II

Collecting locality	Abbreviation	H	Sample size	<i>Hd</i>	<i>II</i> (SD)	Tajima's <i>D</i>	Fu's <i>F_s</i>
Daqing	DQ	H1(10)H2(2)H3(2)	14	0.484(0.142)	0.00142(0.00042)	p>0.05	p>0.05
Dehui	DH	H1(13)H4(1)	14	0.143(0.119)	0.00063(0.00052)	p>0.05	p>0.05
Gongzhuling	GZL	H1(14)H3(1)H5(2) H6(1)	18	0.399(0.138)	0.00117(0.00040)	p>0.05	p>0.05
Harbin	HRB	H1(11)H7(6)	17	0.485(0.079)	0.00071(0.00012)	p>0.05	p>0.05
Heihe	HH	H1(8)H7(4)H8(2) H9(2)	16	0.700(0.090)	0.00620(0.00259)	p>0.05	p>0.05
Jiamusi	JMS	H1(13)H3(2)H7(2) H10(1) H11(1)	19	0.532(0.130)	0.00135(0.00039)	p>0.05	p>0.05
Mudanjiang	MDJ	H1(13) H12(1)	14	0.143(0.119)	0.00042(0.00035)	p>0.05	p>0.05
Qiqihar	QQHR	H1(10)H5(2)H7(4) H13(1)	17	0.618(0.106)	0.00159(0.00050)	p>0.05	p>0.05
Shenyang	SY	H1(12)H7(3) H14(1) H15(1)	17	0.493(0.131)	0.00133(0.00050)	p>0.05	p>0.05
Suihua	SH	H1(11)H4(1) H16(1)	13	0.295(0.156)	0.00150(0.00080)	p>0.05	p>0.05
Total population			159	0.463(0.047)	0.00170(0.00039)	p>0.05	p>0.05

* H, Haplotype distribution; *Hd*, haplotype diversity; *II*, nucleotide diversity; and standard deviation (SD).




FIG.3 Median-joining network of haplotypes of *Leguminivora glycinivorella* based on CO II gene. Each circle represents one unique haplotype, with the area being proportional to the frequency of the haplotype in the ten populations.

RESULTS

Genetic diversity. A 683 bp fragment of the mitochondrial CO II gene and a 415 bp fragment of the mitochondrial Cytb gene were amplified and sequenced from all of the collected samples. The average base composition for CO II gene was as follows: A=35.4%, T=40.5%, C=13.4%, G=10.7%. Another composition was found for Cytb gene: A=32.6%, T=41.0%, C=16.0%, G=10.4%. The results reveal a significantly high value for A/T, which conforms to the composition and structure characteristics of the mtDNA gene sequences of Lepidoptera (Jermin & Crozier 1994, Frati et al. 1997, Nei & Kumar 2002). Under the Kimura 2-parameter model, the estimated transition/transversion bias (R) for CO II gene was 2.3, which conforms to the principle that transition is greater than transversion in more closely related populations (Simon et al. 1994, Frati et al. 1997). Among the 337 individuals, there were no insertions or deletions observed in the examined sequences. Sixteen haplotypes were identified in the 159 samples of CO II gene, and Cytb gene contained fourteen haplotypes in the 178 individuals. The sequences were deposited in GenBank (KJ540178-KJ540193 for CO II haplotypes H1–H16, and KM358122-KM358135 for Cytb haplotypes h1–h14). The number of haplotypes ranged from two to five for each sampled population.

The genetic diversity indices, such as haplotype diversity (H_d) and nucleotide diversity (π) (Nei & Li 1979 Smith et al. 2006) are presented in Table 1 and Table 2. The mean haplotype diversity and nucleotide sequence diversity of CO II gene in the ten populations were 0.463 and 0.00170, respectively. Haplotype diversity ranged from 0.143 (MDJ) to 0.700 (HH), and nucleotide diversity ranged from 0.00042 to 0.00620. Haplotype diversity of Cytb gene ranged from 0.212 (MDJ) to 0.595 (HH), and nucleotide diversity ranged from 0.00068 to 0.01973.

Tests for neutral evolution (Tajima's D and Fu's F_s test) were performed to identify the presence of a selective sweep or a balancing selection in *L. glycinivorella* populations (Harpending et al. 1998). Statistical significance was not found for all populations in both Tajima's D and Fu's F_s test (Table 1 and Table 2), which indicated *L. glycinivorella* did not experience a recent population expansion ($P > 0.05$) among the ten populations. The results suggested a significant correlation between the observed and expected outcomes.

Phylogenetic and network analyses. Based on the principle of selecting a closely related species as the outgroup, *Grapholita molesta* (WQ1)(GenBank: KF552028.1) and *Zeiraphera diniana* (WQ2) (GenBank: FJ647109.1) were chosen as the outgroup




FIG. 4 Mantel test between genetic distance and geographic distance among ten populations of *Leguminivora glycinivorella*. (A) was for CO II, and (B) was for Cytb. Genetic distance of this species was not correlated with the geographical distance between populations.

(WQ1 for CO II and WQ2 for Cytb). The Maximum likelihood trees based on Kimura 2-parameter distances were constructed. All haplotypes in the populations were significantly separated from outgroup, but there was no clear evidence of a geographical spectrum among the haplotypes, and each cluster confidence coefficient was low. The haplotypes (CO II gene) of nearer geographical populations were divided into the same cluster (such as H5 for QQHR and H3 for DQ), but the haplotypes of further geographical populations were also divided into the same cluster (such as H3 for GZL and H10 for JMS), the similar results were found for Cytb. Overall, the phylogenetic trees did not reflect geographical position, indicating a lack of obvious geographic structure (Fig. 2).

To further depict the phylogenetic and geographical relationships among the identified sequences, haplotype networks were constructed with the median-joining method using Network 4.1 software (Fig. 3). The resulting network exhibited an approximate star-like pattern surrounding haplotypes H1 and H7. Haplotype H1 was the most common (72.3%), and was shared by all ten populations. Additionally, H7 was shared by five sample regions and was present in 11.9% of all individuals. Five individuals displayed as haplotype H3, four individuals were haplotype H5, and there were two individuals showing as haplotypes H2, H4, H8 and H9, respectively; whereas the other haplotypes were only

found once and were restricted to a single population, which is indicative of rare haplotypes. Moreover, similar results were found for Cytb (not listed).

Population structure analysis. The AMOVA analysis based on CO II gene haplotype frequencies revealed that 96.36% of the genetic variation occurred within populations, and 3.64% of the genetic variation occurred among populations, the similar result revealed in Cytb (Table 3). Therefore, genetic variation within populations was significantly greater than the genetic variation among populations, which suggests that the genetic differentiation of *L. glycinivorella* populations in northeastern China occurred primarily within populations, whereas the genetic variation among populations was relatively low.

Gene flow and genetic differentiation analysis. The fixation index (F_{st}) is a measure of the variance in gene frequencies between populations (Garcia et al. 2003), and bears a direct relationship to Nm , the product of a population's effective size and its female migration rate per generation (Wright 1969, Goldberg & Ruvolo 1997). Nm is thus the absolute number of female individuals who migrate among populations per generation on average. The genetic distances (F_{st}) and per-generation migration rates (Nm) between pairs of 10 populations are shown in Table 4 and Table 5. The pairwise genetic distances (F_{st}) of 45 pairs of populations ranged from -0.0420 to 0.2225. All of these showed no statistically

TABLE 2. Genetic diversity, Tajima's *D* and Fu's *F_s* test in different geographic populations of *Leguminivora glycinivorella* based on Cytb

Collecting locality	Abbreviation	H	Sample size	<i>Hd</i>	<i>H</i> (SD)	Tajima's <i>D</i>	Fu's <i>F_s</i>
Daqing	DQ	H1(14)H2(4)	18	0.366(0.112)	0.00176(0.00054)	p>0.05	p>0.05
Dehui	DH	H1(15)H2(3)	18	0.294(0.119)	0.00142(0.00057)	p>0.05	p>0.05
Gongzhuling	GZL	H1(16)H3(1)H4(4) H5(1) H6(1)	23	0.502(0.114)	0.01564(0.00495)	p>0.05	p>0.05
Harbin	HRB	H1(14)H4(4) H7(1) H8(2)	21	0.533(0.111)	0.01671(0.00514)	p>0.05	p>0.05
Heihe	HH	H1(11)H4(4) H8(2) H9(1)	18	0.595(0.109)	0.01973(0.00526)	p>0.05	p>0.05
Jiamusi	JMS	H1(14)H8(2)	16	0.233(0.126)	0.00112(0.00061)	p>0.05	p>0.05
Mudanjiang	MDJ	H1(11) H10(2)	13	0.212(0.142)	0.00068(0.00034)	p>0.05	p>0.05
Qiqihar	QQHR	H1(11)H4(2) H11(1) H12(1)	15	0.467(0.148)	0.01308(0.00616)	p>0.05	p>0.05
Shenyang	SY	H1(16)H2(1) H4(3) H11(1) H13(1)	22	0.468(0.123)	0.01373(0.00504)	p>0.05	p>0.05
Suihua	SH	H1(9)H4(4)H14(1)	14	0.538(0.115)	0.002216(0.00536)	p>0.05	p>0.05
Total population			178	0.443(0.044)	0.01143(0.00176)	p>0.05	p>0.05

*H, Haplotype distribution; *Hd*, haplotype diversity; *H*, nucleotide diversity; and standard deviation (SD)

TABLE 3. Analysis of molecular variance (AMOVA) for the CO II and Cytb sequences in *Leguminivora glycinivorella* 10 populations

Source of variation	<i>df</i>	Sum of sequences	Variance components	Percentage of variation(%)
	CO (Cytb)	CO (Cytb)	CO (Cytb)	CO (Cytb)
Among populations	9(9)	8.601(38.310)	0.02256Va(0.11194Va)	3.64(4.70)
With populations	149(168)	89.034(381.594)	0.59754Vb(2.27140Vb)	96.36(95.30)
Total	158(177)	97.635(419.904)	0.62011(2.38333)	

TABLE 4. The *F_{st}* value and gene flow *Nm* among ten populations of *Leguminivora glycinivorella* based on CO II

	DQ	DH	GZL	HRB	HH	JMS	MDJ	QQHR	SY	SH
DQ		7.41	Inf	Inf	9.85	Inf	3.56	35.73	Inf	19.42
DH	0.0632		21.92	3.81	5.69	12.66	Inf	4.40	14.51	Inf
GZL	-0.0105	0.0223		Inf	9.15	Inf	5.92	Inf	Inf	151.01
HRB	-0.0048	0.1159	-0.0013		11.35	Inf	1.75	Inf	Inf	13.83
HH	0.0483	0.0808	0.0518	0.0422		10.30	4.37	11.97	11.55	7.79
JMS	-0.0308	0.0380	-0.0221	-0.0161	0.0463		4.70	53.85	Inf	54.44
MDJ	0.1231	0.0000	0.0779	0.2225	0.1026	0.0961		2.12	4.60	12.80
QQHR	0.0138	0.1020	-0.0017	-0.0242	0.0401	0.0092	0.1906		Inf	15.73
SY	-0.0071	0.0333	-0.0136	-0.0353	0.0415	-0.0188	0.0981	-0.0140		Inf
SH	0.0251	-0.0420	0.0033	0.0349	0.0603	0.0091	0.0376	0.0308	-0.0098	

*Indicated *F_{st}* value's level of significance at P < 0.05; inf is infinite.

The data above the diagonal are *Nm*; the data below the diagonal are *F_{st}*.

TABLE 5. The *Fst* value and gene flow *Nm* among ten populations of *Leguminivora glycinivorella* based on Cytb

	DQ	DH	GZL	HRB	HH	JMS	MDJ	QQHR	SY	SH
DQ		Inf	3.15	2.91	2.40	3.15	2.78	5.18	4.79	1.77
DH	-0.0484		3.32	3.03	2.47	4.72	4.18	5.82	5.14	1.78
GZL	0.1369	0.1309		Inf	12.59	3.56	3.53	Inf	Inf	18.30
HRB	0.1468	0.1416	-0.0441		Inf	3.36	3.20	Inf	Inf	13.31
HH	0.1722	0.1684	0.0382	-0.0481		2.76	2.61	Inf	Inf	Inf
JMS	0.1369	0.0958	0.1233	0.1294	0.1532		Inf	6.26	5.30	1.76
MDJ	0.1525	0.1068	0.1242	0.1353	0.1607	-0.0246		6.16	5.06	1.73
QQHR	0.0881	0.0791	-0.0468	-0.0429	-0.0265	0.0740	0.0751		Inf	81.47
SY	0.0945	0.0887	-0.0388	-0.0375	-0.0247	0.0862	0.0900	-0.0559		Inf
SH	0.2203	0.2189	0.0266	0.0362	-0.0494	0.2212	0.2242	0.0.0061	-0.0026	

*Indicated *Fst* value's level of significance at $P < 0.05$; inf is infinite.
The data above the diagonal are *Nm*; the data below the diagonal are *Fst*.

TABLE 6. The natural logarithm of geographical distance (km) (above the diagonal) and pairwise genetic distance (below the diagonal) among *Leguminivora glycinivorella* populations based on CO II and Cytb (in parentheses).

	DQ	DH	GZL	HRB	HH	JMS	MDJ	QQHR	SY	SH
DQ		7.76	8.41	7.18	8.83	8.03	9.02	9.32	9.23	6.81
DH	0.0675 (0.0000)		7.23	7.06	9.35	8.64	8.65	7.27	9.12	7.85
GZL	0.0000 (0.1586)	0.0228 (0.1506)		8.15	9.63	8.76	8.22	8.45	8.54	8.57
HRB	0.0000 (0.1721)	0.1310 (0.1650)	0.0000 (0.0000)		9.05	9.18	8.65	8.79	7.80	6.71
HH	0.0508 (0.2080)	0.0878 (0.2025)	0.0546 (0.0000)	0.0441 (0.0000)		8.32	8.03	8.22	8.98	8.73
JMS	0.0000 (0.1586)	0.0395 (0.1060)	0.0000 (0.1406)	0.0000 (0.1486)	0.0485 (0.1809)		9.40	8.73	9.98	8.16
MDJ	0.1404 (0.1799)	0.0000 (0.1196)	0.0845 (0.1418)	0.2862 (0.1565)	0.1143 (0.1915)	0.1063 (0.0000)		9.12	9.62	8.32
QQHR	0.0140 (0.0965)	0.1136 (0.0859)	0.0000 (0.0000)	0.0000 (0.0000)	0.0418 (0.0000)	0.0093 (0.0800)	0.2355 (0.0812)		9.17	7.95
SY	0.0000 (0.1044)	0.0344 (0.0975)	0.0000 (0.0000)	0.0000 (0.0000)	0.0433 (0.0000)	0.0000 (0.0943)	0.1088 (0.0990)	0.0000 (0.0000)		9.23
SH	0.0257 (0.2825)	0.0000 (0.2802)	0.0033 (0.0000)	0.0362 (0.0000)	0.0641 (0.0000)	0.0092 (0.2840)	0.0391 (0.2890)	0.0318 (0.0000)	0.0000 (0.0000)	

significant genetic differentiation ($P > 0.05$), suggesting that all of the populations form one genetic group. In addition, gene flow among the 10 populations was estimated by Nm , which is the expected number of migrants exchanged among populations in each generation. All Nm values between pairs of populations were greater than one (Table 4 and Table 5). These results suggest that extensive gene flow occurred among all ten populations of *L. glycinivorella*.

The pairwise CO II and Cytb gene genetic distances were calculated with Kimura 2-parameter (Table 6). Mantel test showed no significant linear dependence between genetic distance and geographic distance both in CO II gene ($r=0.0492$, $p=0.595>0.05$) and Cytb gene ($r=-0.2143$, $p=0.116>0.05$) (Fig. 4), suggesting that long-term geographical isolation did not lead to genetic variation among the populations of *L. glycinivorella*.

DISCUSSION

Although external morphological characteristics are highly similar among *L. glycinivorella* populations, geographic isolation can result in genetic variation. Therefore, the rapid and accurate identification of whether *L. glycinivorella* populations have experienced genetic variation is important for scientific research and agricultural pest control. DNA barcoding offers a standardized system for determining whether species are the same or closely related based on the analysis of small fragments of DNA (Kruse & Sperling 2002, Landry et al. 1999). This study analyzed the genetic variation and gene flow (Nm) in ten *L. glycinivorella* populations based on mtDNA CO II and Cytb, which have a fast evolutionary rate (Meraner et al. 2008, Coates et al. 2004). Genetic variation in populations is created by evolutionary and demographic processes generating either heterogeneity or homogeneity among populations, and this variation determines the evolutionary potential of species. Population genetics provides models and tools for the interpretation of the processes that shape population structure. Sequences encoding mtDNA CO II and Cytb have been shown to be appropriate for intraspecific analysis due to the high degree of polymorphism observed (Pfunder et al. 2004).

Median-joining network for CO II showed that H1 was the most common haplotype (72.3%), and shared by all ten populations (similar result for Cytb), suggesting that potential adaptive exists in *L. glycinivorella* populations due to neutral processes like genetic bottleneck effects. However, the rare haplotype in each population also revealed that genetic differentiation existing among populations to some extent.

The mean haplotype diversity (h_d) of the *L. glycinivorella* populations was medium/low. The ability

to adapt to external environment condition showed by *L. glycinivorella* was relatively weak, which may be related to its biological characteristics. Perhaps limited habitat specialization and abundant host plant may cause relatively low genetic diversity within *L. glycinivorella* populations. In addition, chemical control that is widely used for the prevention and control of *L. glycinivorella*, may reduce the population size such that one common genetic structure persists. Haplotype diversity (h_d) and nucleotide diversity (π) showed differences in each population, and this may due to the wide distribution of *L. glycinivorella* in northeastern China. Genetic structure varies in different geographic population experiencing long term natural selection, and relationship between genetic structure of populations and degree of gene flow has proven to be an important evolutionary and variable (Zu 1999).

In this study, genetic diversity was most abundant in HH population. Studies have shown that *L. glycinivorella* belongs to the "long-day insect" group, diapausing when illumination time less than 15 h per day. The most serious injury of *L. glycinivorella* happens in July to August, when illumination time of high latitudes region (HH) is longer than low latitudes, such as HRB and other regions. Therefore, the beginning period of *L. glycinivorella* outbreak in the HH region occurs earlier than in HRB (Zhang 2013). Thus we may speculate that *L. glycinivorella* originated in the north and there have been founder effects as the *L. glycinivorella* species expanded its range southward, which leads to abundant genetic diversity in HH. In addition, we did obtain the samples randomly, but no reports could be found about whether different host plants varieties will impact on the biological characteristics of *L. glycinivorella*. Studies have shown that the *Thrips* species of *Mesothrips* develop different behavior characteristics and life history on different *Ficus* host (Tree & Walter, 2009).

Genetic variation within and among populations is explained by several processes, such as genetic drift, effective migration, natural selection, range fragmentation, expansion, habitation and low rates of mitochondrial evolution (Slatkin 1985, Avise 2004, Grant et al. 2006). Historical factors play a great important role in population phylogeny and evolution, Tajima's D and Fu's F_s test were both not statistically significant for all populations, indicating that *L. glycinivorella* did not experience a recent population expansion ($P > 0.05$) among the ten populations, population size remained relatively stable state.

Lower F_{st} values indicate a higher level of gene flow (Nm) and lower genetic differentiation exists among populations (Garcia et al. 2003). F_{st} values increase with

greater geographical separation of populations, implying isolation by distance. However, in our study, genetic differentiation in the populations was not highly associated with their geographic distribution. In addition, all populations from the ten locations had no apparent divergence and shared a great amount of gene flow. The N_m for all groups was greater than one, suggesting that temporary reproductive and geographic isolation did not act as a barrier for gene flow. This result was inconsistent with our original expectation where genetic distance of this species should be significantly correlated with the geographical distance between populations. This result also reflects that the level of gene flow among populations is not entirely determined by the migratory flight ability of insect. Fortunately, Fu (Personal communications) from the Chinese academy of agricultural sciences has found a small amount of *L. glycinivorella* in North Huang City island (At the junction of the Yellow Sea and the Bohai Sea which is located dozens of miles from mainland) using light trap for many years. So, we speculate that *L. glycinivorella* might have genetic exchange through other ways such as air-flow spread (passive migratory) or vectored by plant material transportation. Nevertheless, some potential factors should be studied further to understand the population structure and gene flow of *L. glycinivorella*.

The Maximum likelihood cluster results suggested that most of the populations were clustered together, showing no significant phylogeographic structure exists. The resulting network further supported this finding. The data presented here from mtDNA CO II and Cytb gene sequences reveal low genetic diversity and insignificant genetic differentiation existing in the *L. glycinivorella* of northeastern China. However, we examined only two portion of their entire genome in this study. The use of multiple genetic marker systems could increase the resolving power of future genetic studies (Gruenthal et al. 2007). Further studies using nuclear markers and more samples are needed to extend and corroborate the present findings. These studies would help us to understand the comprehensive population evolution and gene flow in *L. glycinivorella*, and thus achieve better prevention and control.

ACKNOWLEDGMENTS

We thank Keith S. Summerville and two anonymous reviewers for helpful reviews and recommendations for improvement, Dr. Wenpeng Sun for acquisition of specimens and helpful insights. We also thank Heilongjiang Academy of Agricultural Sciences and many others for allowing us to examine specimens during visits. This work was supported by the Fund of Common Wealth Industry (agriculture) Special Research (201103002) and the earmarked fund for Modern Agro-industry Technology Research System (CARS-04).

LITERATURE CITED

- AVISE, J. C. 2004. Molecular Markers, Natural History, and Evolution, second ed. Sinauer Associates, Sunderland (Massachusetts) p: 684.
- AVISE, J. C., J. AMOLD, R. M. BALL, E. BERMINGHAM, T. LAMB, J. E. NEIGEL, C. A. REEB, & N. C. SAUNDERS. 1987. Intraspecific phylogeography: The mitochondrial DNA bridge between population genetics and systematic. *Annu. Rev. Ecol. Syst.* 18: 489–522.
- BANDELT, H. J., P. FORSTER, & A. ROHL. 1999. Median-joining networks for inferring intraspecific phylogenies. *Mol. Biol. Evol.* 16: 37–48.
- COATES, B. S., D. V. SUMERFORD, & R. L. HELLMICH. 2004. Geographic and voltinism differentiation among North American *Ostrinia nubilalis* (European corn borer) mitochondrial cytochrome c oxidase haplotypes. *J. Insect Sci.* 4: 35.
- EXCOFFIER, L., G. LAVAL, & S. SCHNEIDER. 2005. Arlequin (version 3.0): an integrated software package for population genetics data analysis. *Evol Bioinformatics* 1: 47–50.
- FELSENSTEIN, J. 1985. Confidence limits on phylogenies, an approach using the bootstrap. *Evolution* 39: 783–791.
- FENG, H. Z., L. WANG, W. C. GUO, Y. CAO, M. L. YANG, H. M. LIU, & J. J. XU. 2012. Cloning and sequencing analysis of the coding moth *Cydia pomonella* COI gene. *Chinese Journal of Applied Entomology* 49: 1503–1507.
- FOLMER, O., M. BLACK, W. HOEH, R. LUTZ, & R. VRIJENHOEK. 1994. DNA primers for amplification of mitochondrial cytochrome c oxidase subunit I from diverse metazoan invertebrates. *Mar. Biol. Biotech.* 3: 294–299.
- FRATI, F., C. SIMON, J. SULLIVAN, & D. L. SWOFFORD. 1997. Evolution of the mitochondrial cytochrome oxidase II gene in Collembola. *J. Mol. Evol.* 44: 145–158.
- GAO, J. B., X. W. ZHANG, G. N. ZHOU, & J. X. LIU. 2011. Genetic structure of pine caterpillar (*Dendrolimus*) populations based on the analysis of Cytb gene sequences. *Acta Ecol. Sinica* 31: 1727–1734.
- GARCIA, B. A., C. MANFREDI, L. FICHERA, & E. L. SEGURA. 2003. Short report: variation in mitochondrial 12s and 16s ribosomal DNA sequences in natural populations of *Triatoma infestans* (Hemiptera: Reduviidae). *Am. J. Trop. Med. Hyg.* 68: 692–694.
- GOLDBERG, T. L. & M. RUVOLO. 1997. The geographic apportionment of mitochondrial genetic diversity in east African chimpanzees, *Pan troglodytes schweinfurthii*. *Mol. Biol. Evol.* 14: 976–984.
- GRANT, W. S., I. B. SPIES, & M. F. CANINO. 2006. Biogeographic evidence for selection on mitochondrial DNA in North Pacific Wall-eye Pollock *Theragra chalcogramma*. *J. Hered.* 97: 571–580.
- GRUENTHAL, K. M., L. K. ACHESON, & R. S. BURTON. 2007. Genetic structure of natural populations of California red abalone (*Haliotis rufescens*) using multiple genetic markers. *Mar. Biol.* 152: 1237–1248.
- HARPENDING, H. C., M. A. BATZER, M. GURVEN, L. B. JORDE, A. R. ROGERS, & S. T. SHERRY. 1998. Genetic traces of ancient demography. *Proc. Natl. Acad. Sci. USA*, 95: 1961–1967.
- HEBERT, P. D., M. Y. STOECKLE, T. S. ZEMLAK, & C. M. FRANCIS. 2004. Identification of birds through DNA barcodes. *PLoS biology* 2 (10), e312. [http://dx. doi. org/10. 1371/ journal. pbio. 0020312](http://dx.doi.org/10.1371/journal.pbio.0020312).
- JERMIN, L. S. & R. H. CROZIER. 1994. The cytochrome b region in the mitochondrial DNA of the ant *Tetraponera rufoniger*: sequence divergence in Hymenoptera may be associated with nucleotide content. *J. Mol. Evol.* 38: 282–294.
- KIMURA, M. 1983. The Neutral Theory of Molecular Evolution. Cambridge University Press, Cambridge, Massachusetts.
- KRUSE, J. J. & F. A. H. SPERLING. 2002. Phylogeny of Nearctic species of the Xylosteana group of *Archips* Hübner (Lepidoptera: Tortricidae) based on combined analysis of morphological and mitochondrial DNA data sets. *Ann. Entomol. Soc. Am.* 95:288–301.
- LANDRY, B., J. A. POWELL, & F. A. H. SPERLING. 1999. Systematics of the *Argyrotaenia franciscana* (Lepidoptera: Tortricidae) species

- group: evidence from mitochondrial DNA. *Ann. Entomol. Soc. Am.* 92:40–46.
- LI, C. X., E. B. MA, & X. Y. ZHENG. 2003. Genetic differentiation of different populations of four locust species in China. *Acta Genetica Sinica* 30: 234–244.
- LI, J., Y. ZHANG, Z. Y. WANG, K. L. HE, & Q. WANG. 2010. Genetic differentiation and gene flow among different geographical populations of the Asian corn borer, *Ostrinia furnacalis* (Guenée) (Lepidoptera: Crambidae) in China estimated by mitochondrial CO II gene sequences. *Acta Entomol. Sinica* 53: 1135–1143.
- LI, L. L., Y. YU, D. GUO, Y. L. TAO, & D. CHU. 2013. Genetic structure of the overwintering Asian corn borer, *Ostrinia furnacalis* (Guenée) collections in Shandong of China based on mtCOII gene sequences. *Acta Ecologica Sinica* 33: 4377–4386.
- LI, Y. T., K. WANG, Y. ZHENG, Q. L. MEN, L. QIAN, Y. L. AN, J. N. FENG, Y. L. ZHANG, & M. H. CHEN. 2013. COI gene analysis based genetic differentiation of *Cydia pomonella* (Lepidoptera: Tortricidae) populations in Hexi Corridor. *J. Northwest A & F Univ. (Nat. Sci. Ed.)* 41: 85–90.
- LIANG, R. X., Z. Y. WANG, K. L. HE, B. CONG, & J. LI. 2011. Genetic diversity of geographic populations of *Monolepta hieroglyphica* (Motschulsky) (Coleoptera: Chrysomelidae) from North China estimated by mitochondrial CO II gene sequences. *Acta Entomol. Sinica* 54: 828–837.
- LIBRADO, P. & J. ROZAS. 2009. DnaSP v5: a software for comprehensive analysis of DNA polymorphism data. *Bioinform.* 25: 1451–1452.
- LYNCH, M. & T. J. CREASE. 1990. The analysis of population survey data on DNA sequence variation. *Mol. Biol. Evol.* 7: 377–394.
- MERANER, A., A. BRANDSTÄTTER, R. THALER, B. ARAY, M. UNTERLECHNER, H. NIEDERSTÄTTER, W. PARSON, R. ZELGER, VIA. J. DALLA, & R. DALLINGER. 2008. Molecular phylogeny and population structure of the codling moth (*Cydia pomonella*) in Central Europe: I. Ancient clade splitting revealed by mitochondrial haplotype markers. *Mol. Phy. Evol.* 48:825–837.
- NEI, M. 1987. *Molecular Evolutionary Genetics*. Columbia University Press, New York, USA, pp. 10–88.
- NEI, M. & S. KUMAR. 2002. (translated by LV, B.Z., ZHONG, Y., GAO, L.P ET AL.) *Molecular Evolution and Phylogenetics*. Higher Education Press, Beijing.
- NEI, M. & W. H. LI. 1979. Mathematical model for studying genetic variation in terms of restriction endonucleases. *Proc. Natl. Acad. Sci.* 76: 5269–5273.
- PFUNDER, M., O. HOLZGANG, & J. E. FREY. 2004. Development of microarray-based diagnostics of voles and shrews for use in biodiversity monitoring studies, and evaluation of mitochondrial cytochrome oxidase I vs. cytochrome b as genetic markers. *Mol. Ecol.* 13: 1277–1286.
- SAMBROOK, J. & D. W. RUSSELL. 2001. *Molecular Cloning: a Laboratory Manual*, third ed. Cold Spring Harbor Laboratory Press, New York.
- SIMON, C., F. FRATI, A. T. BECKENBACH, B. CRESPI, H. LIU, & P. FLOOK. 1994. Evolution, weighting, and phylogenetic utility of mitochondrial gene sequences and a compilation of conserved polymerase chain reaction primers. *Ann. Entomol. Soc. Am.* 87: 651–701.
- SLATKIN, M. 1985. Gene flow in natural populations. *Annu. Rev. Ecol. Syst.* 16: 393–430.
- SMITH, M. A., N. E. WOODLEY, D. H. JANZEN, W. HALLWACHS, & P. D. N. HEBERT. 2006. DNA barcodes reveal cryptic host-specificity within the presumed polyphagous members of a genus of parasitoid flies (Diptera: Tachinidae). *Proc. Natl. Acad. Sci. USA* 103: 3657–3662.
- TAJIMA, F. 1989. Statistical method for testing the neutral mutation hypothesis by DNA polymorphism. *Genetics* 123: 585–595.
- TAKAHATA, N. & S. R. PALUMBI. 1985. Extranuclear differentiation and gene flow in the finite island model. *Genetics* 109: 441–457.
- TAMURA, K., D. PETERSON, N. PETERSON, G. STECHER, M. NEI, & S. KUMAR. 2011. MEGA 5, molecular evolutionary genetics analysis using maximum Likelihood, evolutionary distance, and maximum parsimony methods. *Mol. Biol. Evol.* 28: 2731–2739.
- TREE, D. J. & G. H. WALTER. 2009. Diversity of host plant relationships and leaf galling behaviours within a small genus of thrips *Cynaikothrips* and *Ficus* in south east Queensland, Australia. *Aust. J. Entomol.* 48: 269–275.
- WRIGHT, S. 1969. *Evolution and the Genetics of Populations*. In: *The Theory of Gene Frequencies*, vol. 2. Univ. of Chicago Press, Chicago.
- WU, B. Z. 2001. Study of new methods of prevention and control of *Leguminivora glycinivorella*. *Soybean Bulletin* 3: 9.
- WU, J. X. 2002. *Agricultural Entomology (The North)*. China Agriculture Press, Beijing. 259.
- YANG, R. S., Z. Y. WANG, K. L. HE, & S. X. BAI. 2008. Sequence analysis of mtDNA CO II gene and molecular phylogeny of *Ostrinia* spp. from China (Lepidoptera: Crambidae). *Acta Ecol. Sinica* 51: 182–189.
- ZHANG, W., B. H. LI, H. P. LI, & Y. J. LI. 2013. Occurrence Regularity of *Leguminivora glycinivorella* during 2009–2012 in Heihe Region. *Crops* 4: 146–148.
- ZU, Y. G., M. SUN, & L. KANG. 1999. *Theory, Method, and Application of Molecular Ecology*. Higher Education Press, Beijing.

Submitted for Publication 1 October 2014; revised and accepted 23 December 2014.