

A Simple Technique to Sample Pollen From Moths and Its Applications to Ecological Studies

Authors: Atwater, Montana M., and Lott, Terry

Source: The Journal of the Lepidopterists' Society, 65(4) : 265-267

Published By: The Lepidopterists' Society

URL: <https://doi.org/10.18473/lepi.v65i4.a8>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

A SIMPLE TECHNIQUE TO SAMPLE POLLEN FROM MOTHS AND ITS APPLICATIONS
TO ECOLOGICAL STUDIES**Additional key words:** palynology, natural history, ecology

Pollen is a physical link between many plant-animal interactions and can be a useful tool for biologists in ecological studies. Pollen found on field collected moths can be sampled and identified. Location of pollen, number of pollen grains, and pollen identification can be recorded for individual specimens. These data can be applied to explore a number of life history phenomena such as nectar plant use and pollination ecology. Further, with the option of using museum specimens, researchers can explore ecological questions from a larger geographic and temporal scale otherwise not easily obtained in a limited field season.

Moths and their interactions with their nectar plants are diverse yet notoriously difficult for researchers to investigate. Apart from larger hovering moths (e.g. Sphingidae), moths that visit flowers at night are often small and cryptic. They are known to seek nectar from small white or pale colored flowers often situated at the tips of branches (Oliviera et. al. 2004; Makholela & Manning 2006; Okamoto et. al. 2008; Tasen et. al. 2009). Still, moths nectaring on plants is a phenomenon rarely observed and documenting plant-moth interactions in the field is a formidable task. To complement such studies, investigating pollen found on moth specimens can be applied to further document nectar plants and quantify pollination efficiency.

Researchers have dabbled into exploring these methods to investigate life history questions relating to Lepidoptera pollination ecology (Darwin 1885; Wiklund et. al. 1979; Courtney et. al. 1982; Jennerston 1984; Tasen et. al. 2009), migration (Mikkola 1971; Lingren et. al. 1993), and even overwintering (Berkhouson & Shapiro 1994). Many of these studies used similar techniques to identify pollen with Scanning Electron Microscopy (SEM) and often removed the head of the specimen in order to extract the pollen. The techniques described here offer a less expensive and invasive alternative to using SEM to sample and identify pollen from field collected fresh or museum specimens.

Materials. Stereo microscope (7–15x magnification), hot plate, probe with fine point (see below), glycerin gel stained with pigment (see below), glass microscope slide with cover slips, small weights (~40 grams), slide warmer, reflective microscope with 10–60x objectives,

camera attachment and imaging software (AxioVision Rel. 4.5), pollen keys and regional identification manuals, and/or plant samples from the field.

Probe. Wooden dowel 14 cm long and 0.2 cm in diameter, with an embedded tip of a #2 insect pin.

Glycerin gel with stain. Add 20 g of gelatin (Crescent Chemical Company, Cat. # 23310.02) to 70 ml boiling distilled water, once thoroughly mixed, add 60 ml glycerin (Fisher Scientific, Cat. # 633-4) and 1.2 g phenol (Fisher Scientific, Cat. # G33-4), then after crystals dissolve, add 22 drops of Safranin-O stain (Fisher Scientific, Cat. # S670-25).

Collecting and Preserving Specimens. Field collected specimens should be kept in separate glass vials to prevent pollen contamination. Store moths in vials in a freezer until use in pollen analysis. This will preserve the specimen and help reduce loss of scales or pollen.

Scanning for pollen. To scan a moth for pollen, remove from the vial and pin the specimen through the thorax with an appropriately sized insect pin. Using a small piece of foam to stabilize the pinned specimen, scan the specimen for pollen under a stereomicroscope, paying special attention to the mouthparts (Fig. 1-I). If the proboscis is not visible, use a minuten insect pin to gently bend up from the base of the proboscis until fully exposed. If the specimen is not hydrated, you may need to rehydrate the specimen in a relaxing chamber long enough to rehydrate the proboscis.

Extracting pollen from specimen. Heat the glycerin gel on a hot plate at 52°C in a water bath until it has reached liquid form. Place microscope slide(s) on a slide warmer. Locate pollen on the moth using a stereomicroscope. Pour a small portion of the glycerin gel onto the microscope slide. Gently dip the probe tip into the glycerin gel. For pollen clumps, a small drop of glycerin gel works well. For individual pollen grains, a short streamer of glycerin gel works well, especially for grains embedded between the scales. Once the pollen is adhered to the probe, transfer the pollen from the specimen to the microscope slide into the drop of glycerin gel. Cover the drop with a cover slip (Fig. 1-II), and prepare a label for the microscope slide. Allow the slide to rest on the slide warmer long enough to enable the gel to stain the pollen (at least 24 hours). Seal the edge of the coverslip with clear fingernail polish.




FIG. 1. **I**) Pollen grains found on proboscis and palpi of *Samea ecclesiastis* (Pyraloidea). **II**) Photograph of pollen grain, *Eupatorium* sp. (Asteraceae), taken with AxioVision imaging software. **III**) Microscope slide with pollen sample in glycerin gel stain, cover slip, and weight on warming plate.

Identifying pollen: Using a reflective microscope at 40X (objective) and 1.25X (optivar), locate the pollen on the microscope slide and photograph using an attached digital camera and imaging software (Fig. 1-III). Using this image, identify the pollen with either a pollen key (Kapp et al. 2000), pollen identification manual (Jelks 2001), and/or matching pollen with pollen sample extracted from a properly identified plant (i.e. use of pollen collections can be arranged by collection managers at existing palynology institutions).

If the use of a palynology collection is not feasible, it is also possible to create a pollen library by which to match pollen. Pollen can be sampled from flowers observed in bloom within the same time and vicinity that the moths were collected. This is most beneficial when working in a habitat where the majority of plants are not yet represented in a palynology collection. Creating a pollen library with the plants in the area of study is not only beneficial for identifying pollen found on the moths, but can also be preserved in a local palynology library as a resource for future researchers.

ACKNOWLEDGEMENTS

The authors would like to thank David Jarzen for helpful comments on pollen identification and the use of the pollen collection. Jacqueline Y. Miller, Deborah L. Matthews, and Christian Salcedo provided constructive comments on the manuscript.

LITERATURE CITED

- BERKHOUSEN, A. E., & A. M. SHAPIRO. 1994. Persistent pollen as a tracer for hibernating butterflies: the case of *Hesperia juba* (Lepidoptera: Hesperidae). *Great Basin Naturalist* 54(1): 71-78.
- COURTNEY, S. P., C. H. HILL, & A. WESTERMAN. 1982. Pollen carried for long periods by butterflies. *Oikos* 38 (2): 260-263.
- DARWIN, C. R. 1885. The various contrivances by which orchids are fertilized by insects. John Murray, Albermarle St., London. 316 pp.
- JELKS, M. L. 2001. Allergy pollen key with images. M. L. Jelks publisher, Sarasota, FL, USA. 22 pp.
- JENNERSTON, O. 1984. Flower visitation and pollination efficiency of some North European butterflies. *Oecologia* (Berlin) 63:80-89.
- KAPP, R. O., O. K. DAVIS, & J. E. KING. 2000. Pollen and spores. 2nd ed. American Association of Stratigraphic Palynologists Foundation, College Station, TX, USA. 279 pp.
- LINGREN, P. D., V. M. BRYANT, J. R. RAULSTON, M. PENDLETON, J. WESTBROOK, & G. D. JONES. 1993. Adult feeding host-range and migratory activities of corn-earworm, cabbage-looper, and celery looper (Lepidoptera, Noctuidae) moths as evidenced by attached pollen. *J. Econ. Entomol.* 86: 1429-1439.
- MAKHOLELA, T., & J. C. MANNING. 2006. First report of moth pollination in *Struthiola ciliata* (Thymelaeaceae) in southern Africa. *S. African J. Botany* 72: 597-603.
- MIKKOLA, K. 1971. Pollen analysis as a means of studying the migration of Lepidoptera. *Ann. Entomol. Fenn.* 37: 136-139.
- OKAMATO, T., A. KAWAKITA, & M. KATO. 2008. Floral adaptations to nocturnal moth pollination in *Diplomorpha* (Thymelaeaceae). *Plant Species Biol.* 23: 192-201.
- OLIVIERA, P. E., P. E. GIBBS, & A. A. BARBOSA. 2004. Moth pollination of woody species in the Cerrados of Central Brazil: a case of so much owed to so few? *Plant Systematics and Evolution* 245(1): 41-54.
- TASEN, W., S. TANGMITCHAROEN, M. THAKEA, & K. OGATA. 2009. Insect pollination of *Aquilaria crassna* (Thymelaeaceae): Effect of moths for the fruit setting in Thailand. *J. Faculty Agr. Kyushu Univ.* 54 (2): 321-328.
- WIKLUND, C., T. ERIKSSON, & H. LUNDBERG. 1979. The wood white butterfly, *Leptidea sinapis*, and its nectar plants: a case of mutualism or parasitism? *Oikos* 33: 358-362.

MONTANA M. ATWATER, *McGuire Center for Lepidoptera and Biodiversity, Florida Museum of Natural History, University of Florida, Gainesville, FL 32611-2710, USA; email: matwater@flmnh.ufl.edu, and* TERRY LOTT, *Paleobotany and Palynology Laboratory, Florida Museum of Natural History, University of Florida, Gainesville, FL 32611-7800, US; email: lott@flmnh.ufl.edu*

Received for publication 14 December 2010; revised and accepted 31 May 2011