

On the Finding of Dead Ants Attached to Saturniid Caterpillars: Evidence of Successful Deterrent Chemistry?

Author: Collins, Michael M.

Source: The Journal of the Lepidopterists' Society, 67(1) : 62-63

Published By: The Lepidopterists' Society

URL: <https://doi.org/10.18473/lepi.v67i1.a10>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

ON THE FINDING OF DEAD ANTS ATTACHED TO SATURNIID CATERPILLARS: EVIDENCE OF SUCCESSFUL DETERRENT CHEMISTRY?

Additional key words: ant predation, anti-predator mechanism, chemical defense

Hyalophora cecropia (Saturniidae) is one of the most prominent natural history icons of North America, widely used to teach insect life history in schools and collected and reared by amateur naturalists for generation upon generation. The larvae are as striking as the colorful adults, and display a vivid array of scoli colors, which change with instar. Yet, until very recently, no reported evidence has appeared that the larvae are in any way toxic. As any breeder knows, larvae of all ages appear all too vulnerable to a host of predators and parasitoids (Collins & Weast 1961, p. 84; Tuskes et al. 1996).

Deml & Dettner (2003) discovered an unexpected chemical defense in cecropia larvae. Each scoli is associated with a secretory gland, the complex chemistry of which predictably varies by scoli color and instar. Their conclusion, based also on previous work (Deml & Dettner 1990, 1993, 1997), is that these chemicals are defensive, are released when small spines are broken off a scoli during an attack, and are directed against a sequence of instar-specific predators.

Their work also showed that the chemistry of the body haemolymph in *Hyalophora* does not contain these characteristic chemicals. Human skin seems unaffected by the larval chemistry of scoli, unlike the effect of urticating spines in other saturniid larvae such as the Hemileucinae. There is one obscure record of a robin apparently killed when attempting to eat a mature *H. columbia gloveri* larva (Duncan 1941).

On August 15, 2011 I collected a near-mature *Hyalophora* larva on coyote willow (*Salix exigua*) at 1700

m near a creek flowing into a small canyon along highway 89, just west of US 395. This area is at the eastern edge of a hybrid zone, across Monitor Pass, between *H. c. gloveri* and *H. euryalus*. The larval phenotype was intermediate between these two species; it possessed the spiny scoli typical of *gloveri*, although somewhat reduced in size. Upon close inspection I noticed a dead ant attached to the second thoracic segment, right side, near scoli L₁ (Fig. 1). The ant's abdomen was shrunk, indicating that the ant had remained attached for some days (the fifth instar in *Hyalophora* typically lasts for 7–12 days in the wild). After photographing the larva in the field, I left the ant attached to monitor the health and growth of the larva. It fed normally and spun a cocoon in 5 days. The head of the ant remained attached, but unfortunately the body of the ant became dislodged and lost before I could bring the larva into my lab. Under the microscope I noticed a single spine broken off each of two nearby scoli. From my photographs ant experts Phil Ward (Dept. Entomology, Univ. Calif. Davis) and James Trager (Shaw Nature Res., Gray Summit MO) identified it as belonging to the *Formica rufi* species group, probably either *F. ravida* or *F. moki*.

Surrounding the ant and along that side of the body of the larva were the dried remnants of what appeared to be regurgitated gut contents. *Hyalophora* larvae are not known to regurgitate gut contents unless severely attacked, but when this does occur the larva will curl its body toward the attacker and thrash about. My interpretation is that this action may have killed the ant by



FIG. 1. *Formica* ant (dead) attached to *Hyalophora* larva, Monitor Pass, Mono Co. CA. Note brown stains of dried regurgitant surrounding ant.



FIG. 2. Dead *Formica* (white arrows) attached to *Hemileuca eglanterina*, Monitor Pass, Mono. Co. CA. A similar photograph appears in Tuskes et al., 1996, p.20.

piercing its body with a scoli spine, which broke off and released a toxin. There is no evidence that the gut contents of *Hyalophora* are unpalatable.

In July 1973, coincidentally also on Monitor Pass, near the summit at 2500 m, I photographed a mature larva of *Hemileuca eglanterina* on snowberry (Fig. 2). Attached to the larva were two dead ants, appearing to be the same species of mound-building *Formica* ant as that found on the *Hyalophora*. Again, the simplest interpretation is that the ants died as a result of encountering defensive chemistry, in this case produced by a stinging larva.

Certainly, more careful work needs to be done on the subject of ant predation on lepidopterous larvae. The research on the specialized association of ants with lycaenid butterflies is a fascinating exception, but given the paramount ecological role of ants in biotic communities (Hölldobler & Wilson 1994), the importance of predatory ants in regulating the abundance and distribution of Lepidoptera is poorly understood. Michael Singer (Dept. Biology, Wesleyan U.) is currently researching these topics in temperate regions, and in our discussions pointed out to me two interesting studies: Karhu (1998) on ant exclusion experiments in a boreal forest, and Dyer (1995) on assaying ant predation on protected vs. unprotected Lepidoptera larva in the tropics. Even simple observations of ant predation on captive larvae placed in the field—'staking out lambs in lion country'—would be worthwhile, as suggested for *Eupackardia calleta* (Collins, 2007, p. 41ff), based on its blood chemistry and aposematic coloration. I hope this short note will stimulate further work.

ACKNOWLEDGEMENTS

I thank Phil Ward, University of California, Davis, Entomology and James Trager, Shaw Nature Reserve, Gray Summit, MO for

their efforts to identify the ant species from photographs. Mike Singer, Wesleyan University, Biology Dept., suggested certain references on ant predation and both he and Dave Wagner encouraged the publication of this note. John Rawlins, Carnegie Museum of Natural History – Invertebrate Section, kindly made a digital file from my 1973 35mm slide.

LITERATURE CITED

- COLLINS, M. M. 2007. Moth Catcher. University of Nevada Press, Reno. 200pp.
- COLLINS, M.M. AND R. D. WEAST. 1961. The Wild Silk Moths of the United States. 138pp. Privately published.
- DEML, R. AND K. DETTNER. 1990. Chemical defense of *Eudia* (*Saturnia*) *pavonia* caterpillars. *Naturwissenschaften* 77:588-590.
- DEML, R. AND K. DETTNER. 1993. Biogenic amines and phenolics characterize the defensive secretion of saturniid caterpillars (Lepidoptera: Saturniidae): a comparative study. *J. Comp. Physiol.* 163:123-132.
- DEML, R. AND K. DETTNER. 1997. Chemical defense of emperor moths and tussock moths (Lepidoptera: Saturniidae, Lymantriidae). *Entomol. Gen.* 21:225-251.
- DEML, R. AND K. DETTNER. 2003. Comparative morphology and secretion chemistry of the scoli in caterpillars of *Hyalophora cecropia*. *Naturwissenschaften* 90:460-463.
- DUNCAN, J. B. 1941. The biological control of *Platysamia gloveri* (Str.) with special reference to enemies of the pupal stage (Saturniidae, Lepidoptera). Master's thesis, Univ. Utah, Salt Lake City.
- DYER, L. A. 1995. Tasty generalists and nasty specialists? Antipredator mechanisms in tropical Lepidopteran larvae. *Ecology* 75:1483-1496.
- HÖLLDOBLER, B. AND E. O. WILSON. 1994. Journey to the Ants: A Story of Scientific Exploration. Harvard University Press, Cambridge.
- KARHU, K. J. 1998. Effects of ant exclusion during outbreaks of a defoliator and a sap-sucker on birch. *Ecol. Entomol.* 23:185-194.
- TUSKES, P. M., J. P. TUTTLE, AND M. M. COLLINS. 1996. The Wild Silk Moths of North America. Cornell University Press. 250pp.

MICHAEL M. COLLINS, *Research Associate, Carnegie Museum of Natural History, Section of Invertebrates, Pittsburgh PA. Mailing address: 215 Prospect St., Nevada City, CA 95959. email: michaelmerlecollins@comcast.net*

Submitted for publication 16 September 2011; revised and accepted 5 May 2012.