



## **New species of the genus *Laena* Dejean (Coleoptera: Tenebrionidae) from southeastern Asia (3)1**

Author: Schawaller, Wolfgang

Source: Integrative Systematics: Stuttgart Contributions to Natural History, 1(1) : 71-78

Published By: Stuttgart State Museum of Natural History

URL: <https://doi.org/10.18476/insy.v01.a8>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

# New species of the genus *Laena* Dejean (Coleoptera: Tenebrionidae) from southeastern Asia (3)<sup>1</sup>

WOLFGANG SCHAWALLER

## Abstract

New species of the genus *Laena* Dejean, 1821 (Tenebrionidae: Lagriinae) are described from central India (*L. centroindica* n. sp.), from northeastern India in Arunachal Pradesh and Assam (*L. bomdilaica* n. sp., *L. etalinica* n. sp., *L. gyatsoi* n. sp., *L. monpa* n. sp.), from Myanmar (*L. mandalayica* n. sp., *L. schillhammeri* n. sp.), and from the Malay Peninsula (*L. larutica* n. sp.). *L. bicolor* Schuster, 1926 is rediscovered in Myanmar since the original description, and *Laena kurbatovi* Schawaller, 2006, described from Myanmar, is newly recorded from the adjacent Assam.

Key words: Coleoptera, Tenebrionidae, Lagriinae, *Laena*, new species

## Zusammenfassung

Neue Arten der Gattung *Laena* Dejean, 1821 (Tenebrionidae: Lagriinae) werden aus Zentralindien (*L. centroindica* n. sp.), aus dem nordöstlichen Indien in Arunachal Pradesh und Assam (*L. bomdilaica* n. sp., *L. etalinica* n. sp., *L. gyatsoi* n. sp., *L. monpa* n. sp.), aus Myanmar (*L. mandalayica* n. sp., *L. schillhammeri* n. sp.), und von der Malayischen Halbinsel (*L. larutica* n. sp.) beschrieben. *L. bicolor* Schuster, 1926 ist erstmalig seit der Originalbeschreibung wiedergefunden worden, und *Laena kurbatovi* Schawaller, 2006, beschrieben aus Myanmar, wird erstmalig aus dem angrenzenden Assam gemeldet.

## Contents

1	Introduction .....	71
2	Species from central and north-eastern India .....	73
3	Species from Myanmar (Burma) .....	76
4	Species from the Malay Peninsula (West Malaysia) .....	77
5	References .....	78

## 1 Introduction

The wingless species of genus *Laena* Dejean, 1821 (Tenebrionidae: Lagriinae) are typical elements of the soil fauna of mountainous forests from southeastern Europe throughout continental Asia with usually restricted distributional patterns. The bulk of the species inhabit the Himalayan and Chinese mountain ranges. The adjacent parts of south and southeastern Asia (India, Thailand, Myanmar, Malay Peninsula, Laos and Vietnam, so far unknown from Cambodia) also harbour several species in isolated (and thus endangered) montane and lowland forests, none of them being conspecific with the Himalayan and Chinese species.

The species of southeastern Asia were summarised by SCHAWALLER (2006, 2009, 2012, 2014). These papers contain also full references of the previous publications, check lists and identification keys to species of certain countries, figures, and discussions about species characters and problems in defining natural species groups within

this speciose genus. In the meantime, several additional species were newly collected in central and northeastern India, Myanmar, and in the Malay Peninsula, mostly by sifting or with Winkler-extractors. These are described in the present contribution.


## Acronyms of depositories

ZFMK	Zoologisches Museum A. Koenig, Bonn, Germany
MHNG	Muséum d'Histoire Naturelle, Geneva, Switzerland
NHMB	Naturhistorisches Museum, Basel, Switzerland
NHMW	Naturhistorisches Museum, Wien, Austria
NMP	National Museum of Natural History, Prague, Czech Republic
SMNS	Staatliches Museum für Naturkunde, Stuttgart, Germany

## Acknowledgements

For the trustful and partly long-term loan of specimens from the collections under their care I thank

<sup>1</sup> Contributions to Tenebrionidae, no. 150. – For. no. 149 see: Integrative Systematics 1, 2018.



**Figs. 1–5.** Dorsal view of *Laena* species from southeastern Asia. – 1. *L. bomdilaica* n. sp., holotype NMP. 2. *L. fikaceki*, paratype SMNS. 3. *L. monpa* n. sp., holotype NMP. 4. *L. mandalayica* n. sp., holotype NHMW. 5. *L. schillhammeri* n. sp., holotype NHMW. – Scale: 2 mm.

DIRK AHRENS (Bonn), MICHEL BRANCUCCI (†) (Basel), GIULIO CUCCODORO (Geneva), JIRI HÁJEK (Prague), and HARALD SCHILLHAMMER (Vienna). The photographs were taken by JOHANNES REIBNITZ (Stuttgart) with a Leica DFC320 digital camera on a Leica MZ16 APO microscope and subsequently processed by him with Auto-Montage (Syncroscopy) software. The referees ROLAND GRIMM (Neuenbürg) and OTTÓ MERKL (Budapest) improved the manuscript by their remarks and corrections.

## 2 Species from central and north-eastern India

An identification key for all 28 species known so far from the eastern Himalayas (Darjeeling, Sikkim, Bhutan, Assam, Arunachal Pradesh) was compiled by SCHAWALLER (2012). From central India, isolated from the numerous species in northern India and Nepal, the genus was unknown until now.

### *Laena bomdilaica* n. sp. (Figs. 1, 16)

Holotype (♂): India, Arunachal Pradesh, 1.5 km NE Bomdila, Bomdila Pass, 2600 m, 2.–5.V.2008, leg. M. FIKÁČEK, H. PODSKALSKÁ & P. ŠIPEK, NMP.

**Description:** Body length 3.8 mm. Eyes not prominent. Pronotum with large, coarse confluent punctures, distance less than 1x puncture diameter, most punctures with short adpressed setae; surface convex and feebly shining, often dull by soil incrustation, disc uneven by confluent punctures; lateral margins completely unbordered; posterior corners rounded; propleura with similar but sparser punctation, and with shorter setation as on pronotum. Elytra ovate with rows of punctures, without striae, punctures of rows of similar size as punctures on pronotum, punctures with short adpressed setae; all intervals with fine punctures and similar adpressed setae, interval 9 in posterior part with 1 distinct setiferous pore; intervals 3, 5, and 7 distinctly convex, nearly keel-like with irregular tubercles, intervals 3 and 5 confluent in posterior part and joining to an prominent keel; intervals 1, 2, 4, and 6 nearly flat; surface feebly shining. Apex of elytra not prolonged. All femora without teeth or granules. Male tibiae without peculiarities. Aedeagus see Fig. 16.

**Diagnosis:** *Laena bomdilaica* n. sp. shares with *L. cuccodoroi* Schawaller, 2012 from Assam (Fig. 7) and *L. fikaceki* Schawaller, 2012 from Arunachal Pradesh (Fig. 2) the small body size, the unarmed femora, the unbordered lateral margins of the pronotum, and the general dorsal structure of the elytra with partly keel-like intervals and/or tubercles. However, in *L. bomdilaica* n. sp. the pronotum is widest nearly in the middle (widest in anterior third in *L. cuccodoroi* and *L. fikaceki*), the elytra are much broader, and the structure of the elytral intervals is different with partly keel-like intervals (only a few tubercles in *L. cuccodoroi*, and convex intervals with tubercles in *L. fikaceki*).

In *L. fikaceki* the aedeagus is quite different with a broad apicale and blunt tip, whereas the aedeagi of *L. bomdilaica* n. sp. and *L. cuccodoroi* are similar with triangular apicale and rounded tip (compare figures in SCHAWALLER 2012). *L. interrupta* Schuster, 1935 (Fig. 12) from Darjeeling belongs also in the same group of species.

**Etymology:** Named after the Bomdila Pass, where the holotype was collected.

### *Laena centroindica* n. sp. (Figs. 13, 17)

Holotype (♂): India, Maharashtra, 30 km E Phonda, Dhajipur Forest, 800 m, 2.XI.2000, leg. G. CUCCODORO, MHNG.

**Description:** Body length 2.5 mm. Eyes flat, not prominent. Pronotum with large, but not confluent punctures, distance 1–3x puncture diameters, punctures without setae; surface convex and shining, disc without impressions; lateral margins completely bordered; posterior corners rounded; propleura with similar punctation as on pronotum, and also without setation. Elytra ovate with rows of punctures, without striae, punctures of rows slightly larger than punctures on pronotum, punctures without setae; intervals without punctures and setation, interval 9 with 4 distinct setiferous pores, all intervals feebly convex, nearly flat; surface shining. Apex of elytra not prolonged. All femora without teeth or tubercles. Male tibiae without peculiarities. Aedeagus see Fig. 17.

**Diagnosis:** *Laena centroindica* n. sp. can be recognised by the combination of the following characters: Body size very small (2.5 mm), pronotum with bordered lateral margins, punctures of the elytral rows without setae, all femora without teeth or tubercles, and shape of the aedeagal apicale. No similar species is known from the bulk of species in northern India and Nepal that are separate from central India. A similar tiny species occurs in Thailand (*L. ellenae* Schawaller, 2014), with marked but unbordered lateral margins of pronotum, with long adpressed setae of elytra, and with different aedeagus. Because of the distributional patterns it seems unlikely that both have a closer relation.

**Etymology:** It is the first species of *Laena* from central India.

### *Laena etalinica* n. sp. (Figs. 10, 20)

Holotype (♂): India, Arunachal Pradesh, Etalin, 700 m, 12.–25.V.2012, leg. L. DEMBICKÝ, ZFMK.

Paratypes: Same data as holotype, 8 ex. ZFMK, 3 ex. SMNS. – India, Arunachal Pradesh, Hunli, 1200–1400 m, 26.V.–1.VI. 2012, leg. L. DEMBICKÝ, 6 ex. ZFMK, 2 ex. SMNS.

**Description:** Body length 3.5–6.0 mm. Eyes not prominent. Pronotum with large, not confluent punctures,





**Figs. 6–13.** Dorsal view of *Laena* species from southeastern Asia. – 6. *L. bicolor*, non-type Myanmar SMNS. 7. *L. cuccodoroi*, para-type SMNS. 8. *L. kurbatovi*, non-type Assam SMNS. 9. *L. larutica* n. sp., holotype MHNG. 10. *L. etalinica* n. sp., holotype ZFMK. 11. *L. gyatsoi* n. sp., holotype NMP. 12. *L. interrupta*, non-type Darjeeling SMNS. 13. *L. centroindica* n. sp., holotype MHNG. – Scale: 2 mm.

distance 1–3x puncture diameters, most punctures with long erect setae; surface convex and shining, disc without impressions; lateral margins unbordered; posterior corners rounded; propleura with similar punctation as on pronotum, with a few adpressed microsetae. Elytra ovate with rows of punctures, without striae, punctures of rows of similar size as punctures on pronotum, punctures with long erect setae; intervals with row of sparse small punctures, punctures with long erect setae, interval 9 with 4 distinct setiferous pores, all intervals feebly convex; surface shining. Apex of elytra not prolonged. All femora without teeth or angulations. Posterior tibiae of male somewhat swollen at inner side. Aedeagus see Fig. 20.

**Diagnosis:** In SCHAWALLER'S (2012) key to the species of the eastern Himalayas *Laena etalinica* n. sp. would run to *L. laevipennis* Schuster, 1926 from Sikkim and Darjeeling. Both share the general body size and shape, the unarmed femora, the elytra with punctural rows without striae, and mainly the high convex pronotum with completely unbordered lateral margins. However, in *L. laevipennis* the pronotum is finely and sparsely punctate, the punctures of the elytral rows bear short adpressed setae, and the elytral intervals are completely flat. Unfortunately, the aedeagus of *L. laevipennis* is unknown. Etymology: Named after the village Etalin, in whose vicinity the holotype was collected.

*Laena gyatsoi* n. sp.  
(Figs. 11, 19)

**Holotype** (♂): India, Arunachal Pradesh, 0.7 km W Tawang, 1–1.7 km N monastery, 2950 m, 27.–30.IV.2008, leg. M. FIKÁČEK, H. PODSKALSKÁ & P. ŠÍPEK, NMP.

**Description:** Body length 5.0 mm. Eyes not prominent. Pronotum with large, not confluent punctures, distance 1–2x puncture diameters, most punctures with short adpressed setae; surface convex and shagreened, disc medially with pair of very weak impressions; lateral margins somewhat marked but unbordered; posterior corners rounded; propleura with larger and sparser punctures as on pronotum, without setation. Elytra ovate with rows of punctures, without striae, punctures of rows of similar size as punctures on pronotum, punctures with few adpressed microsetae; intervals with row of sparse small punctures, punctures with adpressed microsetae, interval 9 with 3 distinct setiferous pores, all intervals flat; surface shagreened. Apex of elytra not prolonged. All femora with pair of angulations, on anterior femora more prominent than on middle and posterior femora. Male tibiae without peculiarities. Aedeagus see Fig. 19.

**Diagnosis:** *Laena gyatsoi* n. sp. is similar to *L. holzschuhi* Schawaller, 2012 from Bhutan, and *L. vishnua* Masumoto, 1990 from Darjeeling. They share the general body size and dorsal punctation, short or lacking dor-

sal setation, and the femora with more or less developed angulations. However, in *L. gyatsoi* n.sp. the lateral margins of the pronotum are at mostly marked but unbordered, and the pronotal punctures are large and dense (distinctly bordered and punctures sparser and finer in *L. holzschuhi* and *L. vishnua*). The aedeagi are also quite different with a narrow finger-like apicale in *L. holzschuhi*, or broader finger-like apicale in *L. vishnua* (compare with figures in SCHAWALLER 2012).

**Etymology:** Named in honour of Lama LODRE GYATSO, who founded in 1680–1681 the Tawang Monastery, in whose vicinity the holotype was collected.

*Laenak urbatovi* Schawaller, 2006  
(Figs. 8, 22)

**Specimens examined:** India, Assam, Kaziranga, 75 m, 7.–9.V.1976, leg. C. BARONI-URBANI & W. WITTMER, 1 ♂ 1 ♀ NHMB, 1 ♂ SMNS.

**Remarks:** Body length 3.8–5.2 mm. These specimens from Assam show no external characters different from the single female holotype (thus aedeagus unknown so far), described from the Chin State in Myanmar which lies somewhat south of the Indian Assam. *L. kurbatovi* shares with *L. kenyirica* Schawaller, 2006 and *L. larutica* n. sp. (both from Malaysia) the presence of tooth-like setiferous pores around elytral shoulders.

*Laena monpa* n. sp.  
(Figs. 3, 21)

**Holotype** (♂): India, Arunachal Pradesh, 1.5 km NE Bomdila, Bomdila Pass, 2600 m, 2.–5.V.2008, leg. M. FIKÁČEK, H. PODSKALSKÁ & P. ŠÍPEK, NMP.

**Paratype:** Same data as holotype, 1 ♀ SMNS.

**Description:** Body length 3.5–3.7 mm. Eyes not prominent. Pronotum with fine, scattered punctures, distance 1–4x puncture diameters, most punctures with long erect setae; surface convex and shining, feebly shagreened, disc without impressions; lateral margins not completely bordered, border diminishing near anterior corners; posterior corners rounded; propleura with larger and sparser punctures as on pronotum, without setation. Elytra ovate with rows of punctures, without striae, punctures of rows of similar size as punctures on pronotum, punctures on disc without setae, laterally and distally with a few punctures with longer erect setae; intervals with a row of sparse fine punctures, without setae, interval 9 with 3 indistinct setiferous pores, all intervals flat; surface shining. Apex of elytra not prolonged. All femora without teeth or granules. Tibiae of both sexes without peculiarities. Aedeagus see Fig. 21.

**Diagnosis:** In SCHAWALLER'S (2012) key to the species of the eastern Himalayas *Laena monpa* n. sp. would run to the group of species *L. esagnoli* Kaszab, 1965,

*L. aenea* Schuster, 1926 and *L. affinis* Schuster, 1935, all from Sikkim and Darjeeling. These species share the unarmed femora, the bordered lateral margins of the pronotum, and the short dorsal setation. However, *L. monpa* n. sp. can be recognised by the round and high convex pronotum with shining surface, by not prominent eyes, and by the shape of the aedeagus. In *L. aenea* and *L. affinis*, the dorsal surface is distinctly shagreened, the eyes are prominent, and the aedeagi are different; in *L. espanyoli* the surface is shining and the eyes are not prominent as in *L. monpa* n. sp., however the pronotum is less convex and the aedeagus is completely different.

**Etymology:** Named after the Monpa, the major ethnic tribe of native people in Arunachal Pradesh.

### 3 Species from Myanmar (Burma)

An identification key and distribution map for all five species known so far from Myanmar (Burma) was published by SCHAWALLER (2009).


### *Laena bicolor* Schuster, 1926 (Fig. 6)

**Specimens examined:** Myanmar, Mandalay Region, Mogok, S Panlin, Mt. Taung Mae, west slope, 1870 m, 17.–24.V.2016, leg. H. SCHILLHAMMER, A. BRUNKE, J. JENKINS-SHAW & A. R. JENSEN, 1 ♀ NHMW, 1 ♀ SMNS.

**Remarks:** This species was known so far only by the single holotype of unknown sex. The type locality „Birma: Rubi Mines“ was not exactly localised, but lies probably in the Mandalay Region (SCHAWALLER 2006). Unfortunately, both new specimens are females, thus the aedeagus can not be figured here.

### *Laena mandalayica* n. sp. (Figs. 4, 14)

**Holotype (♂):** Myanmar, Mandalay Region, Mogok, S Panlin, Mt. Taung Mae, west slope, 1870 m, 17.–24.V.2016, leg. H. SCHILLHAMMER, A. BRUNKE, J. JENKINS-SHAW & A. R. JENSEN, NHMW.



**Figs. 14–22.** Aedeagus of *Laena* species from southeastern Asia. – 14. *L. mandalayica* n. sp., holotype NHMW. 15. *L. schillhammeri* n. sp., holotype NHMW. 16. *L. bomdilaica* n. sp., holotype NMP. 17. *L. centroindica* n. sp., holotype MHNG. 18. *L. larutica* n. sp., holotype MHNG. 19. *L. gyatsoi* n. sp., holotype NMP. 20. *L. etalinica* n. sp., holotype ZFMK. 21. *L. monpa* n. sp., holotype NMP. 22. *L. kurbatovi*, non-type Assam SMNS. – Scale: 1 mm.



Paratypes: Same data as holotype, 3 ex. NHMW, 1 ex. SMNS.

**Description:** Body length 7.0–8.0 mm. Eyes not prominent. Pronotum with large, somewhat irregular, not confluent punctures, distance 1–2x puncture diameters, some punctures with short adpressed microsetae; surface convex and feebly shagreened, disc medially with a pair of weak impressions; lateral margins completely bordered; posterior corners rounded; propleura with larger and sparser punctures as on pronotum, without setation. Elytra ovate with rows of punctures, without striae, punctures of rows of similar size as punctures on pronotum, punctures without setae; intervals without punctures and setation, interval 9 with 3 distinct setiferous pores, all intervals regularly convex; surface feebly shagreened. Apex of elytra not prolonged. All femora without teeth or granules. Tibiae of both sexes without peculiarities. Aedeagus see Fig. 14.

**Diagnosis:** *Laena mandalayica* n. sp. is characterised by the following characters: Large body size, shining surface without distinct setation, convex pronotum with bordered lateral margins, convex elytral intervals, and unarmed femora. This character combination is not present in any other species of the area including adjacent Arunachal Pradesh and Yunnan. *L. kachinorum* Schawaller, 2009 from northern Myanmar is also large (9.2–9.6 mm), but the pronotum is less convex and the surface is shagreened, the elytral intervals are flat and distinctly shagreened, the male tibiae medially bear a row of distinct spines, and the aedeagus is different having a short and broad apicale with blunt tip.

**Etymology:** Named after Mandalay, where the type series was collected.

*Laena schillhammeri* n. sp.  
(Figs. 5, 15)

**Holotype** (♂): Myanmar, Mandalay Region, Mogok, S Panlin, Mt. Taung Mae, west slope, 1870 m, 17.–24.V.2016, leg. H. SCHILLHAMMER, A. BRUNKE, J. JENKINS-SHAW & A. R. JENSEN, NHMW.

**Paratypes:** Same data as holotype, 10 ex. NHMW, 2 ex. SMNS. – Myanmar, Mandalay Region, Mogok, S Panlin, Mt. Taung Mae, west slope, 1710–1750 m, 10.–18.VI.2014, leg. A. BRUNKE & H. SCHILLHAMMER, 17 ex. NHMW, 3 ex. SMNS.

**Description:** Body length 4.8–7.0 mm. Eyes slightly prominent. Pronotum with large, but not confluent punctures, distance 1–4x puncture diameters, most punctures with longer erect setae; surface convex and feebly shagreened, disc without impressions; lateral margins completely bordered; posterior corners rounded; propleura with similar but sparser punctures as on pronotum, but without setation. Elytra ovate with rows of punctures in feeble striae, punctures of rows of similar size as punctures on pronotum, most punctures with longer and erect

setae; intervals with a few fine punctures, each bearing an erect seta of equal length, interval 9 with 3 indistinct setiferous pores, all intervals feebly convex, nearly flat; surface feebly shagreened. Apex of elytra not prolonged. All femora without teeth or tubercles. Tibiae of both sexes without peculiarities. Aedeagus see Fig. 15.

**Diagnosis:** *Laena schillhammeri* n. sp. is similar to *Laena kurbatovi* Schawaller, 2006, also from the Chin State in central Myanmar (see above). Both share the general body shape, the long dorsal setation, the bordered lateral margins of the pronotum, and the unarmed femora. However, in *L. kurbatovi*, the pronotum is widest in the middle (widest in anterior part in *schillhammeri* n. sp.), the elytral intervals are distinctly convex (nearly flat in *schillhammeri* n.sp.), and the elytral shoulders are dentate by a higher number of setiferous pores (rounded with few pores in *schillhammeri* n.sp.). Probably, *L. kurbatovi* is also smaller (3.7 mm in the single ♀ holotype) than *L. schillhammeri* n. sp. (4.8–7.0 mm in the type series). The aedeagi are also different with a broad spade-like apicale in *L. schillhammeri* n.sp., and with a short and broad apicale in *L. kurbatovi* (Fig. 22, non-type ♂ from Assam, the holotype from Myanmar is a ♀).

**Etymology:** Named in honor of Dr. HARALD SCHILLHAMMER (Wien), one of the collectors of the type series.

#### 4 Species from the Malay Peninsula (West Malaysia)

The last identification key for all 18 species known so far from the Malay Peninsula was compiled by SCHAWALLER (2014).

*Laenal arutica* n. sp.  
(Figs. 9, 18)

**Holotype** (♂): West Malaysia, Perak, Maxwell Hill, 950 m, 22.XI.1999, leg. G. CUCCODORO & I. LÖBL, MHNG.

**Paratype:** West Malaysia, Perak, Maxwell Hill, 1150 m, 25.XI.1999, leg. G. CUCCODORO & I. LÖBL, 1 ex. SMNS.

**Description:** Body length 3.8–5.0 mm. Eyes normal, not prominent. Pronotum with large, partly confluent punctures, distance 0.5–2x puncture diameters, most punctures with longer erect setae; surface convex and feebly shagreened, disc without impressions; lateral margins irregularly bordered; posterior corners angulate; propleura with sparser punctation as on pronotum, and with shorter setation. Elytra ovate with rows of punctures in feeble striae, punctures of rows of similar size as punctures on pronotum, most punctures with longer and erect setae; intervals with a few fine punctures, each bearing an erect seta of equal length, interval 7 with distinct nearly tooth-like setiferous pore at shoulders, interval 9 with 3 distinct setiferous pores, basal one also nearly tooth-like,



all intervals distinctly convex, surface feebly shagreened. Apex of elytra not prolonged. All femora without teeth or tubercles. Tibiae of both sexes without peculiarities. Aedeagus see Fig. 18.

**Diagnosis:** In the key of the Malayan species by SCHAWALLER (2014), *Laena larutica* n. sp. would run to *L. kenyirica* Schawaller, 2006 from lowland forest in the northeastern part of the peninsula. Both share the general body size and shape, the long and erect dorsal setation, the pronotum widest in anterior third, the presence of tooth-like setiferous pores around elytral shoulders, the convex elytral intervals, and unarmed femora. However, in *L. kenyirica* the pronotum is flatter, its punctation is sparser, and the lateral margin is regularly bordered. The aedeagus is also quite different with short and broad, nearly quadrate apicale with blunt tip. *L. tanahratica* Schawaller, 2014 from the Cameron Highlands belongs to the same group, but in this species, besides other characters, the pronotum is widest in the middle, and the

aedeagus is also different with narrow, finger-like apicale. **Etymology:** Named after Bukit Larut, the modern name of the Maxwell Hill, where the type series was collected.

## 5 References

- SCHAWALLER, W. (2006): New species of the genus *Laena* Latreille (Coleoptera: Tenebrionidae) from southeastern Asia. – *Zootaxa* **1325**: 369–384.
- SCHAWALLER, W. (2009): Three new species of *Laena* Latreille from Upper Burma (Myanmar) (Coleoptera: Tenebrionidae). – *Koleopterologische Rundschau* **79**: 267–272.
- SCHAWALLER, W. (2012): The genus *Laena* Dejean (Coleoptera: Tenebrionidae) in the eastern Himalayas (Darjeeling, Sikkim, Bhutan, Assam, Arunachal Pradesh), with descriptions of seven new species. – *Stuttgarter Beiträge zur Naturkunde A (NS)* **5**: 257–269.
- SCHAWALLER, W. (2014): New species of the genus *Laena* Dejean (Coleoptera: Tenebrionidae) from southeastern Asia (2). – *Annales Zoologici* **64**: 287–298.

Author's address:

Dr. WOLFGANG SCHAWALLER, Staatliches Museum für Naturkunde, Rosenstein 1, 70191 Stuttgart, Germany;  
e-mail: schawaller.ehrenamt@smns-bw.de

Manuscript received: 27.IX.2017. , accepted: 4.XII.2017