



A Triplex DMA Structure of the Polypyrimidine: Polypurine Stretch in the 5' Flanking Region of the Sea Urchin Arylsulfatase Gene

Authors: Sakamoto, Naoaki, Akasaka, Koji, Yamamoto, Takashi, and Shimada, Hiraku

Source: Zoological Science, 13(1) : 105-109

Published By: Zoological Society of Japan

URL: <https://doi.org/10.2108/zsj.13.105>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

A Triplex DNA Structure of the Polypyrimidine: Polypurine Stretch in the 5' Flanking Region of the Sea Urchin Arylsulfatase Gene

Naoaki Sakamoto¹, Koji Akasaka¹, Takashi Yamamoto²
and Hiraku Shimada¹

¹Laboratory of Molecular Genetics, Graduate Department of Gene Science,
Faculty of Science, Hiroshima University, Higashi-Hiroshima 739
and ²Department of Biology, Faculty of Science, Kumamoto
University, Kurokami, Kumamoto 860, Japan

ABSTRACT—Previously we reported that a long (522 bp) polypyrimidine: polypurine stretch in the 5' flanking region of the arylsulfatase gene of the sea urchin, *Hemicentrotus pulcherrimus*, took an unusual, perhaps triplex, DNA structure, when subjected to an acidic pH (pH 5) (Yamamoto *et al.*, 1994). In the present study we have isolated a polypyrimidine: polypurine containing fragment from the arylsulfatase gene and surveyed the sensitivities of the polypyrimidine: polypurine stretch to base modification by diethylpyrocarbonate and osmium tetroxide under various levels of negative supercoiling. Based on the sensitivity of highly negatively supercoiled DNA to these base-modifying reagents, we conclude that, when highly negatively supercoiled, the polypyrimidine: polypurine stretch can take a triplex DNA structure even at a neutral pH and under physiological ionic strength in the presence of Mg²⁺.

INTRODUCTION

Eukaryotic genomes often contain polypyrimidine: polypurine sequences that, in many cases, are located in or around transcription-regulating regions of genes (Mirkin *et al.*, 1987; Sinden, 1994; Wells *et al.*, 1988) and are thus believed to be related to regulation on transcription. Though little evidence is available on the role of these polypyrimidine stretches in gene transcription, many investigators report that a polypyrimidine sequence inserted into a plasmid can take an unusual DNA structure distinct from a usual B-structure (Hentschel, 1982; Htun *et al.*, 1984; Kato *et al.*, 1990; Kohwi-Shigematsu and Kohwi, 1985; Mace *et al.*, 1983; Nickol and Felsenfeld, 1983; Pulleyblank *et al.*, 1985; Schon *et al.*, 1983). Mirkin *et al.* (1987) and Wells *et al.* (1988) showed that a region with a mirror-repeat sequence within a polypyrimidine stretch forms a triplex DNA structure through Hoogsteen base pairings, and stability of triplex DNA structure is dependent on the degree of negative supercoiling of the plasmid as well as pH and ionic strength of the DNA solution (Hanvey *et al.*, 1988; Htun and Dahlberg, 1988; Mirkin *et al.*, 1987; Pulleyblank *et al.*, 1985; Wells *et al.*, 1988).

The arylsulfatase (Ars) gene of the sea urchin, *Hemicentrotus pulcherrimus*, which is expressed in a developmental stage-specific (Sasaki *et al.*, 1988) as well as tissue-specific manner (Akasaka *et al.*, 1990), contains a long polypyrimidine:polypurine (Pyr-Pur) stretch in its 5' flanking region (Akasaka *et al.*, 1994) and in its 1st intron (unpublished

data). Because the DNA region with an unusual structure may have a specific function in regulation of transcription, we have undertaken a series of studies to explore the possible structure of the polypyrimidine stretch in the sea urchin Ars gene. Through an S1 nuclease assay at pH 5, we previously reported that a part of the polypyrimidine stretch of the sea urchin Ars gene can take an unusual DNA structure different from B-DNA, when the stretch is in a negatively supercoiled plasmid (Yamamoto *et al.*, 1994). However, the reaction conditions such as pH 5 and a low ionic strength in an S1 nuclease assay obviously do not reflect *in vivo* conditions.

To approach to the question of the biological function of Pyr-Pur region in the 5' flanking region of the Ars gene, DNA structure of the Pyr-Pur region has been re-examined by measuring sensitivities of the Pyr-Pur region to base modification by osmium tetroxide and diethylpyrocarbonate under physiological ionic strength at neutral pH and in the presence of 5 mM Mg²⁺. The results suggest that, if the stretch is highly negatively supercoiled, a part of the polypyrimidine stretch takes triplex DNA structure even under physiological conditions.

MATERIALS AND METHODS

Plasmid DNA

A region containing 522 bp Pyr-Pur stretch (−2, 201 to −1, 680) of the Ars gene of the sea urchin (*Hemicentrotus pulcherrimus*) was excised and inserted into pBluescript SK− (pBSK) vector (Stratagene Cloning Systems, USA) to construct the plasmid pBSK-AvNs (Fig. 1). Various degrees of negative supercoiling were introduced into pBSK-AvNs by treating 20 µg/ml of pBSK-AvNs with topoisomerase I at 37°C for 1 hr in the presence of various concentrations of ethidium bromide. After extracting the supercoiled

Accepted November 18, 1995

Received September 25, 1995




Fig. 1. Restriction sites around the Pyr-Pur region in the Ars gene and in the pBSK-AvNs. Slashed box; the Pyr-Pur region, open boxes; exons, a thick line; 5' flanking region and introns, a thin line; pBSK vector. A; *Apal*, H; *HindIII*, N; *Nsp7524I*, Av; *Aval*, E; *EcoRI*.

DNA by phenol/chloroform, ethidium bromide was removed by water-saturated *n*-butanol, and DNA was precipitated and washed with ethanol, and dried *in vacuo*.

The density of negative supercoiling was determined by two dimensional electrophoresis (Pulleyblank *et al.*, 1985). A mixture of variously supercoiled plasmids was first electrophoresed on 1.5% agarose in 0.5×TBE, then gels were equilibrated with 1 µg/ml chloroquine in 0.5×TBE. The second electrophoresis was performed in the presence of 1 µg/ml chloroquine in 0.5×TBE. The DNA bands were detected by southern blotting with digoxigenin-labeled pBSK-AvNs as the probe.

Diethylpyrocarbonate (DEPC) and osmium tetroxide reactions

Modification of purine bases (especially adenine) by DEPC and pyrimidine bases (especially thymine) by osmium tetroxide in the negatively supercoiled pBSK-AvNs were carried out as described by Hanvey *et al.* (1988) with some modifications.

Three µg of negatively supercoiled pBSK-AvNs was dissolved in 100 µl of 0.3 M Tris-HCl buffer (pH 7.5) containing 5 mM MgCl₂ and 0.25 M KCl, and incubated for 1 hr at 13°C. The DEPC reaction was started by adding 10 µl of DEPC to the mixture, followed by 1-min of vortexing and a 30-min incubation at 22°C. The osmium tetroxide reaction was started by addition of 2 µl of pyridine and 10 µl of 20 mM OsO₄ to the mixture, followed by 1-min vortexing and 30-min incubation at 22°C. The plasmid DNA was then cleaved at modified nucleotides by piperidine, and sites of modification were determined on a denaturing polyacrylamide gel.

RESULTS AND DISCUSSION

Electrophoretic mobility of negatively supercoiled pBSK-AvNs

pBSK-AvNs was negatively supercoiled by DNA topoisomerase I treatment in the presence of ethidium bromide and electrophoresed on 1.5% agarose gel containing TBE buffer at pH 8.2. Increasing the concentration of ethidium bromide, increased the electrophoretic mobility of pBSK-AvNs (Fig. 2), indicating that the level of negative supercoiling of pBSK-AvNs is positively dependent on the concentration of ethidium bromide in the topoisomerase reaction. Electrophoretic mobility of the plasmid decreased by increasing ethidium bromide from 2 µg/ml to 10 µg/ml, suggesting that the high negative supercoiling ($\sigma < -0.0385$, as determined in the presence of chloroquine by two




Fig. 2. Electrophoretic mobility of pBSK-AvNs changes depending on degree of negative supercoiling. Various degrees of negative supercoiling were produced in pBSK-AvNs by treating the plasmid with topoisomerase I in the presence of various concentrations of ethidium bromide. Electrophoresis was carried out in 1.5% agarose gel. Concentration of ethidium bromide, lane 2; without ethidium bromide, lane 3; 0.2 µg/ml, lane 4; 0.4 µg/ml, lane 5; 0.6 µg/ml, lane 6; 0.8 µg/ml, lane 7; 1.0 µg/ml, lane 8; 1.2 µg/ml, lane 9; 1.4 µg/ml, lane 10; 2 µg/ml, lane 11; 5 µg/ml, lane 12; 8 µg/ml, lane 13; 10 µg/ml. Lanes 1 and 14; pBSK-AvNs before treatment with topoisomerase I.

dimensional electrophoresis, see Materials and Methods) causes a severe conformational change of the plasmid that leads to reduction of the electrophoretic mobility. A plasmid with this level of negative supercoiling is hereafter referred to as highly negatively supercoiled, with the supercoiling density of $\sigma = -0.0357$ as a native level of supercoiling.

Chemical modification of bases in negatively supercoiled pBSK-AvNs

It is known that diethylpyrocarbonate (DEPC) carboxy-ethylates adenines at their N7 and cleaves an imidazole ring, while OsO₄ reacts with thymines at their C5 and C6 to form osmate esters, when DNA is in a non-B structure (Johnston, 1988; Vojtiskova and Palecek, 1987; Wells *et al.*, 1988). Thus the sensitivity of DNA to these reagents is a good measure to estimate its structure.

Modification by DEPC and OsO₄ of nucleotide bases in a linearized pBSK-AvNs, a native pBSK-AvNs (with a native level of supercoiling) and a highly negatively supercoiled pBSK-AvNs was surveyed under physiological ionic strength and Mg²⁺ concentration similar to *in vivo* levels. The experimental results are shown in Figure 3, and the positions of modified bases are summarized schematically in Figure 4.

Clustered adenines in the purine strand of a linear DNA were chemically modified while any bases in the pyrimidine strand were not modified (Fig. 4a). At present we do not have any evidence to explain increased accessibility of DEPC to adenines in linear DNA fragments, but it is conceivable that consecutive adenines tend to cause DNA bending (Sinden, 1994) that would make adenine bases accessible to DEPC.

In the case of a native pBSK-AvNs (Fig. 4b), though the purine strand was modified by DEPC in a manner similar to




Fig. 3. Base modification of Pyr-Pur region in negatively supercoiled plasmid by DEPC or by OsO_4 . pBSK-AvNs in linear, native and highly negatively supercoiled forms were treated with DEPC or with OsO_4 , cleaved by *Hind*III, labeled with ^{32}P at the 3' end of the pyrimidine strand, and the labeled *Hind*III-*Apal* fragments were isolated (a and b). In the case of preparation of the ^{32}P -labeled *Hind*III-*Apal* fragments from the purine strand (c and d), the plasmid was first cut by *Apal*, and the labeled fragment was prepared in a similar way. The labeled fragments were cleaved by piperidine at modified bases and electrophoresed on 4% denaturing polyacrylamide gel (a and b) and on 6% denaturing polyacrylamide gel (c and d). Sites of $(\text{CT})_{11}$ or $(\text{GA})_{11}$ were indicated by vertical lines on the right side of each column. Lanes 1 and 2; Maxam-Gilbert ladders or A+G and C in (a) and (b) and C+T and G in (c) and (d), respectively. Lane 3; linearized pBSK-AvNs, lane 4; native pBSK-AvNs, lane 5; highly negatively supercoiled pBSK-AvNs. (a) DEPC-sensitive sites in the pyrimidine strand. (b) OsO_4 -sensitive sites in the pyrimidine strand. (c) DEPC-sensitive sites in the purine strand. (d) OsO_4 -sensitive sites in the purine strand.

a linear plasmid, thymines in the pyrimidine strand that were not accessible to OsO_4 in a linear plasmid became accessible to the reagent. It is of interest that, while the $(\text{CT})_{11}$ tract was extensively modified by OsO_4 , the complementary $(\text{GA})_{11}$ tract was not modified by DEPC, and thymine stretches flanking both ends of the $(\text{CT})_{11}$ tract were protected from modification. This result suggests that even a native level of negative supercoiling can produce a conformational change in the Pyr-Pur region which results in a DNA structure slightly deviated from B-form.

By imposing higher negative supercoiling than native level, the accessibility of base-modifying reagents to the plasmid increased in the purine strand. As depicted in Figure 4c, while little change was observed in the thymine modification of the pyrimidine strand as compared with a plasmid with a native level of supercoiling, a marked change in adenine modification was observed in a restricted region

of the purine strand. Adenines in the 3' half of $(\text{GA})_{11}$ in the purine strand were modified and the adenines upstream of $(\text{GA})_{11}$ (from -1,827 to -1,807) became more accessible to DEPC. That is, adenines between -1,827 and -1,793 were all modified, indicating the single-stranded nature of the purine strand of this region under high negative supercoiling.

A triplex DNA structure of the Pyr-Pur region

A triplex DNA structure through intramolecular formation of Hoogsteen base-pairings (Hoogsteen, 1963) seems to be a most plausible DNA model that accommodates most of the above experimental results on the accessibility of the Pyr-Pur region to base-modifying reagents. A possible model of a triplex DNA structure taken by the Pyr-Pur region of the sea urchin Ars gene is depicted in Figure 5. In this model, the purine strand between -1,829 and -1,787 is single-stranded, while the pyrimidine counterpart winds up around




Fig. 4. Mapping of sites of base modification in *Hind*III-*Apal* fragments in linear, native and highly negatively supercoiled plasmids. The sites of base modification determined in experiments in Fig. 3 are mapped in the nucleotide sequence of AvNs fragments. Sites of base modification in (a) a linearized plasmid, (b) a native plasmid, and (c) a highly negatively supercoiled plasmid. Thin and short arrows indicate weakly sensitive sites, and thick and long arrows indicate highly sensitive sites. Very thick horizontal arrows indicate a mirror repeat sequence.




Fig. 5. A possible H-DNA model formed in polypyrimidine region of the sea urchin *Ars* gene. Small arrows indicate bases weakly accessible to base-modifying reagents when the polypyrimidine region is highly negatively supercoiled, and large arrows indicate bases highly accessible to the reagents.

the upstream double helical DNA along its major groove to form a triple-stranded DNA structure. Thymine in the third pyrimidine strand of this structure would be eventually protected from attack by OsO_4 .

The existence of a mirror repeat sequence (from -1,849 to -1,804; Fig. 4c) in the Pyr-Pur region supports this model. If strand separation occurs in the right half of the mirror repeat accompanied by folding back of its single-stranded pyrimidine counterpart along a major groove of the left half of the mirror repeat, 21 Hoogsteen base-pairings containing C-G-C and T-A-T would be formed between the purine strand of the double helix and the pyrimidine strand that folded back over it, as suggested by Wells *et al.* (1988). In spite of the presence of Mg^{2+} , H-DNA structure appears to be formed through C-G-C and T-A-T in the Pyr-Pur region of the sea urchin *Ars* gene, while Kohwi and Kohwi-Shigematsu (1988) showed that in the presence of Mg^{2+} an isomer of H-DNA is formed that contains C-G-G in place of C-G-C. Alternately-arranged T-A-T's may stabilize the C-G-C structure.

Biological significance of a triplex DNA structure is not clear though its possible functions in transcription, replication and recombination have been discussed (Sinden, 1994). Michel *et al.* (1993) suggested that a triplex DNA region may act as a transcriptional insulator by insulating some genetic loci from influences of other genes through its unusual steric conformation, while Baran *et al.* (1991) showed that DNA synthesis is arrested at sites of DNA triplex formation. It is also possible that sharp bends of DNA in these structures may help interaction of protein factors that bind to distantly separated *cis*-elements.

ACKNOWLEDGMENTS

A part of this work was carried out in the Center for Gene Research of Hiroshima University. This work was supported in part by Grants-in-Aid for Co-operative Research (A) (No. 04304007), for Scientific Research (B) (No. 05454653) and for Scientific Research on Priority Areas (No. 05277208 and 06270213) to H.S., and for Scientific Research (C) (No. 05680642 and No. 06680719) to K.A., from the Ministry of Education, Science, Sports and Culture, Japan. This research was also supported in part by a Grant Pioneering Research Project in Biotechnology given to H.S. by the Ministry of Agriculture, Forestry and Fisheries, Japan. Research support by a Grant from The Mitsubishi Foundation to H.S. is also acknowledged.

REFERENCES

- Akasaka K, Sakamoto N, Yamamoto T, Morokuma J, Fujikawa N, Takata K, Eguchi S, Shimada H (1994) Corrected structure of the 5' flanking region of arylsulfatase gene of sea urchin, *Hemientrotus pulcherrimus*. *Dev Growth Differ* 36: 633-636
- Akasaka K, Ueda T, Higashinakagawa T, Yamada K, Shimada H (1990) Spatial patterns of arylsulfatase mRNA expression in sea urchin embryo. *Dev Growth Differ* 32: 9-13
- Baran N, Lapidot A, Manor H (1991) Formation of DNA triplexes accounts for arrests of DNA synthesis at $d(\text{TC})_n$ and $d(\text{GA})_n$ tracts. *Proc Natl Acad Sci USA* 88: 507-511
- Harvey JC, Klysik J, Wells RD (1988) Influence of DNA sequence on the formation of non-B right-handed helices in oligopurine:oligopyrimidine inserts in plasmids. *J Biol Chem* 263: 7386-7396
- Hentschel CC (1982) Homocopolymer sequences in the spacer of a sea urchin histone gene repeat are sensitive to S1 nuclease. *Nature* 295: 714-716
- Hoogsteen K (1963) The crystal and molecular structure of a hydrogen-bonded complex between 1-methylthymine and 9-methyl-adenine. *Acta crystallogr* 16: 907-916
- Htun H, Lund E, Dahlberg JE (1984) Human U1 RNA genes contain an unusually sensitive nuclease S1 cleavage site within the conserved 3' flanking region. *Proc Natl Acad Sci USA* 81: 7288-7292
- Htun H, Dahlberg JE (1988) Single strands, triple strands, and kinks in H-DNA. *Science* 241: 1791-1796
- Johnston, BH (1988) The S1-sensitive form of $d(\text{C-T})_n \cdot d(\text{A-G})_n$: Chemical evidence for a three-stranded structure in plasmids. *Science* 241: 1800-1804
- Kato M, Kudoh J, Shimizu N (1990) The pyrimidine/purine-biased region of the epidermal growth factor receptor gene is sensitive to S1 nuclease and may form an intramolecular triplex. *Biochem J* 268: 175-180
- Kohwi-Shigematsu T, Kohwi Y (1985) Poly(dG)-poly(dC) sequences, under torsional stress, induce an altered DNA conformation upon neighboring DNA sequence. *Cell* 43: 199-206
- Kohwi Y, Kohwi-Shigematsu T, (1988) Magnesium ion-dependent triple-helix structure formed by homopurine-homopyrimidine sequences in supercoiled plasmid DNA. *Proc Natl Acad Sci USA* 85: 3781-3785
- Mace HAF, Pelham HRB, Travers AA (1983) Association of an S1 nuclease-sensitive structure with short direct repeats 5' of *Drosophila* heat shock genes. *Nature* 304: 555-557
- Michel D, Chatelain G, Herault Y, Harper F, Brun G (1993) H-DNA can act as a transcriptional insulator. *Cell Mol Biol Res* 39: 131-140
- Mirkin SM, Lyamichev VI, Drushlyak KN, Dobrynin VN, Filippov SA, Frank-Kamenetskii MD (1987) DNA H form requires a homopurine-homopyrimidine mirror repeat. *Nature* 330: 495-497
- Nickol JM, Felsenfeld G (1983) DNA conformation at the 5' end of the chicken adult β -globin gene. *Cell* 35: 467-477
- Pulleyblank DE, Haniford DB, Morgan AR (1985) A structural basis for S1 nuclease sensitivity of double-stranded DNA. *Cell* 42: 271-280
- Sasaki H, Yamada K, Akasaka K, Kawasaki H, Suzuki K, Saito A, Sato M, Shimada H (1988) cDNA cloning, nucleotide sequence and expression of the gene for arylsulfatase in the sea urchin (*Hemicentrotus pulcherrimus*) embryo. *Eur J Biochem* 177: 9-13
- Schon E, Evans T, Welsh J, Efstratiadis A (1983) Conformation of promoter DNA: Fine Mapping of S1-hypersensitive sites. *Cell* 35: 837-848
- Sinden RR (1994) DNA Structure and Function. Academic Press, USA
- Vojtiskova M, Palecek E (1987) Unusual protonated structure in the homopurine: homopyrimidine tract of supercoiled and linearized plasmids recognized by chemical probes. *J Biomol Struct Dynam* 5: 283-296
- Wells RD, Collier DA, Harvey JC, Shimizu M, Wohlrab F (1988) The chemistry and biology of unusual DNA structures adopted by oligopurine-oligopyrimidine sequences. *FASEB J* 2: 2939-2949
- Yamamoto T, Akasaka K, Irie S, Shimada H (1994) A long polypyrimidine: polypurine sequence in 5' flanking region of arylsulfatase gene of sea urchin embryo. *Int J Dev Biol* 38: 337-343