



Japanese Eel: A Model for Analysis of Spermatogenesis

Authors: Miura, Takeshi, and Miura, Chiemi

Source: Zoological Science, 18(8) : 1055-1063

Published By: Zoological Society of Japan

URL: <https://doi.org/10.2108/zsj.18.1055>

BioOne Complete (complete.BioOne.org) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at www.bioone.org/terms-of-use.

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

[REVIEW]

Japanese Eel: A Model for Analysis of Spermatogenesis

Takeshi Miura^{1*} and Chiemi Miura²

¹Marine Bioresources Research Group, Field Science Center for Northern Biosphere,
Hokkaido University, Hakodate 041-8611;

²Division of Marine Biosciences, Graduate School of Fisheries Science,
Hokkaido University, Hakodate 041-8611, Japan

ABSTRACT—The Japanese eel has two characteristics advantageous for the study of the mechanisms controlling spermatogenesis. One is the possibility of artificial induction of the complete process of spermatogenesis from spermatogonial proliferation to spermiogenesis by exogenous gonadotropin injection, and the other is the possibility of inducing this process in an *in vitro* testicular organ culture or germ-Sertoli cell coculture system. Using the eel system, we analyzed the control mechanisms of spermatogenesis. In Japanese eel, the whole process of spermatogenesis is regulated by several sex steroid hormones. Spermatogonial stem cell renewal is promoted by estradiol-17 β (the natural estrogen in vertebrates). Spermatogonial proliferation can be induced by 11-ketotestosterone, the main androgen in teleost. IGF-I is necessary for the action of 11-ketotestosterone in the initiation of spermatogenesis. The action of 11-ketotestosterone is mediated by other factors, such as activin B, produced by Sertoli cells. Although 11-ketotestosterone also induce meiosis and spermiogenesis, the control mechanisms of these processes are not clear. After spermiogenesis, immature spermatozoa undergo sperm maturation, thereby becoming capable of fertilization. Sperm maturation is regulated by 17 α ,20 β -dihydroxy-4-pregnen-3-one (17 α ,20 β -DP), which is progestogen in teleosts. The 17 α ,20 β -DP acts directly on spermatozoa to activate the carbonic anhydrase existed in the spermatozoa. This enzymatic activation causes an increase in the seminal plasma pH, enabling spermatozoa to motile.

Key words: spermatogenesis, Japanese eel, androgen, estrogen, progestin

INTRODUCTION

Spermatogenesis, the formation of sperm that is highly adapted for delivering its genes to an egg, is a complex developmental process. It begins with the mitotic proliferation of spermatogonia, then proceeds through two meiotic divisions followed by spermiogenesis, during which the haploid spermatids develop into spermatozoa. Spermatozoa then undergo maturation, obtaining the ability to fertilize.

Although the process of spermatogenesis is the same in both mammalian and nonmammalian vertebrates, its control mechanisms are not well understood. Spermatogenesis is controlled by numerous hormones and unknown factors (Steinberger, 1971; Hansson *et al.*, 1976; Callard *et al.*, 1978; Billard *et al.*, 1982; Cooke *et al.*, 1998). In higher vertebrates such as mammals, it is difficult to analyze the control mecha-

nisms of spermatogenesis because the seminiferous tubules contain several successive generations of germ cells (i.e., the testicular organization is complex; Clermont, 1972), and few culture systems are available for induction of spermatogenesis *in vitro* (Abe, 1987).

Among species of teleosts, various reproductive styles and gametogenetic patterns. Teleosts constitute the largest phylum (approximately 23,700 species) of living vertebrates (~48,200 species) (Nelson, 1994). Japanese eel, one of such species, has a special spermatogenetic pattern. Under culture conditions, male Japanese eel have immature testes (Fig. 1A) containing only non-proliferated type A and early type B spermatogonia (Miura *et al.*, 1991a). This immature stage of the testis is attributed to insufficient gonadotropin in the eel pituitary (Yamamoto *et al.*, 1972). However, a single injection of human chorionic gonadotropin (hCG) can induce the complete process of spermatogenesis from the proliferation of spermatogonia to spermiogenesis (Fig. 1B,C,D,E) (Miura *et al.*, 1991a). Germ cell development is almost synchronous throughout the testis and the proliferation of spermatogonia,

* Corresponding author: Tel. +81-138-40-5546;
FAX. +81-138-40-5545.
E-mail: takeshi@fish.hokudai.ac.jp




Fig. 1. Light micrographs of eel testis. A) Before hCG injection, and B) 6 days, C) 9 days, D) 12 days, and E) 18 days after hCG injection. Bar, 10 μ m.

meiosis, and spermiogenesis occur at definite times: 3, 12 and 18 days after hCG injection, respectively (Miura *et al.*, 1991a). Furthermore, Japanese eel is the only animal in which complete spermatogenesis has been induced by hormonal treatment *in vitro* using an organ culture system (Fig. 2) and a germ-somatic cells coculture system (Fig. 3), respectively (Miura *et al.*, 1991b,c, 1996). Therefore, the male Japanese eel provides an excellent system for studying the mechanisms of spermatogenesis. This review discusses our recent experimental observations, which indicate the possible molecular control mechanisms of spermatogenesis in Japanese eel.

The endocrine control of eel spermatogenesis

Eel spermatogenesis is also endocrinologically controlled, as is the case in other vertebrates. It is well established that in vertebrates, including fish, gonadotropins (GTHs) are the pri-

mary hormones regulating spermatogenesis (Nagahama, 1987). In most cases, however, it appears that GTHs do not act directly, but rather work through the gonadal biosynthesis of steroid hormones, which in turn mediate various stages of spermatogenesis (Nagahama, 1994).

1) The regulation of spermatogonial stem cell renewal

Spermatogonial mitosis can be categorized by spermatogonial stem cell renewal and spermatogonial proliferation toward meiosis (Clermont, 1972). Recently, it was indicated that estrogen is related to the regulation of the renewal of spermatogonial stem cells in eel (Miura *et al.*, 1999b).

It is widely accepted that "estrogen" is a "female" hormone. However, it has been reported that estrogen exists in some male vertebrates (Schlinger and Arnold, 1992; Fasano and Pieratoni, 1993; Betka and Callard, 1998), and that its receptors are expressed in the male reproductive organs




Fig. 2. The eel testicular organculture system.

(Ciocca and Roig, 1995; Callard and Callard, 1987). Estradiol-17 β (E2), a natural estrogen in vertebrates, was found in Japanese eel serum, and its receptor was expressed in the Sertoli cells (the only non-germinal elements within the seminiferous epithelium of the testes) during the whole process of spermatogenesis. These findings suggested that estrogen is related to the progress of spermatogenesis. We analyzed the action of E2 in spermatogenesis.

Mitosis of eel spermatogonial stem cells was promoted by the implantation of E2, but was suppressed by tamoxifen (an antagonist of estrogen). *In vitro*, 10 pg/ml of E2 was sufficient to induce spermatogonial stem cell division in cultured testicular tissue, confirming the *in vivo* observations. E2 treatments induced only spermatogonial stem cell renewal; they were not found to promote spermatogonial proliferation and meiosis. These findings clearly indicate that estrogen is an indispensable "male hormone", and plays an important role in spermatogonial stem cell renewal.

Generally, E2 induces the target gene expression through its receptor, and the factor translated from this gene affects the biological process. Recently, using gene expression screening, we attempted to clone the cDNA that codes those factors induced or inhibited by E2 stimulation. As a result of this experiment, we obtained three cDNA clones (in preparation). The factor coded by some of these clones may act on spermatogonial renewal.

2) The regulation of spermatogonial proliferation toward meiosis and spermiogenesis

When GTH is secreted from the pituitary, spermatogonial mitosis switches from stem cell renewal to proliferation toward meiosis. As a matter of convenience, we call this point

the initiation of spermatogenesis. It appears that in Japanese eel, GTH does not act directly on germ cells, but rather through the gonadal biosynthesis of 11-ketotestosterone (Miura 1991a,b). 11-ketotestosterone was first identified by Idler *et al.* (1961) as a major androgenic steroid in the male sockeye salmon (*Oncorhynchus nerka*). In various teleost fishes, this steroid has been shown to be synthesized in the testis following GTH stimulation, and high levels were detected in the serum during spermatogenesis (Billard *et al.*, 1982). When 11-ketotestosterone was added to eel testicular organ culture, spermatogenesis from the proliferation of spermatogonia to spermiogenesis was induced (Miura *et al.*, 1991b). The action of 11-ketotestosterone for spermatogenesis is not limited to the Japanese eel; it has been also recognized in goldfish (Kobayashi *et al.*, 1991) and Japanese huchen (Amer *et al.*, 2001). These findings indicate that 11-ketotestosterone is one of the factors involved in the initiation of spermatogonial proliferation toward meiosis.

However, it is believed that the action of 11-ketotestosterone is mediated by other factors produced by Sertoli cells, in which the androgen receptor exists (Ikeuchi *et al.*, in press). It is possible that some of these factors are growth factors, such as insulin-like growth factor-I (IGF-I) and activin B.

IGFs are known to be mediators of growth hormone action in vertebrates. In the rainbow trout testis, IGF-I is expressed in spermatogonia and/or Sertoli cells, and it binds to type 1 IGF receptors. Further, IGF-I stimulates DNA synthesis in spermatogonia (Loir, 1994; Loir and LeGac, 1994; LeGac *et al.*, 1996). Although IGF-I is also necessary for the regulation of eel spermatogenesis, its role is to support the action of 11-ketotestosterone. More specifically, in Japanese




Fig. 3. The eel germ-Sertoli cell coculture system.

eel 11-ketotestosterone is necessary for the induction of spermatogenesis, whereas IGF-I is necessary for the continuation of the process (Nader *et al.*, 1999).

Activin B is a dimeric growth factor belonging to the transforming growth factor-like (TGF β) superfamily, and is composed of two activin β_B subunits. In the Japanese eel, activin B was found in the testis at the initiation of spermatogenesis after hCG stimulation, with its expression site restricted to Sertoli cells. Both transcription and translation of eel activin B were induced by 11-ketotestosterone stimulation *in vitro*. Further, activin B induced proliferation of spermatogonia, but its treatment could not induce meiosis and further spermatogen-

esis (Miura *et al.*, 1995a, b; submitted for publication).

It has been reported that FGF, BMP, PDGF and numerous other growth factors regulate the early stage of spermatogenesis in teleosts and mammals (Watanabe and Onitake, 1995; Zhao *et al.*, 1996; Li *et al.*, 1997; Kim and Fazleabas, 1998). Further investigation is needed for a better understanding of the relationship between growth factors and spermatogenesis in eel.

3) The entry of spermatogonia into meiosis

Following mitotic proliferation, type B spermatogonia differentiate into primary spermatocytes. Generally, the number of mitotic divisions of spermatogonia preceding meiosis are

Table 1. List of cDNA clones obtained by gene expression screening and their expressional patterns.

eSRS No.	homologous protein	size (kb)	up- or down-regulated during hCG induced spermatogenesis	up- or down regulated by 11-KT stimulation <i>in vitro</i>	eSRS No.	homologous protein	size (kb)	up- or down-regulated during hCG induced spermatogenesis	up- or down regulated by 11-KT stimulation <i>in vitro</i>
1	actin β_B	3.4	up	up	15	aquaporin	1.4	up	up
2	unknown	3.4	up	up	16	HMG2	1.1	up	up
3	ZP2	1.4	down	down	17	histone H2A	1.0	up	up
4	ZP3	1.6	down	down	18	tublin α	1.6	up	–
5	cathepsin S	0.85	down	down	19	tublin β	1.5	up	up
6	unknown	1.5	down	–	20	PLK1	3.0	up	up
7	unknown	1.3	down	down	21	TGF β family	2.4	down	down
8	unknown	1.4	down	–	22	carbonic anhydrase	1.5	up	up
9	PCNA	1.6	up	–	23	thymidylate synthase	1.2	up	–
10	unknown	0.5	up	up	24	fatty acid binding protein	1.0	up	up
11	histone H1	0.5	up	up	25	prothymosin α	1.3	up	–
12	CKS1	1.4	up	–	26	ictacalcin	0.8	up	–
13	unknown	–	up	–	27	cytochrome C	0.8	up	–
14	unknown	–	up	–	28	calnexin	2.3	up	–

species specific. In teleosts, a spermatogonial stem cell of medaka (*Oryzias latipes*) yields spermatocytes following 8 mitotic divisions; more specifically, 6 in Sakhalin taimen (*Hucho perryi*), 8 in masu salmon (*Oncorhynchus masou*), 6 in white spotted char (*Salvelinus leucomaenis*), 8 in goldfish (*Carassius auratus*) (Ando *et al.*, 2000), 14 in guppy (*Poecilia reticulata*) (Billard, 1986), 5 or 6 in zebrafish (*Danio rerio*) (Ewing, 1972), and there are 10 mitotic divisions in Japanese eel (Miura *et al.*, 1991a). Although the regulatory mechanisms of the initiation of meiosis are not yet clear, it has been shown that in Japanese eel there is a regulatory stage around the fifth mitotic division of spermatogonia prior to the cells entering meiosis (Miura *et al.*, 1997). To cross this regulatory stage, some factors regulated by 11-ketotestosterone may be required. The key genes coding factors that showed unique expression during spermatogenesis have been considered. To identify these key genes, we isolated cDNA clones of stage-specific genes during eel spermatogenesis using cDNA subtraction and differential display methods (Miura *et al.*, 1998, 1999a). As a result of these experiments, 28 independent cDNA clones showing unique expression patterns during spermatogenesis were obtained (Table 1). As a matter of convenience, we named these clones "eel spermatogenesis related substances (eSRSs)" cDNA. Among these eSRSs, 16 clones are up- or down-regulated by 11-ketotestosterone, the spermatogenesis inducing hormone. The initiation of meiosis may be regulated by some of these factors.

4) Determining whether a control mechanism exists for spermiogenesis

After two meiotic divisions, the germ cells develop into spermatids having small, round, and heterogeneous nuclei. The spermatids transform into spermatozoa through spermiogenesis. This process is characterized by remarkable morphological changes associated with the formation of a spermhead with condensed nucleus, a mid-piece, and a flagellum. Eel spermatozoon has an unusual figure (Fig. 4). It possesses a crescent-shaped nucleus with a flagellum con-

sisting of a 9+0 axonemal structure (generally, the axonemal structure of the flagellum is 9+2), and a single large spherical mitochondrion with developed tubular cristae, which are attached to the caput end at one side of the sperm head (Todd, 1976; Miura *et al.*, 1991a).

In teleosts, it is not yet clear whether regulation mechanisms exist in spermiogenesis. In medaka, it was possible to induce complete spermiogenesis without hormonal treatment in *in vitro* cultured germ cells (Saiki *et al.*, 1997), indicating that there is no regulational mechanism in medaka spermiogenesis. However, the figure of the Japanese eel spermatozoa produced by *in vitro* testicular organ culture and germ cell-Sertoli cell coculture with 11-ketotestosterone is not exactly similar to the fertilizable spermatozoa of eel produced by hCG injection *in vivo*. Based on this discrepancy, it seems conclusive that regulation mechanisms exist in the spermiogenesis of Japanese eel.

5) Induction of sperm maturation

In some species, once the spermatozoa in the testis have completed spermiogenesis, they are not yet capable of fertilizing eggs. In salmonids, the spermatozoa in the testis and in the sperm duct are immotile. If spermatozoa from the sperm duct are diluted with fresh water, they become motile; in contrast, the testicular spermatozoa remain immotile after dilution with fresh water. Thus, spermatozoa acquire the ability of motility during their passage through the sperm duct.

Sperm maturation, the phase during which non-functional gametes develop into mature spermatozoa (fully capable of vigorous motility and fertilization) involves only physiological, not morphological, changes. In salmonids, sperm maturation (the acquisition of sperm motility) has been induced by increasing the seminal plasma pH (approximately to pH 8.0) in the sperm duct, which results in elevation of intrasperm cAMP levels (Morisawa and Morisawa, 1988; Miura *et al.*, 1992). Similar results have been reported for Japanese eel spermatozoa by Miura *et al.* (1995c) and Ohta *et al.* (1997).

Sperm maturation is also regulated by the endocrine sys-




Fig. 4. Electron micrographs of eel spermatozoon. A, The whole image by scanning micrograph (Bar, 1 μ m); B, sagittal section of spermatozoon (Bar, 1 μ m); C, cross section of flagella (Bar, 0.1 μ m); and D, cross sections of sperm heads. Arrowheads indicate the 9 sets of microtubules (Bar, 1 μ m).

tem. In some teleosts including Japanese eel, it is suggested that $17\alpha,20\beta$ -dihydroxy-4-pregnen-3-one ($17\alpha,20\beta$ -DP) is related to the regulation of sperm maturation (Miura *et al.*, 1991d, e, 1992.). $17\alpha,20\beta$ -DP has also been identified as the maturation-inducing hormone of salmonid oocyte (Nagahama and Adachi, 1985). $17\alpha,20\beta$ -DP does not act directly on the sperm; its action is mediated through an increase in the seminal plasma pH, which in turn increases the sperm content of cAMP, thereby allowing the acquisition of sperm motility (Miura *et al.*, 1991d, 1992, 1995c). However, the mechanisms involved in the increase of the seminal plasma pH by $17\alpha,20\beta$ -DP remain unclear. Recently, we attempted to elucidate these mechanisms. As mentioned above, although eSRS22 is one of the factors cloned by the testicular cDNA subtraction method, this factor is related to regulation of the increase in pH (in preparation). In the Japanese eel, eSRS22 is a homo-

logue of carbonic anhydrase (CA). CA catalyzes the reversible hydration of carbon, and is involved in the regulation of ion and acid-base balance in various fluids and tissues (Carter, 1972; Maren, 1967). In Japanese eel, eSRS22/CA protein was expressed in the spermatids and spermatozoa membranes.

In some teleosts including eel, it is suggested that a progesterone receptor exists in the spermatozoon (Gosh and Thomas, 1995; Thomas *et al.*, 1997). If eSRS22/CA is related to sperm maturation, its function seems to be correlated with $17\alpha,20\beta$ -DP and eSRS22/CA activity. Therefore, the relationship between $17\alpha,20\beta$ -DP and eSRS22/CA was examined using intratesticular sperm incubation *in vitro*. The pH value of the artificial seminal plasma (ASP) was increased by $17\alpha,20\beta$ -DP treatment. Moreover, acetazoleamide, a specific inhibitor of CA or anti-eSRS22 specific antibody, suppressed the increase in pH value induced by $17\alpha,20\beta$ -DP stimulation




Fig. 5. A schematic summarizing the possible control mechanisms of spermatogenesis in Japanese eel.

(in preparation). These findings suggested the following possible mechanisms involved in sperm maturation. $17\alpha,20\beta$ -DP acts directly on spermatozoa and induces the activation of eSRS22/CA; this enzymatic activation causes an increase in the seminal plasma pH, and spermatozoa subsequently acquire the motile ability. In masu salmon (*Oncorhynchus masou*), $17\alpha,20\beta$ -DP stimulates the CA activity in spermatozoa, and causes an increase in the pH value of the ASP *in vitro*.

Conclusion

Fig. 5 illustrates the spermatogenic cycle and its predicted regulatory mechanisms in the Japanese eel. By establishment of testicular organ culture and use of molecular biology techniques, analysis of the control mechanisms of eel spermatogenesis has advanced remarkably, and the eel system has proven to be an advantageous system for the study of spermatogenesis. Investigation of the Japanese eel has led to the discovery of several interesting aspects of spermatogenesis. It is highly possible that further investigations of eel spermatogenesis will lead to a better understanding of the general aspects of spermatogenesis.

Recently, environmental pollution by chemicals (collectively known as endocrine disrupters) has been shown to stimulate or block various biological processes (Colborn *et al.*, 1993), and to interfere with the sensitive hormonal pathways that

regulate the reproductive functions. Especially in male animals, exposure to estrogenic compounds (environmental estrogens or exestrogens) can lead to reduced gonad size, feminization of genetic males, and low sperm count and/or quality (Sharpe, 1993; Sumpter, 1995). Japanese eel will also provide an excellent system for analysis of the negative effects of environmental disrupters on spermatogenesis.

ACKNOWLEDGMENTS

A portion of this work was carried out in collaboration with Profs. Yoshitaka Nagahama and Kohei Yamauchi, and we thank them for their encouragement to continue this line of research.

REFERENCES

- Abe S-I (1987) Differentiation of spermatogenic cells from vertebrates *in vitro*. *Internat Rev Cytol* 109: 159–209
- Amer MA, Miura T, Miura C, Yamauchi K (2001) Involvement of sex steroid hormones in the early stages of spermatogenesis in Japanese huchen, *Hucho perryi*. *Biol Reprod* 65: 1057–1066
- Ando N, Miura T, Nader MR, Miura C, Yamauchi K (2000) A method for estimating the number of mitotic divisions in fish testes. *Fisheries Sci* 66: 299–303
- Betka M, Callard GV (1998) Negative feedback control of the spermatogenic progression by testicular oestrogen synthesis: insights from the shark testis model. *APMIS* 106: 252–258

- Billard R, Fostier A, Weil C, Breton B (1982) Endocrine control of spermatogenesis in teleost fish. *Can J Fish Aquat Sci* 39: 65–79
- Billard R (1986) Spermatogenesis and spermatology of some teleost fish species. *Reprod Nutr Develop* 26: 877–920
- Callard I, Callard G, Lance V, Bolaffi J, Rosset J (1978) Testicular regulation in nonmammalian vertebrates. *Biol Reprod* 18: 16–43
- Callard IP, Callard GV (1987) Sex steroid receptors and non-receptor binding proteins. In "Hormones and Reproduction in Fishes, Amphibians and Reptiles" Ed by DO Norris, RE Jones, Plenum Press, New York, pp 355–384
- Carter MJ (1972) Carbonic anhydrase: isoenzymes, properties, distribution, and functional significance. *Biol Rev* 47: 465–513
- Ciocca DR, Roig LMV (1995) Estrogen receptors in human nontarget tissue: Biochemical and clinical implications. *Endocrine Rev* 16: 35–62
- Clermont Y (1972) Kinetics of spermatogenesis in mammals. Seminiferous epithelium cycle and spermatogonial renewal. *Int Rev Cytol* 70: 27–100
- Colborn T, Saal FS, Soto AM (1993) Developmental effects of endocrine-disrupting chemicals in wildlife and humans. *Environ Health Perspect* 101: 378–384
- Cooke HJ, Hargreave T, Elliott DJ (1998) Understanding the genes involved in spermatogenesis: a progress report. *Fertil Steril* 69: 989–995
- Ewing H (1972) Spermatogenesis in zebrafish *Brachydanio rerio* (Hamilton-Buchanan). *Anat Rec* 172: 308
- Fasano S, Pieratoni R (1993). The vertebrate testis. Communication between interstitial and germinal compartments. In "Cellular Communication in Reproduction" Ed by F Facchinetti, IW Henderson, R Pierantoni, A Polzonetti-Magni, Journal of Endocrinology Ltd, Bristol, pp 113–124
- Gosh S, Thomas P (1995) Binding characteristics of 20 β -S to Atlantic croaker sperm membrane receptor. In "the Fifth International Symposium on the Reproductive Physiology of the Fish." Ed by FW Goetz, P Thomas, University of Texas, Austin, pp 239–245
- Hansson D, Calandra R, Purvis K, Ritzen M, French F (1976) Hormonal regulation of spermatogenesis. *Vitam Horm* 34: 187–214
- Idler DR, Bitner II, Schmidt PJ (1961) 11-Ketotestosterone: an androgen for sockeye salmon. *Can J Biochem Physiol* 39: 1737–1742
- Ikeuchi T, Todo T, Kobayashi T, Nagahama Y (in press) Two subtypes of androgen and progesterone receptors in fish testes. *Comp Biochem Physiol*
- Kim JJ, Fazleabas AT (1998) Growth Factors. In "Encyclopedia of Reproduction Vol 2" Ed by E Knobil, JD Neill, Academic Press, San Diego, pp 573–583
- Kobayashi M, Aida K, Stacey NE (1991) Induction of testis development by implantation of 11-ketotestosterone in female goldfish. *Zool Sci* 8: 389–393
- LeGac F, Loir M, Bail PYI, Ollitrault M (1996) Insulin-like growth factor (IGF-I) mRNA and IGF-I receptor in trout testis and in isolated spermatogenic and Sertoli cells. *Mol Reprod Dev* 44: 23–35
- Li H, Papadopoulos V, Vidic B, Dym M, Culty M (1997) Regulation of rat testis gonocyte proliferation by platelet-derived growth factor and estradiol: identification of signaling mechanisms involved. *Endocrinology* 138: 1289–1298
- Loir M (1994) *In vitro* approach to the control of spermatogonia proliferation in the trout. *Mol Cell Endocrinol* 102: 141–150
- Loir M, LeGac F (1994) Insulin-like growth factor I and II binding and action on DNA synthesis in rainbow trout spermatogonia and spermatocytes. *Biol Reprod* 51: 1154–1163
- Maren TH (1967) Carbonic anhydrase: chemistry, physiology, and inhibition. *Physiol Rev* 47: 595–781
- Miura T, Yamauchi K, Nagahama Y, Takahashi H (1991a) Induction of spermatogenesis in male Japanese eel, *Anguilla japonica*, by a single injection of human chorionic gonadotropin. *Zool Sci* 8: 63–73
- Miura T, Yamauchi K, Takahashi H, Nagahama Y (1991b) Hormonal induction of all stages of spermatogenesis *in vitro* in the male Japanese eel (*Anguilla japonica*). *Proc Natl Acad Sci USA* 88: 5774–5778
- Miura T, Yamauchi K, Takahashi H, Nagahama Y (1991c) Human chorionic gonadotropin induces all stages of spermatogenesis *in vitro* in the male Japanese eel (*Anguilla japonica*). *Dev Biol* 146: 258–262
- Miura T, Yamauchi K, Takahashi H, Nagahama Y (1991d) Involvement of steroid hormones in gonadotropin-induced testicular maturation in male Japanese eel (*Anguilla japonica*). *Biomed Res* 12: 241–248
- Miura T, Ninzaki M, Hirai H, Yamauchi K (1991e) Induction of final maturation by injection of chum salmon pituitary homogenate in the male Japanese huchen (*Hucho perryi*). *Bull Fac Fish Hokkaido Univ* 42: 16–25
- Miura T, Yamauchi K, Takahashi H, Nagahama Y (1992) The role of hormones in the acquisition of sperm motility in salmonid fish. *J Exp Zool* 261: 359–363
- Miura T, Miura C, Yamauchi K, Nagahama Y (1995a) Human recombinant activin induces proliferation of spermatogonia *in vitro* in the Japanese eel *Anguilla japonica*. *Fisheries Sci* 63: 434–437
- Miura T, Miura C, Yamauchi K, Nagahama Y (1995b) Activin B is a major mediator of hormone-induced spermatogonial proliferation in the Japanese eel. In "the Fifth International Symposium on the Reproductive Physiology of the Fish." Ed by FW Goetz, P Thomas, University of Texas, Austin, pp 284–286
- Miura T, Kasugai T, Nagahama Y, Yamauchi K (1995c) Acquisition of potential for sperm motility *in vitro* in Japanese eel *Anguilla japonica*. *Fisheries Sci* 61: 533–534
- Miura C, Miura T, Yamashita M, Yamauchi K, Nagahama Y (1996) Hormonal induction of all stages of spermatogenesis in germ-somatic cell coculture from immature Japanese eel testis. *Dev Growth Differ* 38: 257–262
- Miura T, Kawamura S, Miura C, Yamauchi K (1997) Impaired spermatogenesis in the Japanese eel, *Anguilla japonica*: Possibility for the existence of factors that regulate entry of germ cells into meiosis. *Dev Growth Differ* 39: 685–691
- Miura T, Kudo N, Miura C, Yamauchi K, Nagahama Y (1998) Two testicular cDNA clones suppressed by gonadotropin stimulation exhibit ZP2- and ZP3- like structures in Japanese eel. *Mol Reprod Dev* 51: 235–242
- Miura C, Miura T, Kudo N, Yamashita M, Yamauchi K (1999a) cDNA cloning of a stage specific gene expressed during HCG-induced spermatogenesis in the Japanese eel. *Dev Growth Differ* 41: 463–471
- Miura T, Miura C, Ohta T, Nader MR, Todo T, Yamauchi K (1999b) Estradiol-17 β stimulates the renewal of spermatogonial stem cells in males. *Biochem Biophys Res Com* 264: 230–234
- Morisawa S, Morisawa M (1988) Induction of potential for sperm motility by bicarbonate and pH in rainbow trout and chum salmon. *J Exp Biol* 136: 13–22
- Nader MR, Miura T, Ando N, Miura C, Yamauchi K (1999) Recombinant human insulin-like growth factor I stimulates all stages of 11-ketotestosterone-induced spermatogenesis in the Japanese eel, *Anguilla japonica*, *in vitro*. *Biol Reprod* 61: 944–947
- Nagahama Y, Adachi S (1985) Identification of maturation-inducing steroid in a teleost, the amago salmon (*Oncorhynchus rhodurus*). *Dev Biol* 109: 428–435
- Nagahama Y (1987) Gonadotropin action on gametogenesis and steroidogenesis in teleost gonads. *Zool Sci* 4: 209–222
- Nagahama Y (1994) Endocrine regulation of gametogenesis in fish. *Int J Dev Biol* 38: 217–229
- Nelson SS (1994). *Fishes of the world*. John Wiley & Sons, New York
- Ohta H, Ikeda K, Izawa T (1997) Increase in concentrations of potassium and bicarbonate ions promote acquisition of motility *in vitro* by Japanese eel spermatozoa. *J Exp Zool* 277: 171–180

- Saiki A, Tamura M, Matsumoto M, Katowgi J, Watanabe A (1997) Establishment of *in vitro* spermatogenesis from spermatocytes in the medaka, *Oizias latipes*. *Dev Growth Differ* 39: 337–344
- Schlinger BA, Arnold AP (1992) Circulating estrogens in a male songbird originate in the brain. *Proc Natl Acad Sci USA* 89: 7650–7653
- Sharpe RM (1993) Declining sperm counts in men-is there an endocrine cause? *J Endocrinol* 136: 357–360
- Steinberger E (1971) Hormonal control of mammalian spermatogenesis. *Physiol Rev* 51: 1–22
- Sumpter JP (1995) Feminized responses in fish to environmental estrogens. *Toxicology Letters* 82-83: 737–742
- Thomas P, Breckenridge-Miller D, Detweiler C (1997) Binding characteristics and regulation of the 17,20 β ,21-trihydroxy-4-pregnen-3-one (20 β -S) receptor on testicular and sperm plasma membranes of spotted seatrout (*Cynoscion nebulosus*). *Fish Physiol Biochem* 17: 109–116
- Todd PR (1976) Ultrastructure of the spermatozoa and spermiogenesis in New Zealand fresh water eels (Anguillidae). *Cell Tissue Res* 171: 221–232
- Watanabe A, Onitake K (1995) Changes in the distribution of fibroblast growth factor in the teleostean testis during spermatogenesis. *J Exp Zool* 272: 475–483
- Yamamoto K, Hiroi O, Hirano T, Morioka T (1972) Artificial maturation of cultivated male Japanese eel by Synahorin injection. *Bull Jpn Soc Sci Fish* 38: 1083–1090
- Zhao GQ, Deng K, Labosky PA, Liaw L, Hogan BLM (1996) The gene encoding bone morphogenetic protein 8B (BMP8B) is required for the initiation and maintenance of spermatogenesis in the mouse. *Genes Dev* 10: 1657–1669

(Received June 11, 2001 / Invited Review)