



## **Effects of Proteasome Inhibitor (lactacystin) and Cysteine Protease Inhibitor (E-64-d) on Processes of Mitosis in *Xenopus* Embryonic Cells**

Authors: Sekiguchi, Tomohiro, Hosoyama, Yoshiyuki, and Miyata, Shohei

Source: Zoological Science, 19(11) : 1251-1255

Published By: Zoological Society of Japan

URL: <https://doi.org/10.2108/zsj.19.1251>

---

BioOne Complete ([complete.BioOne.org](https://complete.BioOne.org)) is a full-text database of 200 subscribed and open-access titles in the biological, ecological, and environmental sciences published by nonprofit societies, associations, museums, institutions, and presses.

Your use of this PDF, the BioOne Complete website, and all posted and associated content indicates your acceptance of BioOne's Terms of Use, available at [www.bioone.org/terms-of-use](https://www.bioone.org/terms-of-use).

Usage of BioOne Complete content is strictly limited to personal, educational, and non - commercial use. Commercial inquiries or rights and permissions requests should be directed to the individual publisher as copyright holder.

---

BioOne sees sustainable scholarly publishing as an inherently collaborative enterprise connecting authors, nonprofit publishers, academic institutions, research libraries, and research funders in the common goal of maximizing access to critical research.

# Effects of Proteasome Inhibitor (lactacystin) and Cysteine Protease Inhibitor (E-64-d) on Processes of Mitosis in *Xenopus* Embryonic Cells

Tomohiro Sekiguchi, Yoshiyuki Hosoyama and Shohei Miyata\*

Laboratory of Biochemistry, College of Humanities and Sciences, Nihon University,  
Sakurajosui, Setagaya-ku, Tokyo 156-8550, Japan

**ABSTRACT**—At least two different protease pathways have been implicated in the degradation that is required to control the eukaryotic cell cycle; these two pathways center on the activities of ubiquitin/proteasome and cysteine protease. The proteasome inhibitors, lactacystin and AcLLnL and the cysteine protease inhibitor E-64-d were tested for their ability to inhibit the cell cycles of *Xenopus* embryos. Lactacystin, AcLLnL and E-64-d all caused the complete arrest of the cell cycle. To define the specific cell cycle processes that were affected by the two inhibitors, we performed a cytological analysis. Inhibition of the cell cycle by lactacystin and E-64-d occurred during prophase and metaphase. The number of cells that arrested in prophase was 1.4-times higher in the E-64-d-treated group than in the control group and the number of arrested cells in the lactacystin-treated group was 1.4-times higher than in the E-64-d-treated group. The number of cells that arrested in metaphase was 3-to-4-times higher in the E-64-d and lactacystin groups than in the control group. These results indicate that both cysteine protease(s) and proteasomes are involved in the prophase and metaphase stages of cell division.

**Key words:** cysteine protease, E-64-d, proteasome, lactacystin, *Xenopus* embryo

## INTRODUCTION

The degradation of cyclins, p53, cyclin-dependent kinase inhibitor and nuclear oncoproteins is required for multiple processes in mitosis and for the onset of DNA replication (Ciechanover, 1994; King *et al.*, 1996; Hoyt, 1997). Regulated degradation by specific protease pathways is critical for the control of various steps in the cell cycle. Protein degradation mainly occurs via the ubiquitin-dependent proteolytic pathway (Glotzer *et al.*, 1991; King *et al.*, 1995; Sudakin *et al.*, 1995). On the other hand, benzyloxy-carbonyl-Leu-Leu-Tyr diazomethyl ketone, which is thought to inhibit calpain, reduces the proliferation of human TE2 and C-33A cells and WI-38 fibroblasts in the S-phase (Mellgren *et al.*, 1994; Zhang *et al.*, 1997; Mellgren, 1997). N-acetyl Leu-Leu-norleucinal (AcLLnL) inhibits the proliferation of fibroblasts, affecting the progress of the cell cycle from the G1 to the S phase (Sherwood *et al.*, 1993; Dietrich *et al.*, 1996). AcLLnL is not only selective for proteasomes, but it inhibits both cathepsin L and calpain (Roch *et al.*, 1994; Tsubuki *et al.*, 1996). (L-3-*trans*-Ethoxycarbonyloxirane-2

Carbonyl)- L-leucine (3-methylbutyl) amide (E-64-d), an inhibitor of cathepsin and calpain, has been reported to arrest the proliferation of carcinoma A431 cells during metaphase (Shoji-Kasai *et al.*, 1988). We previously reported that E-64-c inhibits the cell cycle of embryonic *Xenopus* cells (Ishii *et al.*, 2000). Consequently, the assumption that proteasomes alone control the cell cycle remains to be verified. To analyze the relation of proteasomes and cysteine proteases to the cell cycle, we studied the effects of a more specific proteasome inhibitor, lactacystin, and the cysteine protease inhibitor, E-64-d, on the process of mitosis in *Xenopus* embryonic cells.

## MATERIALS AND METHODS

### Embryos

Eggs of *Xenopus laevis* were obtained by injection of a gonadotropic hormone and de-jellied with 2.5% cysteine. The embryos were allowed to develop to stage 8 at room temperature.

### Cell culture

Animal cap pieces were dissected from stage 8 blastulae. Single cells from the inner surface of the pieces were separated off by directing a gentle stream of calcium- and magnesium-free medium (50 mM phosphate buffer containing 35 mM NaCl and 1 mM KCl, pH 7.0), as described by Godsave and Slack (1989). Two or three cells were transferred into a well of a Terasaki plate containing 10

\* Corresponding author: Tel. +81-3-5317-9735;  
FAX. +81-3-5317-9433.  
E-mail: miyata@chs.nihon-u.ac.jp

$\mu$ l of 2 mg/ml  $\gamma$ -globulin in a simple salt solution (NAM/2) and cultured in the absence or presence of protease inhibitors for 20 hr at 25°C. Lactacystin and E-64-d were purchased from PEPTIDE INSTITUTE, INC. (Osaka), and AcLLnL was purchased from SIGMA-ALDRICH (Missouri).

### Cytological analysis

To observe the chromosomes, the embryo cells were soaked in 0.02% KCl solution for several min and then fixed with a solution of acetic acid – ethanol (1:3) for 5 min. The cells were then placed on a glass slide, air-dried, and stained with 10% Giemsa in PBS for 30 min.

### Preparation of enzyme samples

To assay the proteasome activities of the *Xenopus* embryos *in vitro*, aliquots of enzyme samples were prepared using the method reported by Tokumoto and Ishikawa (1993). Five hundred embryos in the blastula stage were homogenized in 5 ml of 25 mM Tris-HCl buffer (pH 7.9) containing 10 mM of 2-mercaptoethanol and 20% glycerol. The homogenate was then centrifuged at 36,000 g for 30 min at 4°C

### Enzyme assays

Native gel electrophoresis was performed, as described by Mahaffey *et al.* (1993), on a 4.5% resolving gel and a 2.5% stacking gel. Gels were cast in 90 mM Tris-HCl (pH 8.3), 1.6 mM borate, and 0.08 mM EDTA. After electrophoresis, the gels were soaked for 30 min at room temperature in 10 ml of 30 mM Tris-HCl (pH 7.8), 5 mM  $MgCl_2$ , 10 mM KCl, 0.5 mM DTT, 2 mM ATP, and 50  $\mu$ M Suc-Leu-Leu-Val-Tyr-MCA (sLLVY-MCA), with or without a protease inhibitor. The fluorescent gels were transilluminated by a UV light and photographed with a Polaroid camera.

## RESULTS

The effects of E-64-d, AcLLnL and Lactacystin on cell division in isolated cells were examined *in vitro*. Single cells from mid-blastula-stage *Xenopus* embryos were shown to divide in a non-nutritive medium. Under the standard conditions of the present study, most cells divided between 4 and 10 times. The occurrence of cleavage arrest increased with increasing concentrations of E-64-d (Table 1). A concentration of 400  $\mu$  of E-64-d resulted in the arrest of cleavage in 77% of the treated cells, and a concentration of 800  $\mu$ M resulted in the arrest of cleavage in 96% of the treated cells. A concentration of 1200  $\mu$ M of E-64-d completely blocked the cell cycle.

Treatment with 100  $\mu$ M of AcLLnL resulted in cleavage

arrest in 19% of the treated cells; AcLLnL concentration of 200  $\mu$ M, 400  $\mu$ M and 800  $\mu$ M resulted in cleavage arrest in 79%, 91% and 96% of the treated cells, respectively (Table

**Table 2.** Effects of AcLLnL treatment on cells isolated from blastula embryos. Isolated cells from blastulae were cultured in a medium containing AcLLnL. After 20 hr, blastomere cleavage was observed under a microscope.


Concentration of AcLLnL	No. of non-cleavage cell	No. of cleavage cell
Control	3 (4%)	66 (96%)
100 $\mu$ M	11 (19%)	46 (81%)
200 $\mu$ M	55 (79%)	15 (21%)
400 $\mu$ M	74 (91%)	7 (9%)
800 $\mu$ M	98 (96%)	4 (4%)

**Table 3.** Effects of lactacystin treatment on cells isolated from blastula embryos. Isolated cells from blastulae were cultured in a medium containing lactacystin. After 20 hr, blastomere cleavage was observed under a microscope.

Concentration of Lactacystin	No. of non-cleavage cell	No. of cleavage cell
Control	8 (8%)	88 (92%)
10 $\mu$ M	5 (9%)	61 (91%)
50 $\mu$ M	46 (53%)	41 (47%)
100 $\mu$ M	80 (99%)	1 (1%)
200 $\mu$ M	83 (98%)	2 (2%)

**Table 1.** Effects of E-64-d treatment on cells isolated from blastula embryos. Isolated cells from blastulae were cultured in a medium containing E-64-d. After 20 hr, blastomere cleavage was observed under a microscope.

Concentration of E-64-d	No. of non-cleavage cell	No. of cleavage cell
Control	11 (5%)	202 (95%)
200 $\mu$ M	49 (48%)	53 (52%)
400 $\mu$ M	72 (77%)	22 (23%)
800 $\mu$ M	154 (96%)	6 (4%)
1200 $\mu$ M	80 (100%)	0 (0%)




**Fig. 1.** Effects of lactacystin and AcLLnL on proteasomes in embryonic cells at stage 8. Extract samples of embryos at stage 8 were electrophoresed on a native 4.5% gel and the gels were treated with a 50  $\mu$ M sLLVY-MCA solution with or without a protease inhibitor. The fluorescent proteolytic product was visualized by illumination with UV light. Lane 1, 400  $\mu$ M AcLLnL; lane 2, 100  $\mu$ M lactacystin; lane 3, no protease inhibitor. The arrow indicates the proteasome band, which co-migrates with the same protease from the reticulocyte lysate.

2). Control blastomers were treated with an equivalent volume of solution in which the protease inhibitor was dissolved, which affected 4% of the treated cells by arresting the cell cycle. However, 96% of the treated cells developed normally. The effects of lactacystin on the division of these cells were also studied (Table 3). A concentration of 50  $\mu$ M of lactacystin resulted in the inhibition of cleavage in 53% of the individual cells, and a concentration of 100  $\mu$ M of lactacystin inhibited cleavage in 99% of the cells.

We measured the proteasome activity after gel electro-

phoresis (Fig 1). The fluorescent band of lane 3 showed hydrolytic activity by the proteasome in embryonic cells at stage 8 toward sLLVY-MCA in the presence of ATP. The protease activity was inhibited by lactacystin (Lane 2) and AcLLnL (Lane 1). We have already reported that 400  $\mu$ g/ml of E-64 inhibited the activity of calcium-activated protease in embryonic cells, but E-64 did not inhibit the proteasome activity, even at the same concentration (Ishii *et al.*, 2000).

To analyze the effects of protease inhibitors on specific stages of the cell cycle, we observed the behavior of the



**Fig. 2.** Nuclei and chromosomes of cells arrested in prophase and metaphase by E-64-d and lactacystin. Embryonic cells were treated with 800  $\mu$ M E-64-d or 100  $\mu$ M lactacystin for 1 hr, soaked in a hypotonic solution, and stained with Giemsa. 1 and 2, early prophase and metaphase in arrested cells following treatment with E-64-d; 3, metaphase in arrested cells following treatment with E-64-d; 4 and 5, early prophase and metaphase in cells arrested with lactacystin; 6, metaphase in a cell arrested with lactacystin.

**Table 4.** Number of cells arrested in prophase and metaphase in control preparations without protease inhibitors, colchicine, lactacystin or E-64-d. Isolated cells were treated with a control preparation, 0.5 µg/ml colchicine, 800 µM E-64-d and 100 µM lactacystin for 1 hr. Embryonic cells were treated as described in Fig. 2. The number of cell in prophase includes cells in early prophase and metaphase.

Treatment	No. of Prophase	No. of Metaphase	Total number
Control	125 (26%)	6 (1.2%)	474
	129 (36%)	4 (1.1%)	355
	152 (29%)	5 (0.94%)	528
Colchicine	144 (30%)	82 (17%)	481
	80 (30%)	51 (19%)	269
E-64-d	188 (43%)	12 (2.8%)	437
	168 (41%)	12 (2.9%)	410
Lactacystin	302 (50%)	22 (3.6%)	610
	342 (67%)	23 (4.5%)	512
	248 (60%)	16 (3.9%)	411

nuclei and chromosomes in *Xenopus* embryo cells 1 hr after the addition of protease inhibitors. Fig. 2 shows the prophase (Figs. 2–1, 2, 4, 5) and metaphase (Figs. 2–3, 6) stages of cell division. Following treatment with colchicine, 30% of the cells arrested in prophase and about 18% arrested in metaphase (Table 4). About 30% of the total number of control cells arrested in prophase, while about 42% of the E-64-d-treated cells and about 59% of the lactacystin-treated cells arrested in prophase (Table 4). Thus, the number of cells that arrested in prophase was 1.4 times greater in the E-64-d group than in the control group and 1.4 times greater in the lactacystin group than in the E-64-d group. About 1% of the control cells arrested in metaphase, while 3% of the E-64-d-treated cells and 4% of the lactacystin-treated cells arrested in metaphase (Table 4). Thus, E-64-d and lactacystin arrested 3 to 4 times as many cells in metaphase, compared with the controls.

## DISCUSSION

The cell division cycle in eukaryotes contains three major control or transition points: the conversion of quiescent cells to a state of active proliferation, the initiation of DNA synthesis in the growing cell, and the induction of mitosis in cells with newly replicated genomes (Glutzer *et al.*, 1991; King *et al.*, 1995; Sudakin *et al.*, 1995). Proteolysis of cell cycle-related proteins by proteasomes occurs at least twice during mitosis. Proteasomes are activated during prophase and metaphase and are related to both the entry into the mitotic phase and the transition from metaphase to anaphase (Kawahara *et al.*, 1992; Georgi *et al.*, 2002). Mitotic cyclins A and B are synthesized for the entry into mitosis, and kinase p34 cdk2-cyclin (MPF) induces the various ultrastructural changes required for cell division: namely,

the breakdown of the nuclear envelope, chromatin condensation, and the construction of the mitotic spindle (Minshull *et al.*, 1989). If the continued presence of cyclin is required to activate MPF, then the destruction of cyclin A and B during the stages of mitosis would lead to the inactivation of MPF.

To analyze how proteasomes and cysteine proteases are related to the stages of mitosis, we used two protease inhibitors and assayed the G2/M transition by visual examination of the nuclear morphology. Prophase nuclei were distinguished from interphase nuclei by the condensation of DNA to form mitotic chromosomes and the breakdown of the nuclear envelope (Fig. 2). Metaphase was distinguished from the other stages by the appearance of condensed chromosomes (Fig. 2). Inhibition by the cysteine protease inhibitor, E-64-d, and the proteasome inhibitor, lactacystin, increased the number of cells that arrested in prophase and metaphase, compared with the controls. These results showed that proteasomes and cysteine proteases are related to prophase and metaphase.

DNA condensation and the breakdown of the nuclear envelope depend on the degradation of Weel by ubiquitin/proteasome through a Cdc 34-dependent reaction (Michael and Newport 1998). Degradation of cyclin A always precedes that of cyclin B and occurs just after nuclear envelope breakdown; furthermore, non-degradable cyclin A arrests cells with condensed chromosomes (Holloway *et al.*, 1993). Both cyclin A and B are degraded in mitosis by ubiquitin-mediated proteolysis (Krek 1998; Peters 1999). However, when the cyclin A-type destruction box was substituted for the normal one in cyclin B1 or B2, the resulting constructs could not be digested by proteasomes (Klotzbucher *et al.*, 1996). Thus, while ubiquitination is probably necessary for cyclin destruction, it is not solely responsible. Other proteases are likely to be involved in the degradation of cyclin A.

The initiation of anaphase is known to depend on the onset of sister chromatid separation, by the ubiquitination of the anaphase inhibitor; this reaction activates the anaphase-promoting complex, a cell cycle-regulated ubiquitin protein ligase that targets proteins for destruction by the proteasome (Nasmyth *et al.*, 2000). The injection of calpain II during late metaphase promoted the precocious disassembly of the mitotic spindle and the onset of anaphase (Schollmeyer 1988), and E-64 can arrest carcinoma cells during metaphase (Shoji-Kasai *et al.*, 1988). In this study, we showed that cysteine proteases are not only related to the onset of anaphase, but also to the progression of prophase.

## ACKNOWLEDGMENTS

This work was supported in part by a grant from the Ministry Education, Science, Sports and Culture to Promote Advanced Scientific Research.



## REFERENCES

- Ciechanover A (1994) The ubiquitin-proteasome proteolytic pathway. *Cell* 79: 13–21
- Dietrich C, Bartsch T, Schanz F, Oesch F, Wieser RJ (1996) p53-Dependent cell cycle arrest induced by N-acetyl-L-leucyl-L-leucyl-L-norleucinal in platelet-derived growth factor-stimulated human fibroblasts. *Proc Natl Acad Sci USA* 93: 10815–10819
- Georgi AB, Stukenberg PT, Kirschner MW (2002) Timing of events in Mitosis. *Curr Biol* 12: 105–114
- Glutzer M, Murray AW, Kirschner MW (1991) Cyclin is degraded by the ubiquitin pathway. *Nature* 349: 132–138
- Godsave SF, Slack JMW (1989) Clonal analysis of mesoderm induction in *Xenopus laevis*. *Dev Biol* 134: 486–490
- Holloway SL, Glutzer M, King RW, Murray AW (1993) Anaphase is initiated by proteolysis rather than by the inactivation of maturation-promoting factor. *Cell* 73: 1393–1402
- Hoyt MA (1997) Eliminating all obstacles: Regulated proteolysis in the eukaryotic cell cycle. *Cell* 91: 149–151
- Ishii T, Aoki T, Hosoyama Y, Miyata S (2000) *In vivo* and *in vitro* effects of a protease inhibitor (E-64) on cell division of *Xenopus* eggs. *Zool Sci* 17: 1283–1287
- Kawahara H, Sawada H, Yokosawa H (1992) The 26 S proteasome is activated at two points in the ascidian cell cycle. *FEBS Letter* 310: 119–122
- King RW, Peters J-M, Tugendreich S, Rolf M, Hieter P, Kirschner MW (1995) A 20S complex containing CDC27 and CDC16 catalyzed the mitosis-specific conjugation of ubiquitin to cyclin B. *Cell* 81: 279–288
- King RW, Deshaies RJ, Peters J-M, Kirschner MW (1996) How proteolysis drives the cell cycle. *Science* 274: 1652–1659
- Klotzbucher A, Stewart E, Harrison D, Hunt T (1996) The 'destruction box' of cyclin A allows B-type cyclins to be ubiquitinated, but not efficiently destroyed. *EMBO J* 15: 3053–3064
- Krek W (1998) Proteolysis and the G1-S transition: the SCF connection. *Curr Opin Genet Dev* 8: 36–42
- Mahaffey D, Yoo Y, Rechsteiner M (1993) Ubiquitin metabolism in cycling *Xenopus* egg extracts. *J Biol Chem* 268: 21205–21211
- Mellgren RL (1997) Evidence for participation of a calpain-like cysteine protease in cell cycle progression through late G1 phase. *Biochem Biophys Res Commun* 236: 555–558
- Mellgren RL, Shaw E, Mericle MT (1994) Inhibition of growth of human TE2 and C-33A cells by the cell-permeant calpain inhibitor benzyloxycarbonyl-Leu-Leu-Tyr diazomethyl ketone. *Exp Cell Res* 215: 164–171
- Michael WM, Newport J (1998) Coupling of mitosis to the completion of S phase through Cdc34-mediated degradation of Wee1. *Science* 282: 1886–1889
- Minshull J, Pines J, Golsteyn R, Standart N, Mackie S, Colman A, Blow J, Ruslman J, Wu M, Hunt T (1989) The role of cyclin synthesis, modification and destruction in the control of cell division. *J Cell Sci* 12: 77–97
- Nasmyth K, Peters J-M, Uhlmann F (2000) Splitting the chromosome: cutting the ties that bind sister chromatids. *Science* 288: 1379–1385
- Peters J-M (1999) Subunits and substrates of the anaphase-promoting complex. *Exp Cell Res* 248: 339–349
- Rock KL, Gramm C, Rothstein L, Clark K, Stein R, Dick L, Hwang D, Goldberg AL (1994) Inhibitors of proteasome block the degradation of most cell proteins and the generation of peptides presented on MHC class I molecules. *Cell* 78: 761–771
- Schollmeyer JE (1988) Calpain II involvement in mitosis. *Science* 240: 911–913
- Sherwood SW, Kung AL, Roitelman J, Simoni RD, Schimke RT (1993) *In vivo* inhibition of cyclin B degradation and induction of cell-cycle arrest in mammalian cells by the neutral cysteine protease inhibitor N-acetyl-leucylleucyl-norleucinal. *Proc Natl Acad Sci USA* 90: 3353–3357
- Shoji-Kasai Y, Senshu M, Iwashita S, Imahori K (1988) Thiol protease-specific inhibitor E-64 arrests human epidermoid carcinoma A431 cells at mitotic metaphase. *Proc Natl Acad Sci USA* 85: 146–150
- Sudakin V, Ganioth D, Dahan A, Heller H, Hershko J, Luca FC, Ruderman JV, Hershko A (1995) The cyclosome, a large complex containing cyclin-selective ubiquitin ligase activity, targets cyclins for destruction at end of mitosis. *Mol Biol Cell* 6: 185–197
- Tokumoto T, Ishikawa K (1993) A novel "Active" form of proteasomes from *Xenopus laevis* ovary cytosol. *Biochem Biophys Res Commun* 192: 1106–1114
- Tsubuki S, Saito Y, Tomioka M, Ito H, Kawashima S (1996) Differential inhibition of calpain and proteasome activities by peptidyl aldehydes of di-leucine and tri-leucine. *J Biochem* 119: 572–576
- Zhang W, Lu Q, Xie Z-J, Mellgren RL (1997) Inhibition of the growth of WI-38 fibroblasts by benzyloxycarbonyl-Leu-Leu-Tyr diazomethyl ketone: evidence that cleavage of p53 by a calpain-like protease is necessary for G1 to S-phase transition. *Oncogene* 14: 255–263

(Received June 14, 2002 / Accepted August 30, 2002)